



ARTÍCULO DE INVESTIGACIÓN / RESEARCH ARTICLE

# SEXUAL DIMORPHISM IN *Lysapsus bolivianus* (ANURA: HYLIDAE) IN AN AMAZONIAN FLOODPLAIN

## Dimorfismo sexual en *Lysapsus bolivianus* (Anura:Hylidae) en una llanura amazónica

Patrick RIBEIRO SANCHES<sup>1</sup> , Mayara Fabiana MELO FURTADO<sup>2</sup> , Carlos Eduardo COSTA-CAMPOS<sup>1,2\*</sup>

<sup>1</sup>. Programa de Pós-Graduação em Biodiversidade e Biotecnologia (Rede-BIONORTE), Universidade Federal do Amapá. Campus Marco Zero do Equador, 68903-419, Macapá, AP, Brazil, patricksanchs@gmail.com

<sup>2</sup>. Departamento de Ciências Biológicas e da Saúde, Laboratório de Herpetologia, Universidade Federal do Amapá. Campus Marco Zero do Equador, 68903-419, Macapá, AP, Brazil, mayarafabiana@gmail.com

\* For correspondence: dudueducampos@gmail.com

**Received Date:** 19<sup>th</sup> March 2024. **Revised Date:** 02<sup>th</sup> September 2024. **Accepted Date:** 23<sup>th</sup> November 2024.

**Associate Editor:** Xavier Maquinez

**Please cite this article as / Cite as:** Sanches, P. R., Furtado, M. F. M. and Costa-Campos, C. E. (2025). Sexual dimorphism in *Lysapsus Bolivianus* (Anura: Hylidae) in an amazonian floodplain. *Acta Biol. Colomb.*, 30(1), 89-94. <https://doi.org/10.15446/abc.v30n1.112939>

### ABSTRACT

Sexual dimorphism (SD) in morphology may be attributed to a variety of ecological pressures, including sexual selection, differential growth rate, age at maturity, longevity, and niche partitioning. This morphological variation between the sexes can help us better understand the evolution of life histories and fill gaps about species/populations from less-explored regions. We investigated sexual dimorphism in morphometric characters of *Lysapsus bolivianus* from a floodplain environment in the Eastern Amazon based on 273 specimens collected between 2013 and 2019. Females of *L. bolivianus* attain larger body sizes (Snout-to-Vent Length), but males have proportionally larger heads in length than females. Previous studies indicate that SSD in *L. bolivianus* can be linked to differential growth rate and age to maturity between sexes, whereas differences in head morphometry can be explained by differences in prey volume ingested by males and females. However, the species inhabits a highly seasonal environment and bear pressure predation. In this sense, males and females may cope with different selective pressures due to size differences.

**Keywords:** Eastern Amazon, growth rate, head length, morphometry, shape dimorphism.

### RESUMEN

El dimorfismo sexual (SD) en la morfología puede atribuirse a una variedad de presiones ecológicas, incluida la selección sexual, la tasa de crecimiento diferencial, la edad de madurez, la longevidad y la división de nichos. Esta variación morfológica entre sexos puede ayudarnos a comprender mejor la evolución de las historias de vida y llenar vacíos sobre especies/poblaciones de regiones menos exploradas. Investigamos el dimorfismo sexual en caracteres morfométricos de *Lysapsus bolivianus* de un ambiente de llanura aluvial en el este de la Amazonia con base en 273 especímenes recolectados entre 2013 y 2019. Las hembras de *L. bolivianus* alcanzan tamaños corporales más grandes (longitud del hocico a la ventilación), pero los machos tienen tamaños corporales proporcionalmente mayores. cabezas que las hembras. Estudios anteriores indican que la SSD en *L. bolivianus* puede estar relacionada con la tasa de crecimiento diferencial y la edad hasta la madurez entre sexos, mientras que las diferencias en la morfometría de la cabeza pueden explicarse por diferencias en el volumen de presas ingeridas por machos y hembras. Sin embargo, la especie habita en un entorno altamente estacional y soporta una alta presión de depredación. En este sentido, machos y hembras pueden afrontar diferentes presiones selectivas debido a diferencias de tamaño.

**Palabras clave:** Amazonía oriental, dimorfismo de forma, longitud de la cabeza, morfometría, tasa de crecimiento.

## INTRODUCTION

Sexual size dimorphism (SSD) represents the expressed differences in body morphology related to distinct selective pressures acting on each sex that can be attributed to natural or sexual selection (De Lisle and Rowe, 2013; Rudh and Qvarnström, 2013). It has been documented in a wide range of taxa, including reptiles (Cox *et al.*, 2007), birds (Owens and Hartley, 1998), and mammals (McPherson and Chenoweth, 2012). In anurans, 90 % of SSD is female-biased, i.e. females are larger than males in total length, and it is usually associated with fecundity selection (Shine, 1979; Kupfer, 2007). Besides overall size, differences in body morphology may include other body components such as the size of the head and limbs (Berns, 2013). These traits, often overlooked, can reflect shape differences between sexes which may also be the result of selective mechanisms affecting species individually (Monnet and Cherry, 2002; Arantes *et al.*, 2015). For example, species inhabiting different environments, or possessing distinct ecological adaptations, may exhibit differing degrees of sexual dimorphism in different body parts and traits (Zajitschek *et al.*, 2020).

Aquatic species have evolved unique morphological adaptations to aquatic lifestyle, such as hydrodynamic bodies, elongated pelvic girdles, and relatively long, muscular hind legs with extensive webbing on the hind feet (Citadini *et al.*, 2018). Particular microhabitats may lead to variation in the degree of SSD in anurans. Females with aquatic lifestyles, for instance, are associated with larger egg clutches, which may substantially exacerbate fecundity selection, resulting in a higher degree of SSD (Juarez, 2023). Similarly, ecological pressures of aquatic lifestyle may be important in driving sexual differences in shape and related morphological traits in anurans.

Despite anurans being one of the key targets of SSD investigations, there is still a shortage of empirical data for many taxa. Although external morphology characterization has been extremely valuable for Paradoxical frogs (Hylidae, Pseudoeurycea) systematics (Garda and Cannatella, 2007; Garda *et al.*, 2010), there is still a gap in the extant literature regarding sexual dimorphism for this taxon in less-explored regions. In addition, paradoxical frogs exhibit unique morphological adaptations to cope with their aquatic lifestyle (Turazzini and Gomez, 2023) but very few studies have addressed sex as a factor of intraspecific morphological variation in Pseudoeurycea species (genus *Lysapsus* Cope, 1862 and *Pseudis* Wagler, 1830). Garda *et al.* (2007) reported on sexual dimorphism in *L. limellum*, including specimens of *L. bolivianus* from Amapá state treated as a synonym of *L. limellum*, and found evidence of sexual dimorphism in body size without a significant difference in shape variables between sexes. In this sense, we check here for morphometric differences among males and females of *L. bolivianus* in a floodplain environment at the eastern portion of the Brazilian Amazon. By addressing this topic, we expect to fill gaps


regarding intraspecific phenotypic variation and provide in-depth knowledge about the life-history of this species and its populations.

## MATERIALS AND METHODS

The investigation took place in Rio Curiaú Environmental Protection Area (APA Curiaú), north of Macapá City, Amapá state, Brazil. It covers approximately 40 % of the 21.700 ha of the Curiaú river basin. Vegetation is predominantly composed of lowland marshes and swamps, called várzea forests, with permanent and temporary floodplains and lakes supporting a rich anuran community (Lima *et al.* 2017, Baia *et al.* 2020). We collected specimens at Fazenda Toca da Raposa (00°09'00.7" N, 51°02'18.5" W), one of the floodplain environments of APA Curiaú dominated by herbaceous and shrubs vegetation and surrounded by várzea forest. We surveyed the study site randomly in space by visual and auditory encounter methods (Heyer *et al.*, 1994). We conducted fieldwork once every five months, from February 2013 to October 2019, and each survey lasted 3h (18:00 h-21:00 h). We killed the specimens with a 5 % lidocaine topical solution, fixed in 10 % formaldehyde and preserved in 70 % ethanol to be deposited in the herpetological collection of the Universidade Federal do Amapá (voucher numbers: CECC 538-555, 1042-1057, 2220-2250, 2654-2666, 3352-3371, 3701-3732, 3857-3871, 3929-3957, 4003-4062, 4290-4331).

We measured 273 specimens (114 males and 159 females, see Fig. 1) using a Mitutoyo 500-196-30 digital caliper with 0.01 mm precision and a Carson eFlex MM-840 digital magnifying glass with a millimeter lens for measurements less than 10 mm. We included only adult specimens in the morphometric analysis based on adult snout-to-vent length (Garda *et al.*, 2010). We identified sex by direct observation of the gonads, which we exposed by abdominal dissection, and by secondary sexual characteristics, such as the coloration of the vocal sac in males and the presence of oocytes in females.

We took thirteen morphometric measurements based on Garda *et al.* (2007): snout-vent length (SVL), head length (HL), head width (HW), tympanum diameter (TD), eye diameter (EYDM), interorbital distance (IOD), eye-to-nostril distance (EYNO), snout-to-nostril distance (SND), internarial distance (IND), arm length (AL), forearm length (FAL), thigh length (THL), and tibia length (TL) (Table 1). We applied a cluster and a classification analysis to distinguish between sexes of *L. bolivianus*, respectively: (1) Principal Component Analysis (PCA); and (2) Random Forest (RF) (Breiman, 2001). We conducted these analyses in the R statistical software environment version 4.3 (R Core Team 2018) using SVL and 12 morphometric ratios (HL/SVL, HW/SVL, TD/SVL, EYDM/SVL, IOD/SVL, EYNO/SVL, SND/SVL, IND/SVL, AL/SVL, FAL/SVL, THL/SVL, and TL/SVL). Ratios are generally used to distinguish between morphologically




**Figure 1.** Adult males (a) and females (b) of *Lysapsus bolivianus* collected at Río Curiaú Environmental Protection Area (APA Curiaú), north of Macapá city, Amapá state, Brazil.

similar species which often differ significantly in body proportions, which can be applied to distinguish morphological differences between sexes.

**Table 1.** Summary of morphometric variables measured for *Lysapsus bolivianus* according to sex.

Variables	<i>Lysapsus bolivianus</i>	
	Males=168	Females=122
Snout-vent length (SVL)	14.01 - 20.92 (17.5 ± 1.13)	16.4 - 23.48 (19.7 ± 1.40)
Head length (HL)	3.98 - 8.93 (5.81 ± 0.76)	4.26 - 7.91 (5.86 ± 0.63)
Head width (HW)	4.24 - 7.35 (6.01 ± 0.58)	5.41 - 8.14 (6.86 ± 0.64)
Diameter of tympanum (DT)	0.94 - 2.14 (1.45 ± 0.22)	0.89 - 2.28 (1.55 ± 0.26)
Eye diameter (EYDM)	1.12 - 6.58 (2.27 ± 0.46)	1.71 - 3.22 (2.39 ± 0.26)
Interorbital distance (IOD)	1.13 - 3.8 (2.37 ± 0.75)	1.2 - 3.91 (2.49 ± 0.81)
Eye-to-nostril distance (EYNO)	0.8 - 4.14 (1.52 ± 0.32)	0.74 - 3.02 (1.66 ± 0.30)
Snout-nostril distance (SND)	0.25 - 1.87 (0.96 ± 0.23)	0.43 - 1.54 (1.01 ± 0.20)
Internarial distance (IND)	0.71 - 2.27 (1.17 ± 0.22)	0.74 - 6.95 (1.23 ± 0.54)
Arm length (AL)	1.28 - 8.85 (2.64 ± 0.8)	0.65 - 4.96 (2.77 ± 0.74)
Forearm length (FAL)	1.5 - 4.27 (2.75 ± 0.5)	1.77 - 4.55 (2.96 ± 0.46)
Thigh length (THL)	6.52 - 12.55 (9.76 ± 1.07)	7.47 - 13.41 (10.7 ± 1.12)
Tibia length (TL)	6.98 - 13.3 (10.3 ± 1.07)	7.97 - 13.71 (11.5 ± 1.13)

We performed PCA using the “prcomp” function of the stats package version 4.3.1 using the parameters “scale = T” and “center = T” to scale and center the morphometric variables. We conducted RF with 1000 trees using the “randomForest” function of the randomForest package version 4.7 (Liaw and Wiener, 2002). Random Forest is a machine learning classification algorithm that generates trees from bootstrap samples of a database to aggregate results and classifies samples (specimens) into groups (sex) by choosing the best splits based on a random sample of predictors. To test the hypothesis that the two sexes are statistically distinct in terms of morphometry, we conducted a Multivariate Analysis



**Figure 2.** Cluster analysis based on morphometric variables for males and females *Lysapsus bolivianus*. a) Plot of the first and second multidimensional axis of the Principal Component Analysis, b) charts of variable importance scores of Random Forest and c) confusion matrix showing the classification error of the individuals generated by RandomForest.

of Variance (MANOVA) using the first two axes of the PCA. Additionally, we conducted a univariate approach on SVL and the most important variables (Table 2) indicated by PCA and RF using Analyses of Variance (ANOVA) considering that the analysis exhibited a normal distribution of residuals based on the Shapiro–Wilk test (Shapiro and Wilk, 1965).

**Table 2.** Contribution of each variable to the first two axes of the PCAs in multidimensional morphometric space to explain the variation between males and females of *Lysapsus bolivianus* from APA do Curiaú, Amapá state, Brazil. Bold numbers highlight the variables with the highest contribution values for each Principal Component (PC).

Morphometric measurements	PC1	PC2
Snout-vent length (SVL)	-0.43964	0.052614
Head length ratio (HL/SVL)	0.396395	-0.09906
Head width ratio (HW/SVL)	0.141651	-0.45538
Diameter of tympanum ration (TD/SVL)	0.203381	-0.15397
Eye diameter ratio (EYDM/SVL)	0.257746	-0.21955
Interorbital distance ratio (IOD/SVL)	0.343854	0.416092
Eye-to-nostril distance ratio (EYNO/SVL)	0.195645	0.107765
Snout-nostril distance ratio (SND/SVL)	0.260847	0.053582
Internarial distance ratio (IND/SVL)	0.310285	0.07756
Arm length ratio (AL/SVL)	0.236180	0.354423
Forearm length ratio (FAL/SVL)	0.331883	0.076313
Thigh length ratio (THL/SVL)	0.177411	-0.3811
Tibia length ratio (TL/SVL)	0.059096	-0.48768
Explanation	19.3%	18.4%

## RESULTS

Morphometric multidimensional space occupied by males and females of *L. bolivianus* overlapped substantially for the first two axes of PCA (Fig. 2a). The variables that contributed most to the PC1 were, SVL and HL/SVL. In PC2 the biggest loadings were TL/SVL and HW/SVL (Table 2). The MANOVA test did not indicate difference between sexes for the first two axes of PCA (Pillai trace = 0.27, degrees of freedom (d.f.) = 2 and 270,  $p < 0.0001$ ), with a low Pillai's trace ( $< 0.5$ ) indicating a weak effect of sex on the morphometric variation of individuals.

The capacity to reliably classify morphometrics to the correct sex was moderated, as assessed by the classification error in the confusion matrix of Random Forest (Fig. 2c). In males, 10 % of individuals were misclassified. In comparison, the model exhibited 22 % of incorrect predictions for females. Based on Mean Decrease Accuracy, SVL and HL/SVL were the variables that most contributed to the model's accuracy. Similarly, Mean Decrease Gini indicated that SVL has the largest contribution to the correct split between sexes, while HL/SVL also contributed to the split rule, but to a lesser extent (Fig. 2b).

Our univariate approach confirmed significant differences in morphometric variables analyzed separately, females were significantly larger than males in SVL (ANOVA,  $F_{1,271} = 247.3$ ,  $P < 0.0001$ ; Fig. 3), while males have a proportionally larger head length ratio (HL/SVL) (ANOVA,  $F_{1,271} = 53.8$ ,  $P < 0.0001$ , Fig. 4).


## DISCUSSION

Similar to the findings of Garda et al. (2007) and Sá-Oliveira et al. (2020), our results indicated that females of *L. bolivianus* from different populations in Amapá state are larger than males in SVL. These findings agree with previous


studies of sexual size dimorphism in *L. limellum* in the Brazilian Pantanal (Prado and Uetanabaro, 2000) and *L. laevis* in the Brazilian Amazon (Vaz-Silva et al., 2005). Such a pattern of sexual dimorphism has been the root of the hypothesis involving SSD in anurans and sexual selection (Shine 1979). Woolbright (1983) hypothesized that larger female size is an adaptation to increasing clutch size capacity and the chances of offspring survival. For *L. limellum*, a positive relationship was found between female SVL and the number of deposited eggs (Prado and Uetanabaro, 2000), but the same was not observed by analyzing the number of eggs in ovaries and ovisacs (Garda et al., 2007). Nevertheless, a comprehensive investigation of female fecundity is necessary to confirm such a correlation in *L. bolivianus*.

Alternative explanations for a female-biased SSD are differential growth or age at maturity, which can be explained by a higher male mortality rate or an earlier male maturation (Trivers, 1974; Monnet and Cherry, 2002). *Lysapsus bolivianus* from APA Curiaú is characterized by rapid growth to maturity with males growing faster than females (Sá-Oliveira et al., 2020). However, both sexes have similar longevity and predation is one of the main causes of mortality in this population. Therefore, the differential growth to maturity between sexes associated to mortality can also explain a female-biased SSD observed in *L. bolivianus*. Moreover, the allometric relationships between size and shape can also be influenced by selective pressures acting differently upon each sex (Melo-Moreira et al., 2021).

Some studies report sexual dimorphism in body shape, in which the size of a given morphometric attribute is proportionally different between sexes. For instance, males of *Leptodactylus fuscus* bear broader heads with longer and spatulate-shaped snouts in comparison to females due to their burrowing behavior (Melo-Moreira et al., 2021). Likewise, in the Neotropical Savanna, where males of *Rhinella rubescens*



**Figure 3.** Boxplot representing snout-to-vent length (SVL). In females (F), SVL is greater than in males (M), and values were significantly different.



**Figure 4.** Boxplot representing head length ratio (head length/snout-to-vent length). In males (M), HL ratio is greater than in females (F), and values were significantly different.

were found to have narrower heads, females of *R. schneideri* retained higher heads, likely associated to niche partitioning among sexes as head size is correlated to diet (Arantes *et al.*, 2015). However, the causes of sexual dimorphism in body shape are hard to track, given the impact of several life-history traits on the evolution of sexual dimorphism in anurans (Shine, 1989; Han and Fu, 2013). Contrary to the findings of Garda *et al.* (2007) for *L. limellum*, which found no evidence of shape dimorphism, the data gathered here suggest males may have proportionally larger heads in length. A previous study with the same population of *L. bolivianus* detected difference in prey volume ingested by males and females (Furtado and Costa-Campos, 2020). However, evidence is poor to support the formulation of a definitive conclusion relating diet with sexual dimorphism in head morphology in anurans. In addition, the time and conditions at which aquatic species reach maturity are crucial in determining survival, development, and growth between sexes have the potential to influence dimorphism (Zhang and Lu, 2013). Studies conducted on *L. bolivianus* in the same population reported marked seasonal fluctuations in water levels at the same study site, as well as environmental instability, which can affect sexes differently (Sá-Oliveira *et al.*, 2020).

## AUTHOR'S PARTICIPATION

PRC and CECC collected the data; PRC and CECC wrote the original draft; CECC and MFMF revised the final draft.

## ACKNOWLEDGMENTS

This study is part of the results of the “História Natural de Anfíbios Anuros na Amazônia Oriental”. We are grateful to the colleagues of “Laboratório de Herpetologia” for the assistance on the fieldwork. Instituto Chico Mendes de Conservação da Biodiversidade (ICMBio) for providing collection permits (SISBIO # 34238-1). Mateus Ramos for permission where this study was conducted (Toca da Raposa Farm). Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) for MFMF's scholarship (Proc. 88887.669490/2022-00). CECC thanks the Conselho Nacional de Desenvolvimento Científico e Tecnológico – CNPq for the research grant (Proc. 307697/2022-3).

## CONFLICT OF INTEREST

The authors declare no conflicts of interest.

## REFERENCES

- Arantes, I. D. C., Vasconcellos, M. M., Boas, T. C., Veludo, L. B. A. and Colli, G. R. (2015). Sexual dimorphism, growth, and longevity of two toad species (Anura, Bufonidae) in a Neotropical Savanna. *Copeia*, 103(2), 329-342. <https://doi.org/10.1643/CH-14-092>.
- Baia, R. R., Sanches, P. R., Santos, F. P., Florentino, A. C. and Costa-Campos, C. E. (2020). Diet overlap of three sympatric species of *Leptodactylus* Fitzinger (Anura: Leptodactylidae) in a Protected area in the Brazilian Amazon. *Cuadernos de Herpetología*, 34, 175-184.
- Berns, C. M. (2013). The evolution of sexual dimorphism: understanding mechanisms of sexual shape differences. In: Moriyama, H. (Ed.), *Sexual Dimorphism*. Rijeka: InTech. pp. 1-16.
- Breiman, L. (2001). Random forests. *Machine Learning*, 45, 5-32. <https://doi.org/10.1023/A:1010933404324>.
- Citadini, J. M., Brandt, R., Williams, C. R., and Gomes, F. R. (2018). Evolution of morphology and locomotor performance in anurans: relationships with microhabitat diversification. *Journal of Evolutionary Biology*, 31(3), 371-381. <https://doi.org/10.1111/jeb.13228>.
- Cox, R. M., Butler, M. A. and John-Alder, H. B. (2007). The evolution of sexual size dimorphism in reptiles. In: Fairbairn, D. J., Blanckenhorn, W. U. and Székely, T. (Eds.), *Sex, size and gender roles: evolutionary studies of sexual size dimorphism*. pp. 38-49. Oxford University Press. <https://doi.org/10.1093/acprof:oso/9780199208784.003.0005>.
- De Lisle, S. P. and Rowe, L. (2013). Correlated evolution of allometry and sexual dimorphism across higher taxa. *The American Naturalist*, 182(5), 630-639. <https://doi.org/10.1086/673282>.
- Figueiredo, V. A. M. B., Santos, E. M., Pedroso-Santos, F., Sanches, P. R. and Costa-Campos, C. E. (2022). Diet and clutch size of *Rhinella castaneotica* (Anura: Bufonidae) from a forest area in Serra do Navio, Amapá, Brazil. *North-Western Journal of Zoology*, 18, 35-39.
- Furtado, M. F. M. and Costa-Campos, C. E. (2020). Diet composition of *Lysapsus bolivianus* Gallardo, 1961 (Anura, Hylidae) of the Curiaú Environmental Protection Area in the Amazonas river estuary. *Herpetology Notes*, 13, 113-123.
- Garda, A. A. and Cannatella, D. C. (2007). Phylogeny and biogeography of paradoxical frogs (Anura, Hylidae, Pseudae) inferred from 12S and 16S mitochondrial DNA. *Molecular Phylogenetics and Evolution*, 44, 104-114. <https://doi.org/10.1016/j.ympev.2006.11.028>.
- Garda, A. A., Costa, G. C., França, F. G. R. and Mesquita, D. O. (2007). Ecology of *Lysapsus limellum* in the Brazilian Amazon river basin. *The Herpetological Journal*, 17(3), 141-148.
- Garda, A. A., Santana, D. J. and São-Pedro, V. A. (2010). Taxonomic characterization of Paradoxical frogs (Anura, Hylidae, Pseudae): geographic distribution, external morphology, and morphometry. *Zootaxa*, 2666(1), 1-28. <https://doi.org/10.11646/zootaxa.2666.1.1>.
- Han, X. and Fu, J. (2013). Does life history shape sexual size dimorphism in anurans? A comparative analysis. *BMC Evolutionary Biology*, 13, 1-11. <https://doi.org/10.1186/1471-2148-13-27>.
- Heyer, W. R., Donnelly, M. A., McDiarmid, R. W., Hayek, L. C. E. and Foster, M. S. (1994). *Measuring and monitoring biological diversity. Standard methods for Amphibians*. Smithsonian Institution Press, Washington.

- Juarez, B. H., Moen, D. S. and Adams, D. C. (2023). Ecology, sexual dimorphism, and jumping evolution in anurans. *Journal of Evolutionary Biology*, 36(5), 829-841. <https://doi.org/10.1111/jeb.14171>.
- Kupfer, A. (2007). Sexual size dimorphism in amphibians: an overview. In: Fairbairn, D. J., Blanckenhorn, W. U. and Székely, T. (Eds.), *Sex, size and gender roles: evolutionary studies of sexual size dimorphism* (pp. 50-60). Oxford University Press.
- Liaw, A. and Wiener, M. (2002). Classification and regression by randomForest. *R News*, 2, 18-22.
- Lima, J. R. F., Lima, J. D., Lima, S. D., Silva, R. B. L. and Andrade, G. V. D. (2017). Amphibians found in the Amazonian Savanna of the Rio Curiaú Environmental Protection Area in Amapá, Brazil. *Biota Neotropica*, 17(2), e20160252. <https://doi.org/10.1590/1676-0611-BN-2016-0252>.
- McPherson, F. J. and Chenoweth, P. J. (2012). Mammalian sexual dimorphism. *Animal Reproduction Science*, 131(3-4), 109-122. <https://doi.org/10.1016/j.anireprosci.2012.02.007>.
- Melo-Moreira, D. D. A., Murta-Fonseca, R. A., Galdino, C. A. B. and Nascimento, L. B. (2021). Does a male nest builder have the same head shape as his mate? Sexual dimorphism in *Leptodactylus fuscus* (Anura: Leptodactylidae). *Zoologischer Anzeiger*, 295, 23-33. <https://doi.org/10.1016/j.jcz.2021.09.004>.
- Monnet, J. M. and Cherry, M. I. (2002). Sexual size dimorphism in anurans. *Proceedings of the Royal Society of London*, 269(1507), 2301-2307. <https://doi.org/10.1098/rspb.2002.2170>.
- Owens, I. P. F. Hartley, I. R. (1998). Sexual dimorphism in birds: why are there so many different forms of dimorphism? *Proceedings of the Royal Society of London*, 265, 397-407. <https://doi.org/10.1098/rspb.1998.0308>.
- Prado, C. P. A. and Uetanabaro, M. (2000). Reproductive biology of *Lysapsus limellus* Cope, 1862 (Anura, Pseudidae) in the Pantanal, Brazil. *Zoocriaderos*, 3, 25-30.
- R Core Team (2018). R: A language and environment for statistical computing. R Foundation for statistical computing, Vienna, Austria.
- Rudh, A. and Qvarnström, A. (2013). Adaptive colouration in amphibians. *Seminars in Cell & Developmental Biology*, 24(6-7), 553-561. <https://doi.org/10.1016/j.semcd.2013.05.004>.
- Sá-Oliveira, J. C., Costa-Campos, C. E., Araujo, A. S. and Ferrari, S. F. (2020). Patterns of growth and natural mortality in *Lysapsus bolivianus* (Anura, Hylidae, Pseudae) in an environmental protection area in the estuary of the Amazon River. *Amphibian & Reptile Conservation*, 14, 156-162.
- Shapiro, S. S. and Wilk, M. B. (1965). An analysis of variance test for normality (complete samples). *Biometrika*, 52(3-4), 591-611. <https://doi.org/10.1093/biomet/52.3-4.591>.
- Shine, R. (1979). Sexual selection and sexual dimorphism in the Amphibia. *Copeia*, 1979(2), 297-306. <https://doi.org/10.2307/1443418>.
- Shine, R. (1989). Ecological causes for the evolution of sexual dimorphism: a review of the evidence. *The Quarterly Review of Biology*, 64(4), 419-461. <https://doi.org/10.1086/416458>.
- Trivers, R. L. (1974). Parent-offspring conflict. *Integrative and Comparative Biology*, 14(1), 249-264. <https://doi.org/10.1093/icb/14.1.249>.
- Turazzini, G. F. and Gómez, R. O. (2023). Comparative osteology of paradoxical frogs (Hylidae: Pseudae) with comments on diagnostic features, evolutionary trends and potential aquatic adaptations. *Zoologischer Anzeiger*, 303, 47-70. <https://doi.org/10.1016/j.jcz.2023.01.011>.
- Vaz-Silva, W., Frota, J. G., Prates-Junior, P. H. and Silva, J. S. B. (2005). Dieta de *Lysapsus laevis* Parker, 1935 (Anura: Hylidae) do médio rio Tapajós, Pará, Brasil. *Comunicações do Museu de Ciência e Tecnologia, PUCRS*, 18, 3-12.
- Woolbright, L. L. (1983). Sexual selection and size dimorphism in Anuran Amphibia. *The American Naturalist*, 121(1), 110-119. <https://doi.org/10.1086/284042>.
- Zajitschek, S. R., Zajitschek, F., Bonduriansky, R., Brooks, R. C., Cornwell, W., Falster, D. S., Lagisz, M., Mason, J., Senior, A. M., Noble, D. W. and Nakagawa, S. (2020). Sexual dimorphism in trait variability and its eco-evolutionary and statistical implications. *elife*, 9, e63170. <https://doi.org/10.7554/eLife.63170.sa2>.
- Zhang, L. and Lu, X. (2013). Sexual size dimorphism in anurans: ontogenetic determination revealed by an across-species comparison. *Evolutionary Biology*, 40, 84-91. <https://doi.org/10.1007/s11692-012-9187-2>.