

**Accepted Manuscript / Manuscrito aceptado**

Local and Landscape Drivers of Bird Community Shifts in Urban Parks in a Neotropical City

Doi: <https://doi.org/10.15446/abc.v30n1.113739>

To appear in / Para aparecer en: Acta Biológica Colombiana

Received Date / Fecha de recibido: 27<sup>th</sup> June 2024 / 22 de junio de 2024

Revised Date / Fecha de revisado: 19<sup>th</sup> September 2024 / 19 de septiembre de 2024

Accepted Date / Fecha de aceptado: 04<sup>th</sup> November 2024 / 04 de noviembre de 2024

**Please cite this article as / Cite así:** Almazán-Núñez, R.C., Alvarez-Alvarez, E.A., Poblete-López, D.K., Rodríguez-Godínez, R., Sierra-Morales, P. (2025). Local and Landscape Drivers of Bird Community Shifts in Urban Parks in a Neotropical City. *Acta Biol Colomb.*, 30(1), XX-XX.  
<https://doi.org/10.15446/abc.v30n1.113739>

This is a PDF file of an unedited manuscript accepted for publication. As a service to our readers, we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Este es un archivo PDF de un manuscrito sin editar que ha sido aceptado para su publicación. Como servicio a nuestros lectores, ponemos a su disposición esta versión preliminar del manuscrito. Este manuscrito se someterá a corrección de estilo, composición tipográfica y revisión de la prueba resultante antes de ser publicado en su forma definitiva. Tenga en cuenta que durante el proceso de producción pueden descubrirse errores que podrían afectar al contenido, y que se aplican todas las exenciones de responsabilidad legales que se aplican a la revista.

Acta Biol Colomb (2025); **30(1)**: pag–pag

Doi: <https://doi.org/10.15446/abc.v30n1.113739>

Available online at: <http://www.revistas.unal.edu.co/index.php/actabiol>



Copyright © 2025 Author(s) retain the copyright of this article. This article is published under the terms of the [Attribution-NonCommercial-ShareAlike 4.0](https://creativecommons.org/licenses/by-nc-sa/4.0/) international (CC BY-NC-SA 4.0)

ARTÍCULO DE INVESTIGACIÓN / RESEARCH ARTICLE

## LOCAL AND LANDSCAPE DRIVERS OF BIRD COMMUNITY SHIFTS IN URBAN PARKS IN A NEOTROPICAL CITY

**Impulsores locales y paisajísticos de los cambios en la comunidad de aves en parques  
urbanos de una ciudad neotropical**

**Running head: Local and landscape drives of Neotropical urban bird shifts**

R. Carlos ALMAZÁN-NÚÑEZ<sup>a\*</sup>, Edson A. ALVAREZ-ALVAREZ<sup>b</sup>, Diana K. POBLETE-  
LÓPEZ<sup>c</sup>, Rosalba RODRÍGUEZ-GODÍNEZ<sup>d</sup>, Pablo SIERRA-MORALES<sup>e</sup>

Laboratorio de Ecología y Biogeografía de la Conservación, Facultad de Ciencias Químico-  
Biológicas, Universidad Autónoma de Guerrero, Av. Lázaro Cárdenas s/n, Ciudad Universitaria  
Sur, 39090, Chilpancingo de los Bravo, Guerrero, México, [alvarez.ea@outlook.com](mailto:alvarez.ea@outlook.com);  
[dianapoblete23@hotmail.com](mailto:dianapoblete23@hotmail.com); [goretti.merced3@gmail.com](mailto:goretti.merced3@gmail.com); [sierra02pix@hotmail.com](mailto:sierra02pix@hotmail.com).

<sup>a</sup> ORCID: <https://orcid.org/0000-0002-9913-2737>

<sup>b</sup> ORCID: <https://orcid.org/0000-0002-3771-9946>

<sup>c</sup> ORCID: <https://orcid.org/0000-0003-1183-1646>

<sup>d</sup> ORCID: <https://orcid.org/0000-0001-9326-7414>

<sup>e</sup> ORCID: <https://orcid.org/0000-0002-6232-1278>

\* **For correspondence:** [rcarlos.almazan@gmail.com](mailto:rcarlos.almazan@gmail.com)

**Received:** 27<sup>th</sup> June 2024. **Revised:** 19<sup>th</sup> September 2024. **Accepted:** 04<sup>th</sup> November 2024

**Associate Editor:** Xavier Marquínez

**Citation/ citar este artículo como:** Almazán-Núñez, R.C., Alvarez-Alvarez, E.A., Poblete-López, D.K., Rodríguez-Godínez, R., Sierra-Morales, P. (2025). Local and Landscape Drivers of Bird Community Shifts in Urban Parks in a Neotropical City. *Acta Biol Colomb.*, 30(1), XX-XX. <https://doi.org/10.15446/abc.v30n1.113739>

## ABSTRACT

Urban growth affects bird community composition and diversity. However, these effects may vary based on urbanization level and season. We explored how local and landscape variables influenced urban avoiders, exploiters, and adapters birds during two seasons in urban and peri-urban parks in a neotropical city in southern Mexico. Between 2017 and 2018, we performed bird sampling during non-breeding (August–September) and breeding (February–March) seasons. We compared urban bird diversity between parks using Hill numbers. 67 bird species were recorded—38 adapters, 24 avoiders, and five urban exploiters. Overall, peri-urban parks had the highest bird diversity in both seasons compared to urban parks. Urban exploiters were associated with urbanization at the local and landscape levels in both seasons, while urban avoiders preferred parks with structurally complex vegetation in the non-breeding season. Adapters used urban and peri-urban parks interchangeably. Our findings suggest that densely urbanized parks increase the dominance of exploited bird

species while drastically decreasing the presence of native birds. The findings also highlight the role of peri-urban parks in the conservation and planning of neotropical urban centers, reducing the effects of urbanization.

**Keywords:** bird diversity, greenspaces, non-native birds, urban settlement, vegetation complexity.

### RESUMEN

El crecimiento urbano afecta la composición y diversidad de las comunidades de aves. Sin embargo, estos efectos pueden variar según el nivel de urbanización y la temporada del año. Exploramos cómo variables locales y paisajísticas influyeron en las aves urbanas evasoras, explotadoras y adaptadoras durante dos temporadas en parques urbanos y periurbanos de una ciudad neotropical en el sur de México. Entre 2017 y 2018, realizamos muestreos de aves durante las temporadas no reproductiva (agosto-septiembre) y reproductiva (febrero-marzo). Comparamos la diversidad de aves urbanas entre parques utilizando los números de Hill. Se registraron un total de 67 especies de aves urbanas: 38 adaptadoras, 24 evasoras y cinco explotadoras. En general, los parques periurbanos tuvieron la mayor diversidad de aves en ambas temporadas en comparación con los parques urbanos. Las aves explotadoras urbanas se asociaron con la urbanización a nivel local y paisajístico en ambas temporadas, mientras que las evasoras urbanas prefirieron parques con vegetación estructuralmente compleja en la temporada no reproductiva. Las aves adaptadoras utilizaron parques urbanos y periurbanos indistintamente. Nuestros hallazgos sugieren que los parques densamente urbanizados aumentan la dominancia de especies de aves explotadoras al tiempo que disminuyen drásticamente la presencia de aves nativas. Los hallazgos también destacan el

papel de los parques periurbanos en la conservación y planificación de los centros urbanos neotropicales, reduciendo los efectos de la urbanización.

**Palabras clave:** asentamiento humano, aves no nativas, complejidad de la vegetación, , diversidad de aves.

## INTRODUCTION

Urbanization is a primary driver of the current global biodiversity crisis (Elmqvist et al., 2021). The annual human population growth rate is approximately 1.5 % (7.7 billion people), with an expected increase of 2.3 % by 2100 (10.9 billion people; United Nations, 2019). This exponential growth has led to the modification and loss of natural habitats, resulting in decreased quality of ecosystem services, such as water availability and food production (Seto and Ramankutty, 2016). Furthermore, alterations to natural environments have contributed to a decline in native species and an increase in non-native species, promoting biotic homogenization in urban areas (Clergeau et al., 2006).

Latin America is one of the world's most biodiverse and urbanized regions, with over 80 % of the population residing in metropolises (United Nations, 2019). High population density has significantly impacted Latin American tropical ecosystems, reducing their functions and dynamics (Dobbs et al., 2019). In Mexico, one of the most densely populated countries in the Neotropics, urban development has been both constant and disorderly (United Nations, 2019; INEGI, 2020). Rapid human population growth has also triggered an increase in urban greenspaces, which replace forest areas and affect biodiversity (Puga-Caballero et al., 2014; MacGregor-Fors et al., 2021). However, evidence suggests that urban parks can play a crucial role in supporting bird species diversity if

structural vegetation elements that provide feeding, reproduction, and shelter sites are maintained (Zúñiga-Palacios et al., 2020).

The level of urbanization differentially affects the distribution of urban birds. Urban avoider birds (i.e., species sensitive to human-induced changes) and certain species that have adapted to moderate levels of urbanization may be influenced by urban traits such as noise levels, the number of buildings, human population density, and the presence of non-native bird species (Ortega-Álvarez and MacGregor-Fors, 2009; Carral-Murrieta et al., 2020). Increased urbanization generally leads to decreased species richness. Typically, avoider birds benefit from greenspaces such as recreational parks, vacant lots, and gardens that provide natural microhabitats (Puga-Caballero et al., 2014; Zúñiga-Palacios et al., 2020). Conversely, urbanization mainly favors some urban adapters (i.e., species that tolerate moderately developed urban areas) and exploiter bird species (i.e., species entirely dependent on human activities). An increase in the number of birds in these groups in human settlements has been linked to their ability to exploit anthropogenic resources, such as food and artificial nesting sites, as well as their environmental tolerance to urbanized areas (Fischer et al., 2015). Thus, urban adapter and exploiter populations are increasing yearly, promoting biotic homogenization in urban landscapes (Clergeau et al., 2006).

Bird communities can also be seasonally affected by the degree of urbanization (Zhou and Chu, 2014). During the breeding season (February to June), birds tend to be negatively impacted by buildings that replace vegetation cover, reducing the availability of natural sites for nesting, reproduction, and feeding (Leveau and Leveau, 2016). In contrast, during the non-breeding season, the arrival of migratory birds increases the richness and

abundance of bird species in neotropical urban parks. Urban ecology studies that evaluate the spatial-temporal relationships of bird communities with different urban traits support the development of conservation and planning strategies for biodiversity in urban parks (MacGregor-Fors et al., 2020).

The primary aim of this study was to evaluate the influence of vegetation and urbanization-related local and landscape characteristics on bird communities in urban and peri-urban parks in a city in southern Mexico during both breeding and non-breeding seasons. We hypothesized that bird diversity decreases in urban parks due to higher human disturbance (e.g., people, noise, buildings) and increases in peri-urban areas, which are adjacent to vacant lots and forest areas. However, we also considered that the vegetation structure and landscape features of each park affect bird diversity (Campos-Silva and Piratelli, 2021), regardless of its location. Thus, urban parks with structurally complex vegetation could support greater diversity, even when surrounded by a densely urbanized matrix.

Additionally, due to the variation in resources available in urban and peri-urban parks and the food, reproduction, and shelter requirements of bird species, we predict that the structure and composition of the bird community would differ between urban and peri-urban parks and between breeding and non-breeding seasons. Since urban avoider birds are less tolerant of urbanization and their habitat preferences are positively associated with vegetation complexity, we expect this bird group to be related to greater vegetation complexity at the local level and areas of greater forest coverage at the landscape level. Conversely, we expect urban exploiters and adapters to be more associated with

urbanization variables at both the local and landscape levels, as they have more general habitat requirements and can utilize the urban matrix. The study findings can guide future decisions regarding urban planning and urban park management in cities that are continuously growing but lack orderly urban development practices, a common scenario throughout the Neotropics.

## MATERIALS AND METHODS

### Study area and sampling sites

This study was conducted in Chilpancingo city, the capital of Guerrero state in southern Mexico ( $17^{\circ}33'05''\text{N}$   $-99^{\circ}30'03''\text{W}$ ; Fig. 1a). Chilpancingo has an average elevation of 1 250 m.a.s.l. The predominant climates are semi-warm subhumid and humid with mean annual temperatures ranging from 15 to 24 °C. Predominant vegetation surrounding Chilpancingo is tropical dry forest under different land-use regimes, such as crops and livestock areas. Other vegetation types include oak and pine forests. Chilpancingo is located within the biogeographical province of the Sierra Madre del Sur, one of Mexico's most important biodiversity hotspots (Luna-Vega et al., 2016).

The city has accelerated demographic growth and disorderly urban development—in the last ten years, its population increased by 2.5 %, and it currently has 225,728 inhabitants (INEGI, 2020). This increase places it as one of the cities with the highest population growth rate at the national level (Pisanty et al., 2009). Despite this population increase, it occupies 26<sup>th</sup> position out of the 32 state capitals in the country in terms of population size (INEGI, 2020), so it can still be considered a small to medium-sized city, despite its recent and constant growth.



Six sampling sites were selected within urban greenspaces in Chilpancingo city, separated by an average distance of 1 km to avoid pseudoreplication and ensure data independence. Three of these parks—Margarita Maza (PMM), Alameda (ALA), and Jardín Botánico (JBO)—are located in the city's interior, the most densely urbanized area (hereafter urban parks; Fig. 1b). These parks are surrounded by buildings, residential houses, streets, and avenues that facilitate a greater flow of people and vehicular traffic. Urban parks include management activities such as gardening or tree pruning, as well as human activities like walking, jogging, and picnicking, which peak on weekends. These parks are, on average, 2.1 km from forested areas surrounding the city. The dominant plant species found there include *Bursera linanoe*, *B. submoniliformis*, *B. suntui*, *B. aptera*, *Jacaranda mimosifolia*, *Pithecellobium dulce*, *Casuarina equisetifolia*, *Eucalyptus globulus*, *Tabebuia rosea*, *Ficus* spp., and *Tecoma* spp.

The other three sites are located on the city periphery (hereafter peri-urban parks; Figs. 1a y 1b): Club Deportivo Mitsumaru (CDM), Universidad Sentimientos de la Nación (USN), and Parque Ecológico los Manantiales (PEM). Peri-urban parks have greater vegetation cover and less urban infrastructure. These parks are situated near vacant lots with low levels of human activity. The dominant plant species in these parks include *Guazuma ulmifolia*, *Ipomoea murucoides*, *Jacaranda mimosifolia*, *Pithecellobium dulce*, *Tabebuia* spp., *Ficus* spp., and *Salix* spp. Peri-urban parks are located close to surrounding forest areas (average 0.8 km), allowing for the presence of some native plant species such as *Taxodium mucronatum*, *Cascabela thevetioides*, *Plumeria rubra*, *B. fagaroides*, and *I.*

*murucoides*. The surface area of each urban and peri-urban park was measured using Google Earth® satellite images and geographic information systems (ArcGIS 10.5; Fig. 1b).

### Local and landscape-level variables

We associated the presence of urban bird species with vegetation and urbanization variables at two spatial scales—local and landscape. The local scale refers to the entire park, while the landscape scale refers to a radius of 200 m from the boundaries of each park. At the local level, we measured vegetation structure and composition variables in a 30-m radius plot; at the landscape level, we measured the land-use area covered by each urban and peri-urban park.

In parks smaller than 1.5 ha (PMM, ALA, CDM, and PEM), three circular plots were established, while four circular plots were established in parks larger than 3 ha (JBO and USN). The number of plots varied among parks according to their size. Each plot had a 30 m radius (0.28 ha). To measure vegetation structure and composition variables, two 30 m perpendicular lines were demarcated in each park, oriented toward the four cardinal points. All tree and shrub individuals with a diameter at breast height ( $DBH \geq 15$  cm) whose branches intersected the rope were measured and identified. Vegetation cover was estimated for each tree and shrub using the ellipse formula based on maximum and minimum diameter lengths (Muller-Dombois and Ellenberg, 1974). Foliage stratification was estimated using an optical square marked with two perpendicular axes, allowing the height of objects above to be assessed when viewed horizontally through the device. In each plot, foliage heights and contacts at the point of intersection were recorded 60 times,

at 50 cm intervals along the two axes. The recorded heights were then grouped into 1 m intervals. Foliage height diversity was evaluated using the Shannon-Wiener index. Finally, the number of trees and shrubs per unit area (density) of each plot was estimated. The structure and composition variables obtained in the plots at each park included: 1) plant richness, 2) plant abundance, 3) plant dominance (Simpson index;  $D'$ ), 4) vegetation cover ( $m^{2-1}$ ), 5) diameter at breast height (cm), 6) foliage height diversity ( $H'$ ), 7) mean plant height (m), and 8) plant density ( $ind*ha^{-1}$ ). Additionally, during each bird survey, several urbanization variables in each plot were measured: 1) noise level (expressed in dB), using a sound level digital device to measure noise pollution twice, 5 min apart, in each plot; 2) the number of people per minute, considering only those who crossed within each plot; 3) the number of pets (i.e., dogs and cats) observed per minute; 4) the number of electric wires; and 5) the number of electric poles.

At the landscape level, we delimited buffer areas with a radius of 200 m from the limits of each park (Fig. 1b). This 200-m buffer area was used to assess the effect of urban landscape composition on urban bird communities in each urban and peri-urban park. We considered that this radius would capture the main urban characteristics surrounding the parks that could influence the bird communities. Additionally, the city is small to medium-sized, which limits the use of larger buffer sizes that would cause overlap between them. Other studies have also employed a similar buffer size, demonstrating a favorable response to the effects of urban variables on bird diversity (De Castro et al., 2017). We mapped five land-cover classes: 1) asphalt streets, 2) dirt roads, 3) buildings, 4) bare ground (including areas with herbaceous vegetation), and 5) shrub and tree vegetation (this category included

both native and non-native plants, as they were generally combined in the parks; Fig. 1b). The area of each urban and peri-urban park ranged from 0.5 ha (PEM) to 3.36 ha (JBO; Fig. 1b). Overall, because the city is medium-sized, recreational parks are typically not very large.

### **Urban bird surveys**

Urban bird surveys were conducted over two seasons—non-breeding (August–September 2017) and breeding (February–March 2018). A total of 42 days of fieldwork were spread across 21 days in each season. Three to four sampling points were designated within each park. Bird observations were conducted at intervals of one and a half hours during peak bird activity: mornings (between 07:00 and 08:30 am) and afternoons (between 05:00 and 6:30 pm). Each park received 14 visits (84 visits in total), with seven during each season. The 50-m radius point count method was used to record bird species. This distance was chosen because it generally allowed for good visibility of birds within each park's specific vegetation and urbanization conditions. Point counts were situated in the same circular plots where vegetation and urbanization variables were measured. To ensure data independence, these plots were separated by at least 200 m (Reynolds et al., 1980). Only bird species that actively used the urban habitat (resting, foraging, or nesting) during the surveys were recorded. Species that did not interact with park vegetation or urban structures were excluded from the study. Additionally, bird observations from weekends were omitted due to the potential influence of higher human activity in some urban parks compared to peri-urban areas, which could compromise data comparability and consistency. Each point

received 10 minutes of observation, a timeframe deemed sufficient to count most urban bird species while minimizing the chance of double-counting individuals (Reynolds et al., 1980). Bird identification relied on binoculars ( $8 \times 42$  and  $10 \times 50$ ) and field guides (e.g., Howell and Webb, 1995).

Finally, each recorded bird species was categorized according to its behavior in urban environments. This classification was based on direct field observations and Blair's proposal (1996). The classification system is as follows: 1) urban exploiters, species that can exploit available urban resources (e.g., food, buildings, artificial nests) and are almost entirely dependent on human activities, reaching high population densities in urban environments; 2) urban adapters, species that rely on natural vegetation but can utilize human subsidies in moderately developed urban areas; and 3) urban avoiders, species that disappear with increasing urbanization due to their high sensitivity to human-induced changes (Blair, 1996). The scientific nomenclature and systematic arrangement of urban bird species followed the guidelines of the American Ornithological Society (Chesser et al., 2024).

### **Data analyses**

All analyses, except for the sampling effort evaluation and canonical correspondence analysis, were performed using the *iNEXT*, *MASS*, and *vegan* libraries (Hsieh et al., 2016; Oksanen et al., 2022; Ripley et al., 2022) in R 4.0.3 (R Development Core Team, 2020).

### **Vegetation variables at the local and landscape levels**

Mean  $\pm$  standard error (SE) values for the thirteen vegetation and urbanization variables at the local and landscape levels were obtained.

### **Sampling effort**

The potential number of urban bird species in each urban and peri-urban park was estimated using the Chao 2 richness estimator in EstimateS 9.1 (Colwell, 2013). Chao 2 is reliable for relatively small sampling units (i.e., point counts; 0.25 km<sup>2</sup>) and is less dependent on sampling effort than other estimators (Hortal et al., 2006).

### **Urban bird species richness and diversity**

Urban bird diversity in both urban and peri-urban parks was estimated across two seasons (breeding and non-breeding) using the effective number of species, or Hill numbers (Jost, 2006). Diversity was assessed at three levels: first-order diversity, measured as urban bird species richness ( $q = 0$ ), second-order diversity, or the exponential of Shannon entropy, which accounted for species' proportional abundance ( $q = 1$ ), and third-order diversity, or the inverse Simpson concentration, which reflected the effective number of dominant bird species in the community ( $q = 2$ ; Jost, 2006). We compared all diversity values using 84 % confidence intervals (CI), as they robustly mimic  $P < 0.05$  statistical tests for both symmetric and asymmetric CI (MacGregor-Fors and Payton, 2013). If the 84 % CI for the three diversity values does not overlap, differences between the urban and peri-urban groups are interpreted as significant.

### Structure of urban bird communities

To compare bird community structure (dominance/evenness) between urban and peri-urban parks during the breeding and non-breeding seasons, we employed species rank/abundance plots. Steep curves indicate low evenness and high dominance, while less pronounced curves indicate increasing uniformity and decreasing dominance (Magurran, 2004). We compared the slopes of the rank/abundance plot regression lines using a covariance analysis (ANCOVA) to test whether the proportion of dominant and rare urban bird species varied between urban and peri-urban parks during the two seasons. Urban bird abundance data were log-transformed ( $\log_{10}$ ) as recommended by Magurran (2004).

### Influence of habitat at the local and landscape levels

A principal component analysis (PCA) was performed to synthesize the thirteen vegetation and urbanization variables at the local level through varimax rotation to redistribute the variance of factors. To fulfill the assumptions of normality and homogeneity of data variance, vegetation and urbanization variables at the local and landscape levels were standardized to zero means with unit standard deviation and  $\sqrt{\arcsine(x/100)}$ , respectively. A canonical correspondence analysis (CCA) was then conducted to analyze the relationship between urban bird species and transformed variables at the local and landscape levels during the breeding and non-breeding seasons (Ter Braak and Verdonschot, 1995). The factors obtained in the vegetation and urbanization PCAs were used in the CCAs and plotted as local-level vectors. The five land-cover classes—asphalt

streets, dirt roads, buildings, bare ground, and shrub and tree vegetation—were used in the CCAs as landscape-level vectors. Urban bird species with fewer than three individuals were excluded from the CCAs, as low individual numbers may result from random observations in urban and peri-urban parks. We performed Monte Carlo permutations (999 simulations,  $p < 0.05$ ) to assess the relationship between urban bird species and vegetation and urbanization variables at the local and landscape levels. Two statistical tests were used, one based on the first ordination axis and the other on all canonical axes (Ter Braak and Verdonschot, 1995). These analyses were carried out using CANOCO 4.5 and CANODRAW 4.0 software (Ter Braak and Šmilauer, 2002).

## RESULTS

### Description of the parks at the local and landscape levels

At the local level, JBO urban park had the most diverse tree species ( $10 \pm 3.47$ ; mean  $\pm$  standard error). Plant abundance and density were similar across all parks. PEM and CDM parks had the largest trees and the most varied foliage heights. These two parks, along with JBO, had the most vegetation cover. PMM and ALA parks exhibited the most noise, pets, people, electric poles, and wires.

At the landscape level, PMM and JBO parks had the most asphalt streets and buildings, respectively. USN and CDM parks had the barest ground and vegetation cover, respectively. USN also had the most dirt roads (Table S1 for details).

During the breeding season, vegetation and urbanization variables were synthesized into three principal components (PCs) that explained 89 % of the variance at the local level.



PC1 represented urbanization variables (52.7 % of the explained variance). PC2 and PC3 explained 20.8 % and 15 % of the variance (73.6 % and 89 % of the cumulative variance, respectively), representing vegetation composition and structure variables. In the non-breeding season, the three PCs explained 88 % of the variance. PC1 accounted for 56.3 % of the variance, grouping urbanization variables, while PC2 and PC3 explained 18.7 % and 12.7 % of the variance (75.1 % and 88 % of the cumulative variance, respectively). Both components represented vegetation composition and structure variables (Table S2).

### **Urban bird composition, richness, and diversity**

We recorded 67 urban bird species belonging to 23 families across the six parks (Table S3). The peri-urban and JBO urban parks exhibited the highest urban bird species richness (Table 1). Based on the Chao 2 estimator, 73 %–92 % of the expected urban bird species were recorded in the urban and peri-urban parks. Further, 38 urban bird species were classified as urban adapters, 24 as avoiders, and five as exploiters. The number of urban exploiters increased slightly in the peri-urban parks. The peri-urban and JBO urban parks had the highest number of urban adapters and avoiders (Table 1). The USN and CDM peri-urban parks had the highest urban bird species richness ( $q_0$ ) and diversity ( $q_1$  and  $q_2$ ) during both breeding and non-breeding seasons (Fig. 2).

### **Structure of urban bird communities**

Bird community structure significantly differed between urban and peri-urban parks during the breeding and non-breeding seasons (ANCOVA: both seasons  $p < 0.05$ ; Fig. 3).

The House Sparrow (*Passer domesticus*) was dominant in two urban parks (PMM and ALA) in both seasons and during the breeding season in the PEM peri-urban park. The Great-tailed Grackle (*Quiscalus mexicanus*) was dominant in the JBO urban park and the CDM peri-urban park during both seasons, as well as in the PEM peri-urban park during the non-breeding season. While *Turdus rufopalliatus* was dominant in USN during the breeding season, *Sporophila torqueola* was dominant in this peri-urban park during non-breeding seasons (Fig. 3).

#### **Distribution of urban bird community at the local and landscape levels**

At the local level, the first two axes of the CCA accounted for 82 % (axis 1 = 51 %, axis 2 = 82 %) and 94 % of the cumulative variance (axis 1 = 66 %, axis 2 = 94 %) for breeding and non-breeding seasons, respectively. Only the first axis was significant for both breeding (Monte Carlo: F-ratio = 1.57,  $p = 0.02$ ) and non-breeding seasons (Monte Carlo: F-ratio = 1.61,  $p = 0.01$ ). Urbanization variables positively influenced urban exploiters, while some adapters and avoiders were positively associated with vegetation structure and composition during both seasons (Fig. 4a–b).

At the landscape level, two main axes of the CCA explained 68 % (axis 1 = 43 %, axis 2 = 68 %) and 82 % of the cumulative variance (axis 1 = 53 %, axis 2 = 82 %) for breeding and non-breeding seasons, respectively. The Monte Carlo tests for all canonical axes were significant for both breeding (F-ratio = 1.53,  $p = 0.04$ ) and non-breeding seasons (F-ratio = 1.72,  $p = 0.01$ ). Most urban avoider species were positively associated with vegetation cover, while urban exploiters were primarily related to buildings, and adapter

species were associated with vegetation and urbanization covers throughout the survey periods (Figs. 4c y 4d).

## DISCUSSION

### Bird diversity among urban parks

As hypothesized, bird species diversity decreased in parks with the greatest influence of urbanization during both breeding and non-breeding seasons. This lower species diversity in urban parks can be attributed to the scarcity of resources and inadequate conditions for the survival of native birds that are intolerant of urbanization (Blair, 2004). Another contributing factor is that two urban parks (ALA and PMM) are public spaces adjacent to schools and commercial centers, characterized by high levels of human presence, buildings, and noise pollution. Previous studies have shown that these factors associated with densely urbanized areas significantly decrease bird species diversity (Carral-Murrieta et al., 2020; Escobar-Ibáñez et al., 2020; Schrimpf et al., 2021). Therefore, parks situated within a densely urbanized matrix pose threats to sensitive bird species due to anthropogenic disturbances (Amaya-Espinel et al., 2019).

However, we also found that vegetation structure, regardless of the park's urban or peri-urban location, appears to be a critical factor in enhancing species diversity. For instance, the JBO urban park exhibited species diversity comparable to that obtained in the peri-urban parks. This urban park, while surrounded by an urban matrix, has greater vegetation cover and plant richness, positively influencing the presence of certain bird species typical of less urbanized areas (e.g., Northern Beardless-Tyrannulet [*Camptostoma*

*imberbe*], White-tipped Dove [*Leptotila verreauxi*], and Summer Tanager [*Piranga rubra*]).

This finding aligns with other studies highlighting the importance of local conditions for maintaining bird species diversity in neotropical urban greenspaces (Puga-Caballero et al., 2014; Zúñiga-Palacios et al., 2020). Notably, the JBO park, located within the Guerrero Autonomous University, represents a space characterized by the native flora of southern Mexico. Even though this park is situated within the densely urbanized matrix, the native flora promotes the visitation and permanence of bird species seeking food, shelter, and breeding sites (Castro-Torreblanca and Blancas-Calva, 2014). Latin American university campuses are considered potential refuges for biodiversity, as they maintain greenspaces with native woody plants that provide resources for birds within densely urbanized matrices (Lessi et al., 2016). In the JBO, some native plant species, such as *Bursera linanoe*, *B. submoniliformis*, and *B. aptera*, provide fruits for the Tyrannidae, Turdidae, and Vireonidae families. It has been shown that fruits of *Bursera* spp. contain lipids that constitute a valuable nutrient source for both resident and migratory bird species (Almazán-Núñez et al., 2016).

In contrast to the patterns observed in most urban parks, peri-urban parks exhibited the most diverse bird species diversity during the breeding and non-breeding seasons. As previously documented, this is due to greater vegetation cover and lower human density compared to urban parks (Canedoli et al., 2018). Additionally, these parks, located peripherally in Chilpancingo city, are closer to forest areas, allowing birds can move toward these peri-urban parks (Puga-Caballero et al., 2014). Some bird groups typical of forest areas, such as the urban avoiders White-tipped Dove, Rusty-crowned Ground-

Sparrow (*Melospiza kieneri*), and Banded Wren (*Thryophilus pleurostictus*), were primarily observed in peri-urban parks. This highlights the biological and ecological importance of these parks as alternative refuges and biological corridors for avian maintenance and conservation in highly urbanized areas (Lessi et al., 2016; Canedoli et al., 2018).

### **Urban bird community structure among urban parks**

Bird community structure varied significantly between parks during both breeding and non-breeding seasons. These differences are primarily related to the presence of urban exploiter birds, such as the House Sparrow and Great-tailed Grackle. Both species dominated in nearly all parks during both seasons, demonstrating their adaptability to different anthropogenic environments (García-Arroyo et al., 2020). The House Sparrow and Great-tailed Grackle are generalist species associated with human disturbance, utilizing building facades and roofs as resting and nesting sites (MacGregor-Fors and Schondube, 2011; Faggi and Caula, 2017). Additionally, in urban parks, human activities indirectly provide these birds with food and other waste necessary for their survival, increasing their dominance. Conversely, because food and nesting resources derived from natural vegetation are more abundant in peri-urban parks, and human activity is considerably reduced, bird communities tend to be more equitably distributed in these areas (Puga-Caballero et al., 2014; Lessi et al., 2016).

Although House Sparrows and Great-tailed Grackles were abundant in two peri-urban parks (PEM and CDM), their dominance was not as pronounced as in urban parks. This indicates that local conditions within these parks and the matrix of adjacent habitats

are not entirely favorable for the settlement of these species (MacGregor-Fors and Schöndube, 2011). However, as the populations of these two urban exploiters increase, native birds become less adapted to urbanization, leading to their replacement (García-Arroyo et al., 2020) and avian homogenization in the urban parks of Chilpancingo City, as documented globally (Clergeau et al., 2006).

### **Distribution of urban bird community at the local and landscape levels**

As expected, urban avoider birds were positively related to vegetation complexity, while exploiters and some urban adapters were primarily associated with urbanization at both the local and landscape levels. The biological traits, life histories, and habitat preferences of each urban species drive these associations between urban bird groups and different environmental characteristics. For example, urban exploiters and adapters are generalist species. These species tend to be larger with broad diets, increasing their probability of adaptation and survival across a wide range of environmental conditions (Palacio, 2020). Our results indicated that these species are predominantly distributed in densely urbanized parks. Most urban exploiters, such as the House Sparrow, Rock Pigeon (*Columba livia*), and Eurasian Collared-Dove (*Streptopelia decaocto*), are invasive birds, while others, such as the Great-tailed Grackle, are native but exhibit invasive behavior. This finding supports the urban tolerance hypothesis, which states that urbanization favors species based on their biological traits, enabling them to utilize resources and adapt to the adverse conditions posed by urban landscapes (Sol et al., 2014).

Urban exploiters and several urban adapters are both granivorous scavenger species, with a vast food spectrum (i.e., euryphagic species; Møller, 2014). Consequently, such species can successfully use anthropogenic resources in urban areas (Blair, 1996; Fischer et al., 2015), explaining the positive association of these species with urbanization. Particularly, urban exploiters were favored by urban variables at the local (number of people, electric poles, and wires) and landscape level (buildings and bare ground), which is explained by their remarkable physiological plasticity and behavior. This plasticity allows urban exploiters to use any resource (e.g., bare ground, buildings, electric poles, and wires) to carry out different functions in their life cycle in densely urbanized matrices (Palacio, 2020). It has been shown, for example, that these species use electric poles and wires as perching places and sites for reproduction (Mikami et al., 2022). This pattern coincides with previous studies describing how urban exploiters and some adapter species are primarily associated with urbanization due to their general behavior in urban landscapes (Ortega-Álvarez and MacGregor-Fors, 2009; MacGregor-Fors and Schondube, 2011).

By contrast, greater vegetation complexity favored urban avoiders at both the local and landscape levels. Blue Mockingbird (*Melanotis caerulescens*), Cinnamon-bellied Saltator (*Saltator grandis*), and Golden Vireo (*Vireo hypochryseus*) were associated with parks exhibiting more complex vegetation during both breeding and non-breeding seasons, preferring peripheral sites adjacent to the urban matrix (Blair, 1996; Fischer et al., 2015). This bird group is associated with wooded areas that provide food, nesting sites, shelter, and reproduction opportunities, situated further away from noise, vehicular traffic, buildings, and the continuous flow of people (Polak et al., 2013). Continuous human

activity poses a threat to urban evasive species (Palacio, 2020). Recently, Schrimpf et al. (2021) demonstrated that decreased human mobility due to the COVID-19 pandemic increased the abundance of some bird species compared to pre-pandemic levels in urban habitats in North America. This underscores how accelerated and uncontrolled urban growth, particularly in Latin American countries, could jeopardize the survival of native bird species sensitive to anthropogenic disturbance.

## CONCLUSIONS

This study provides evidence of the effects of urbanization on the structure and composition of bird communities. Parks that are densely urbanized at the local and landscape levels show an increase in the abundance of exploiters (primarily non-native invasive species) and some urban adapters. Conversely, peripheral parks with a greater supply of resources can retain the local character of their native birds, resulting in a better representation of urban avoiders in these sites. Under certain structural vegetation conditions, parks located on the periphery of cities can serve as connectors for birds found in parks situated within the urban matrix and are essential components of conservation and planning in Neotropical urban centers. This is particularly important for the Latin American region, which contains countries with the greatest biodiversity and some of the highest urbanization rates in the world. Moreover, we demonstrate that seasonality affects differentiation in urban bird community structure, even though general trends in bird diversity patterns were maintained across the different parks.



### ACKNOWLEDGMENTS

We thank Isela Salazar and Marisaby Morales for their help in the fieldwork. We also wish to thank Facultad de Ciencias Químico Biológicas of the Universidad Autónoma de Guerrero for the facilities to carry out this research.

### DISCLOSURE OF INTERESTS

The authors declared no potential conflicts of interest concerning the research, authorship, and/or publication of this article.

### AUTHORS CONTRIBUTIONS

RCA-N: Conceptualization, Methodology, Formal analysis, Writing - Original Draft, Supervision, Project administration. EAA-A: Conceptualization, Methodology, Formal analysis, Writing - Review & Editing. DKP-L: Conceptualization, Writing - Review & Editing. RR-G: Conceptualization, Writing - Review & Editing. PS-M: Conceptualization, Writing - Review & Editing.

### REFERENCES

- Almazán-Núñez, R. C., Eguiarte, L. E., Arizmendi, M. C. and Corcuera, P. (2016) *Myiarchus* flycatchers are the primary seed dispersers of *Bursera longipes* in a Mexican dry forest. *PeerJ*, 4, e2126. <https://doi.10.7717/peerj.2126>
- Amaya-Espinel, J. D., Hostetler, M., Henríquez, C. and Bonacic, C. (2019). The influence of building density on Neotropical bird communities found in small urban parks.

*Landscape and Urban Planning*, 190, 103578.

<https://doi.10.1016/j.landurbplan.2019.05.009>

Blair, R. B. (1996). Land use and avian species diversity along an urban gradient.

*Ecological Applications*, 6(2), 506–519. <https://doi.10.2307/2269387>

Blair, R. B. (2004). The effects of urban sprawl on birds at multiple levels of biological organization. *Ecology and Society*, 9(5), 2. <https://doi.10.5751/ES-00688-090502>

Campos-Silva, L. A. and A. J. Piratelli. 2021. Vegetation structure drives taxonomic diversity and functional traits of birds in urban private native forest fragments. *Urban Ecosystems*, 24, 375-390. <https://doi.org/10.1007/s11252-020-01045-8>

Canedoli, C., Manenti, R. and Padoa-Schioppa, E. (2018). Birds biodiversity in urban and periurban forests: environmental determinants at local and landscape scales. *Urban Ecosystems*, 21, 779–793. <https://doi.10.1007/s11252-018-0757-7>

Carral-Murrieta, C. E., García-Arroyo, M., Marín-Gómez, O. H., Sosa-López, J. R. and MacGregor-Fors, I. (2020). Noisy environments: untangling the role of anthropogenic noise on bird species richness in a Neotropical city. *Avian Research*, 11(32). <https://doi.10.1186/s40657-020-00218-5>

Castro-Torreblanca, M. and Blancas-Calva, E. (2014). Aves de Ciudad Universitaria campus Sur de la Universidad Autónoma de Guerrero, Chilpancingo, Guerrero, México. *Huitzil*, 15, 82–92. <https://doi.org/10.28947/hrmo.2014.15.2.57>

Chesser, R. T., Billerman, S. M., Burns, K. J., Cicero, C., Dunn, J. L., Hernández-Baños, B. E. Jiménez, R. A., Kratter, A. W., Mason, N. A., Rasmussen, P. C. and Remsen, J. V.

- (2024). Check-list of North American birds (online). Retrieved on 07 October 2024 from: <https://checklist.americanornithology.org/taxa/>
- Clergeau, P., Croc, S., Jokimäki, J., Kaisanlahti-Jokimäki, M. L. and Dinetti, M. (2006). Avifauna homogenisation by urbanisation: analysis at different European latitudes. *Biological Conservation*, 127(31), 336–344. <https://doi.org/10.1016/j.biocon.2005.06.035>
- Colwell, R. K. (2013). EstimateS: Statistical estimation of species richness and shared species from samples, Version 9.1. Retrieved on 05 February 2023 from <http://viceroy.eeb.uconn.edu/estimates> ).
- De Castro, J. C., Martello, F., Ribeiro, M., Armitage, R. A., Young, R. J. and Rodrigues, M. (2017). Street trees reduce the negative effects of urbanization on birds. *PLoS ONE*, 12, e0174484. <https://doi.org/10.1371/journal.pone.0174484>
- Dobbs, C., Escobedo, F. J., Clerici, N., de la Barrera, F., Eleuterio, A. A., MacGregor-Fors I., Reyes-Paecke, S., Vásquez, A., Camacho, J. D. and Hernández, J. (2019). Urban ecosystem services in Latin America: mismatch between global concepts and regional realities? *Urban Ecosystems*, 22, 173–187. <https://doi.org/10.1007/s11252-018-0805-3>
- Elmqvist, T., Andersson, E., McPhearson, T., Bai, X., Bettencourt, L., Brondizio, E., Colding, J., Daily, G., Folke, C., Grimm, N., Haase, D., Ospina, D., Parnell, S., Polasky, S., Seto, K. C. and Leeuw V. D. (2021). Urbanization in and for the Anthropocene. *npj Urban Sustainability*, 1(6). <https://doi.org/10.1038/s42949-021-00018-w>

- Escobar-Ibáñez, J. F., Rueda-Hernández, R. and MacGregor-Fors, I. (2020). The greener the better! avian communities across a neotropical gradient of urbanization density. *Frontiers in Ecology and Evolution*, 8, 500791. <https://doi.org/10.3389/fevo.2020.500791>
- Faggi, A. and Caula, S. (2017). ‘Green’ or ‘gray’? Infrastructure and bird ecology in urban Latin America. In: MacGregor-Fors, I. and Escobar-Ibáñez J.F. (Eds) Avian ecology in Latin American cityscapes. Pp. 79–98. Springer International Publishing AG, Cham.
- Fischer, J. D., Schneider, S. C., Ahlers, A. A. and Miller, J. R. (2015). Categorizing wildlife responses to urbanization and conservation implications of terminology. *Conservation Biology*, 29(4), 1246–1248. <https://doi.org/10.1111/cobi.12451>
- García-Arroyo, M., Santiago-Alarcón, D., Quesada, J. and MacGregor-Fors, I. (2020). Are invasive House Sparrows a nuisance for native avifauna when scarce? *Urban Ecosystems*, 23, 793–802. <https://doi.org/10.1007/s11252-020-00963-x>
- Hortal, J., Borges, P. A. V. and Gaspar, C. (2006). Evaluating the performance of species richness estimators: sensitivity to sample grain size. *Journal of Animal Ecology*, 75(1), 274–287. <https://doi.org/10.1111/j.1365-2656.2006.01048.x>
- Howell, S. N. G. and Webb, S. (1995). A guide to the birds of Mexico and Northern Central America. Oxford University Press, New York.  
<https://doi.org/10.1093/oso/9780198540137.001.0001>
- Hsieh, T. C., Ma, K. H. and Chao, A. (2016). iNEXT: An R package for interpolation and extrapolation of species diversity (Hill numbers). *Methods in Ecology and Evolution*, 7(12), 1451–1456. <https://doi.org/10.1111/2041-210X.12613>

- INEGI (2020). Página electrónica institucional, Instituto Nacional de Estadística y Geografía. Retrieved on 01 February 2023 from [www.inegi.org.mx](http://www.inegi.org.mx)
- Jost, L. (2006). Entropy and diversity. *Oikos*, 113(2), 363–375.  
<https://doi.org/10.1111/j.2006.0030-1299.14714.x>
- Lessi, B. F., Pires, J. S. R., Batisteli, A. F. and MacGregor-Fors, I. (2016). Vegetation, urbanization, and bird richness in a Brazilian periurban area. *Ornitología Neotropical*, 27, 203–210. <https://doi.org/10.58843/ornneo.v27i0.63>
- Leveau, L. M. and Leveau, C. M. (2016). Does urbanization affect the seasonal dynamics of bird communities in urban parks? *Urban Ecosystems*, 19, 631–647.  
<https://doi.org/10.1007/s11252-016-0525-5>
- Luna-Vega, I., Espinosa, D. and Contreras-Medina, R. (2016). Biodiversidad de la sierra Madre del Sur: una síntesis preliminar. Universidad Nacional Autónoma de México, México.
- MacGregor-Fors, I. and Payton, M. E. (2013). Contrasting diversity values: statistical inferences based on overlapping confidence intervals. *PLoS ONE*, 8, e56794.  
<https://doi.org/10.1371/journal.pone.0056794>
- MacGregor-Fors, I. and Schondube, J. E. (2011). Gray vs. green urbanization: relative importance of urban features for urban bird communities. *Basic and Applied Ecology*, 12(4), 372–381. <https://doi.org/10.1016/j.baae.2011.04.003>
- MacGregor-Fors, I., Escobar-Ibáñez, J. F., Schondube, J. E., Zuria, I., Ortega-Álvarez, R., Sosa-López, J. R., Ruvalcaba-Ortega, I., Almazán-Núñez, R. C., Arellano-Delgado, M., Arriaga-Weiss, S. L., Calvo, A., Chapa-Vargas, L., Silvestre, P. X.,

- García-Chávez, J. H., Hinojosa, O., Koller-González, J. M., Lara, C., López de Aquino, S., López-Santillán, D., Maya-Elizarrarás, E. and Vega-Rivera, J. H. (2021). The urban contrast: a nationwide assessment of avian diversity in Mexican cities. *Science of The Total Environment*, 753, 141915.  
<https://doi.org/10.1016/j.scitotenv.2020.141915>
- MacGregor-Fors, I., Gómez-Martínez, M. A., García-Arroyo, M. and Chávez-Zichinelli, C. A. (2020). A dead letter? Urban conservation, management, and planning strategies from the Mexican urban bird literature. *Urban Ecosystems*, 23, 1107–1115.  
<https://doi.10.1007/s11252-020-00970-y>
- Magurran, A. E. (2004). Measuring biological diversity. Blackwell Publishing, Oxford.
- Mikami, K., Morimoto, G., Ueno, Y. and Mikami, O. K. (2022). Vertical space utilization by urban birds and their relationship to electric poles and wires. *Landscape and Ecological Engineering*, 18, 19–30. <https://doi.10.1007/s11355-021-00479-2>
- Møller, A. P. (2014). Life history, predation and flight initiation distance in a migratory bird. *Journal of Evolutionary Biology*, 27(6), 1105–1113.  
<https://doi.10.1111/jeb.12399>
- Muller-Dombois, D. and Ellenberg, H. (1974). Aims and methods of vegetation ecology. John Wiley and Sons, New York.
- Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B., Solymos, P., Stevens, M. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M., Durand, S., Antoniazzi, H., FitzJohn, R., Friendly, M., Furneaux, B., Hannigan, G., Hill,

- M. O., Lahti, L., McGlinn, D., Ouellette, M. H., Cunha, E., Smith, T., Stier, A. and Oksanen, C. J. (2022). Vegan: community ecology package, R package version 2.6-2. Retrieved on 01 February 2023 from <https://cran.r-project.org/web/packages/vegan/index.html>
- Ortega-Álvarez, R. and MacGregor-Fors, I. (2009). Living in the big city: effects of urban land-use on bird community structure, diversity, and composition. *Landscape and Urban Planning*, 90(1-3), 189–195. <https://doi.10.1016/j.landurbplan.2008.11.003>
- Palacio, F. X. (2020). Urban exploiters have broader dietary niches than urban avoiders. *Ibis*, 162(1), 42–49. <https://doi.10.1111/ibi.12732>
- Pisanty, I., Mazari, M. and Ezcurra, E. (2009). El reto de la conservación de la biodiversidad en zonas urbanas y periurbanas. In: Dirzo, R., González, R. and March I. J. (Eds.) Capital natural de México, vol. II: estado de conservación y tendencias de cambio. Pp. 719–759. CONABIO, México.
- Polak, M., Wiącek, J., Kucharczyk, M. and Orzechowski, R. (2013). The effect of road traffic on a breeding community of woodland birds. *European Journal of Forest Research*, 132, 931–941. <https://doi.10.1007/s10342-013-0732-z>
- Puga-Caballero, A., MacGregor-Fors, I. and Ortega-Álvarez, R. (2014). Birds at the urban fringe: avian community shifts in different peri-urban ecotones of a megacity. *Ecological Research*, 29(4), 619–628. <https://doi.10.1007/s11284-014-1145-2>
- R Development Core Team. (2020). R: A language and environment for statistical computing. Retrieved on 01 February 2023 from <https://www.r-project.org>

- Reynolds, R. T., Scott, J. M. and Nussbaum, R. M. (1980). A variable circular-plot method for estimating bird numbers. *The Condor*, 82(3), 309–313. <https://doi.10.2307/1367399>
- Ripley, B., Venables, B., Bates, D. M., Hornik, K., Gebhardt, A. and Firth, D. (2022). Support functions and datasets for venables and Ripley's MASS. R package version 7.3-57, Retrieved on 01 February 2023 from <https://cran.r-project.org/web/packages/MASS/index.html>
- Schrimpf, M. B., Des Brisay, P. G., Johnston, A., Smith, A. C., Sánchez-Jasso, J., Robinson, B. G., Warrington, M. H., Mahony, N. A., Horn, A. G., Strimas-Mackey, M., Fahrig, L. and Koper, N. (2021). Reduced human activity during COVID-19 alters avian land use across North America. *Science Advance*, 7, eabf5073. <https://doi.10.1126/sciadv.abf5073>
- Seto, K. C. and Ramankutty, N. (2016). Hidden linkages between urbanization and food systems. *Science*, 352(6288), 943–945. <https://doi.10.1126/science.aaf7439>
- Sol, D., González-Lagos, C., Moreira, D. and Lapiedr, O. (2014). Urbanization tolerance and the loss of avian diversity. *Ecology Letters*, 17(8), 942–950. <https://doi.10.1111/ele.12297>
- Ter Braak, C. J. F. and Šmilauer, P. (2002) CANOCO Reference manual and CanoDraw for Windows user's guide: software for canonical community ordination (version 4.5). Wageningen University and Research, New York.
- Ter Braak, C. J. F. and Verdonschot, P. F. M. (1995). Canonical correspondence analysis and related multivariate methods in aquatic ecology. *Aquatic Sciences*, 57, 255–289. <https://doi.10.1007/BF00877430>




United Nations. (2019). World population prospects: highlights. Retrieved on 05 February 2023 from <https://population.un.org/wup/>

Zhou, D. and Chu, L. M. (2014). Do avian communities vary with season in highly urbanized Hong Kong? *Wilson Journal Ornithology*, 126(1), 69–80.  
<https://doi.10.1676/13-097.1>


Zúñiga-Palacios, J., Zuria, I., Moreno, C. E., Almazán-Núñez, R. C. and González-Ledesma, M. (2020). Can small vacant lots become important reservoirs for birds in urban areas? A case study for a Latin American city. *Urban Forestry and Urban Greening*, 47, 126551. <https://doi.10.1016/j.ufug.2019.126551>

**Table 1.** Observed (Robs), estimated (Chao 2) species richness, and urban bird types in urban and peri-urban parks in a Neotropical Mexican city. Urban and peri-urban parks are named in Fig. 1.


	Urban parks			Peri-urban parks		
	PMM	ALA	JBO	PEM	CDM	USN
Robs	26	24	47	41	46	46
% Chao 2	76	92	73	87	90	90
Urban exploiter birds	5	6	6	7	7	7
Urban adapter birds	20	18	30	23	28	29
Urban avoider birds	1	-	11	11	11	10




**Fig. 1.** Location of Chilpancingo City in Guerrero state, southern Mexico. (a) urban (1-3) and peri-urban parks (4-6): 1) Parque Margarita Maza (PMM), 2) Alameda (ALA), 3) Jardín Botánico (JBO), 4) Parque Ecológico Manantiales (PEM), 5) Club Deportivo Mitsumarú (CDM), 6) Universidad Sentimientos de la Nación (USN). (b) Land-cover classes in the urban and peri-urban parks delimited by 200 m buffer areas.



**Fig. 2.** Mean values of richness (q0) and diversity (q1 and q2) of urban bird species in urban and periurban parks during the breeding and non-breeding seasons in a Neotropical Mexican city. Error bars represent 84 % IC. Urban and peri-urban parks are named in (Fig. 1).



**Fig. 3.** Rank-abundance curves for urban bird species in urban and peri-urban parks during the breeding and non-breeding seasons in a Neotropical city in a Neotropical Mexican city. Urban bird groups: exploiters (blue), adapters (red), avoiders (green). Urban and peri-urban parks are named in (Fig. 1).



**Fig. 4.** Canonical ordination of exploiter (blue circles), adapter (red diamonds), avoider birds (green triangles) at the local (upper panels) and landscape level (lower panels) during the breeding (a and c) and non-breeding seasons (b and d) in a Neotropical Mexican city. Urban bird codes are shown in Table S3. Landscape-scale variables: asphalt streets (ASP), dirt road, build-up areas (BUI), bare ground (BGR), vegetation (VEG).

Supplementary Material

**Table S1.** Local and landscape variables in urban and peri-urban parks in a Neotropical Mexican city. Urban and peri-urban parks are named in (Fig. 1).

Variables	Urban parks			Peri-urban parks		
	PMM	ALA	JBO	PEM	CDM	USN
<b>Local-level</b>						
Vegetation traits						
Plant richness	6 ± 3.54	7 ± 1.42	10 ± 3.47	6 ± 0.71	4 ± 1.42	5 ± 0.58
Plant abundance	16 ± 3.54	16 ± 4.25	15 ± 5.04	16 ± 4.95	12 ± 0.71	16 ± 4.51
Plant dominance (D')	0.45 ± 0.40	0.25 ± 0.1	0.15 ± 0.07	0.34 ± 0.07	0.43 ± 0.24	0.30 ± 0.10
Cover (m <sup>2-1</sup> )	2834.21 ±	3321.07 ±	6269.27 ±	5389.40 ±	6800.79 ±	4235.67 ±
	525.92	2459.22	409.87	333.59	5714.3	1347.17
Density (m <sup>2</sup> )	0.13 ± 0.03	0.13 ± 0.04	0.12 ± 0.04	0.13 ± 0.04	0.10 ± 0.01	0.13 ± 0.04
Height (m)	8.03 ± 1.89	5.39 ± 0.24	5.64 ± 0.84	6.76 ± 0.36	7.67 ± 2.23	5.78 ± 0.81
Diameter at breast height (cm)	71.13 ±	71.63 ±	67.35 ±	95.88 ±	128.61 ±	47.24 ±
	10.36	15.35	18.29	14.79	99.81	10.36
Foliage height diversity (H')	1.63 ± 0.09	1.61 ± 0.10	1.58 ± 0.28	1.73 ± 0.11	1.65 ± 0.08	1.23 ± 0.14
Urbanization traits						
Noise (db)	55.78 ± 1.04	50.57 ± 1.28	25 ± 21.66	36.07 ± 1.33	34 ± 2.07	42.31 ± 1.22
N° people	5.89 ± 0.21	33.07 ± 2.8	0.6 ± 0	2.36 ± 0.51	1.82 ± 0.38	1.74 ± 0.42
N° pets	4 ± 1.41	1.36 ± 0.14	0 ± 0	1.75 ± 0.37	0.46 ± 0.14	0.45 ± 0.11
N° wires	9 ± 4.24	22 ± 4.24	6.33 ± 3.21	11.50 ± 0.71	7.5 ± 2.12	9 ± 4.58
N° light posts	7.5 ± 4.94	9.5 ± 2.12	3.33 ± 1.52	6.50 ± 2.13	6.5 ± 2.12	1.66 ± 0.57
<b>Landscape-level</b>						
Asphalt streets	38.7	22.5	32.5	11.6	28	28.9
Dirt road	0.8	-	-	2.1	1.7	2.6
Built-up areas	37	67.3	69.9	17.2	12	16.8
Bare ground	16.7	6.5	15	31.9	21.7	60.4
Vegetation	6.7	1.1	9.5	21	33.6	8.2

**Table S2.** Ordination values of the principal components (PC) of the 11 vegetation and urbanization variables in the urban and peri-urban parks during breeding and non-breeding seasons in a Neotropical Mexican city. Significant correlations are as follows:  $p \leq 0.05$  (\*),  $p \leq 0.01$  (\*\*),  $p \leq 0.001$  (\*\*\*).

Variables	Breeding season			Non-breeding season		
	PC1	PC2	PC3	PC1	PC2	PC3
Species richness	-0.15	0.08	-0.89**	0.13	-0.01	-0.9***
Plant abundance	0.38	-0.75	-0.25	0.36	0.81**	-0.06
Plant dominance	0.69	-0.48	0.49	0.5	0.41	0.7
Cover (m <sup>2-1</sup> )	-0.53	0.8**	-0.26	-0.45	-0.74	-0.49
Height (m)	-0.19	0.13	0.95***	0.4	-0.1	0.77*
Foliage height diversity (H')	-0.32	0.74*	-0.12	-0.4	-0.59	-0.30
Diameter at breast height (cm)	0.18	0.95***	0.2	0.07	-0.98***	0.11
Noise (db)	0.58	-0.71	0.22	0.75	0.62	0.05
Nº people	0.91***	-0.32	-0.2	0.91***	0.37	-0.002
Nº wires	0.9***	-0.33	-0.13	0.89**	0.33	0.15
Nº light posts	0.96***	0.004	0.15	0.88**	-0.08	0.37
Eigenvalues	5.8	2.29	1.67	6.2	2.05	1.4
Explained variance (%)	52.73	20.86	15.2	56.36	18.77	12.7
Cumulative variance (%)	52.73	73.6	89	56.36	75.13	88

**Table S3.** Composition and relative abundance of urban bird species in urban and peri-urban parks during breeding and non-breeding seasons in a Neotropical Mexican city.

Urban and peri-urban parks are named in (Fig. 1). Urban classification (UC): urban exploiters (UrEx), urban adapters (UrAd), urban avoiders (UrAv). Urban and peri-urban parks are named in (Fig. 1).

Taxon	UC	Code	Breeding season						Non-breeding season					
			PMM	ALA	JBO	PEM	CDM	USN	PMM	ALA	JBO	PEM	CDM	USN
Columbidae														
<i>Columba livia</i>	UrEx	Cv		0.018					0.017					
<i>Streptopelia decaocto</i>	UrEx	Sd	0.047	0.096	0.018	0.044	0.003	0.068	0.027	0.164	0.024	0.066		0.008
<i>Columbina inca</i>	UrAd	Ci	0.037	0.158	0.023	0.092	0.131	0.073	0.011	0.144	0.018	0.023	0.069	0.059
<i>Columbina passerina</i>	UrAd	Cp					0.015	0.006			0.008			0.003
<i>Leptotila verreauxi</i>	UrAv	-					0.002				0.002			
<i>Zenaida asiatica</i>	UrAd	Za			0.002	0.012			0.015		0.024	0.027	0.008	0.028
Cuculidae														
<i>Crotophaga sulcirostris</i>	UrAd	Cs			0.014	0.029	0.029				0.012	0.015	0.003	0.034
<i>Piaya cayana</i>	UrAd	Pc					0.003				0.006		0.005	
Trochilidae														
<i>Cynanthus doubledayi</i>	UrAd	Cd				0.002		0.002		0.006		0.004		0.003
<i>Basilinna leucotis</i>	UrAv	Bl	0.007	0.004		0.002	0.002	0.003						
<i>Ramosomyia violiceps</i>	UrAd	Rv			0.005		0.003		0.004	0.023	0.01	0.008	0.005	0.02
Strigidae														
<i>Glaucidium brasilianum</i>	UrAd	Gb			0.005								0.005	0.003
Momotidae														
<i>Momotus mexicanus</i>	UrAd	Mm			0.023	0.012	0.02	0.011			0.008	0.008	0.005	0.02
Picidae														
<i>Melanerpes chrysogenys</i>	UrAd	Mc	0.02		0.075	0.029	0.033	0.017	0.004		0.026	0.023	0.046	0.028
Psittacidae														
<i>Eupsittula canicularis</i>	UrAd	Ec			0.002				0.015	0.023				
Tyrannidae														
<i>Camptostoma imberbe</i>	UrAv	-			0.002									
<i>Myarchus nuttingi</i>	UrAv	Mn									0.008			0.006
<i>Myiarchus tyrannulus</i>	UrAd	-	0.007											
<i>Pitangus sulphuratus</i>	UrAd	Ps		0.005	0.046	0.01	0.028	0.041		0.006	0.014	0.012	0.02	0.039
<i>Megarynchus pitangua</i>	UrAv	-					0.003							
<i>Myiozetetes similis</i>	UrAd	Ms				0.002	0.003	0.003				0.004		0.011
<i>Tyrannus melancholicus</i>	UrAd	Tm	0.044	0.042	0.041	0.027	0.03	0.032	0.054	0.073	0.037	0.035	0.043	0.048
<i>Tyrannus vociferans</i>	UrAv	Tv												0.014
<i>Tyrannus crassirostris</i>	UrAv	Tc					0.002					0.004		0.006
<i>Tyrannus verticalis</i>	UrAd	Te				0.014	0.011		0.027		0.006		0.005	0.008
<i>Empidonax minimus</i>	UrAv	-												0.006
<i>Pyrocephalus rubinus</i>	UrAd	Pu						0.027	0.011	0.096				0.02



Vireonidae														
<i>Vireo hypochryseus</i>	UrAv	Vh					0.013		0.004	0.003	0.002		0.003	
<i>Vireo gilvus</i>	UrAd	Vg									0.01			
Laniidae														
<i>Lanius ludovicianus</i>	UrAd	Ll					0.005						0.005	0.006
Hirundinidae														
<i>Stelgidopteryx serripennis</i>	UrAd	Ss			0.005	0.01	0.011	0.013				0.023	0.13	0.006
<i>Hirundo rustica</i>	UrAd	Hr	0.081	0.031	0.091	0.137	0.136	0.063	0.027			0.139	0.064	0.017
Ptiliognathidae														
<i>Ptiliognathus cinereus</i>	UrAd	Pi	0.02	0.071	0.002		0.052	0.016	0.057	0.011	0.002		0.015	0.025
Poliophtidae														
<i>Poliophtila caerulea</i>	UrAd	Pa							0.042	0.054	0.142	0.039	0.054	0.048
Troglodytidae														
<i>Catherpes mexicanus</i>	UrAd	-				0.002	0.002							
<i>Thryophilus sinaloa</i>	UrAv	Ts				0.002	0.003					0.008		
<i>Thryophilus pleurostictus</i>	UrAv	-										0.004		
Mimidae														
<i>Melanotis caerulescens</i>	UrAv	Me			0.002		0.003					0.008		
<i>Toxostoma curvirostre</i>	UrAd	Tu		0.004	0.007	0.002	0.026	0.003				0.008	0.003	0.008
<i>Mimus polyglottos</i>	UrAv	-												0.003
Turdidae														
<i>Turdus rufopalliat</i>	UrAd	Tr		0.031	0.094	0.037	0.059	0.041	0.011	0.051	0.209	0.085	0.084	0.11
Passeridae														
<i>Passer domesticus</i>	UrEx	Pd	0.359	0.427	0.027	0.17	0.013	0.09	0.31	0.274	0.002	0.224	0.01	0.11
Fringillidae														
<i>Haemorhous mexicanus</i>	UrAd	Hm	0.003	0.005	0.009	0.002	0.003	0.022	0.027	0.003			0.003	
<i>Spinus psaltria</i>	UrAd	Sp							0.008		0.004			
Passerellidae														
<i>Peucaea ruficauda</i>	UrAd	Pf			0.002	0.039	0.033	0.097				0.012	0.013	0.082
<i>Peucaea humeralis</i>	UrAv	Ph						0.006						
<i>Melospiza kieneri</i>	UrAv	Mk				0.002	0.003							0.006
Icteridae														
<i>Icterus wagleri</i>	UrAv	Iw					0.003	0.003				0.046		0.008
<i>Icterus spurius</i>	UrAd	Is								0.028	0.014	0.004	0.005	0.014
<i>Icterus pustulatus</i>	UrAd	Ip			0.005	0.012	0.007	0.028	0.008	0.003	0.037	0.031	0.013	0.048
<i>Icterus bullockii</i>	UrAd	Ib							0.004		0.004			0.014
<i>Icterus pectoralis</i>	UrAv	Ie				0.006	0.002	0.002						
<i>Icterus gularis</i>	UrAv	-			0.002									
<i>Icterus galbula</i>	UrAv	Ig									0.008	0.019		0.006
<i>Molothrus aeneus</i>	UrEx	Ma				0.002	0.028	0.003				0.004	0.087	
<i>Molothrus ater</i>	UrAd	Mt			0.005		0.008				0.002		0.008	
<i>Quiscalus mexicanus</i>	UrEx	Qm	0.322	0.089	0.429	0.194	0.154	0.111	0.28		0.211	0.081	0.209	0.093
Parulidae														
<i>Mniotilta varia</i>	UrAv	Mn										0.004		
<i>Leiothlypis ruficapilla</i>	UrAd	Lr							0.015	0.006	0.014	0.015	0.018	0.006
<i>Setophaga coronata</i>	UrAd	So							0.019	0.003	0.099	0.004	0.038	0.011
<i>Cardellina pusilla</i>	UrAd	-									0.002	0.004		
Cardinalidae														

<i>Piranga rubra</i>	UrAv	Pr									0.006			
<i>Piranga ludoviciana</i>	UrAd	Pl							0.004	0.006	0.002			0.008
<i>Pheucticus melanocephalus</i>	UrAv	Pm									0.016			
<i>Passerina versicolor</i>	UrAv	Pv	0.003		0.002	0.004								
Thraupidae														
<i>Sporophila torqueola</i>	UrAd	St	0.04	0.015	0.048	0.08	0.048	0.158	0.015	0.008	0.008	0.008	0.02	0.006
<i>Saltator grandis</i>	UrAv	Sg			0.009	0.02	0.041	0.013				0.004	0.003	0.003