

A NEW GENUS OF RHYTIRRHININI FROM COLOMBIA (COLEOPTERA: CURCULIONIDAE)

JUAN J. MORRONE

Laboratorio de Sistemática y Biología Evolutiva (LASBE), Museo de La Plata, Paseo del Bosque, 1900 La Plata, Argentina.

Resumen

Rupanius, gen. n., comprende la única especie *R. carinatus*, sp. n., endémica de la provincia biogeográfica del Páramo de Colombia. Este género se reconoce por los élitros subcuadrados y con una conspicua carena declivital y es considerado similar a *Puranus* Germain y *Acrorius* Kirsch. El género y especie son descritos, ilustrados y comparados con los otros Rhytirrhinini de Colombia.

Abstract

Rupanius, new genus, comprises the single species *R. carinatus*, new species, endemic to the Páramo biogeographic province of Colombia. It is diagnosed based on the elytra subquadrate and with a conspicuous declivital carina, and is considered similar to *Puranus* Germain and *Acrorius* Kirsch. The genus and species are described, illustrated, and compared with the other Colombian Rhytirrhinini.

The South American Rhytirrhinini (Coleoptera: Curculionidae, Somatodinae) are basically distributed along the Andean range, with their highest diversity in the Subantarctic region of southern Argentina and Chile (MORRONE, 1993). I have recently revised some genera from the northern Andes, assigning species from Colombia to *Acrorius* Kirsch (MORRONE, 1994a) and *Macrostyphlus* Kirsch (MORRONE, 1994b). The main objective of this paper is to describe a new taxon from Colombia, *Rupanius carinatus* new genus, new species, based on material loaned to me by Bob Anderson (Canadian Museum of Nature, Ottawa, Canada). In addition to the descriptions, a key for the genera and species of Colombian Rhytirrhinini is provided.

Key to the Colombian Rhytirrhinini

1. Body vestiture of rounded scales; pronotum subcylindrical, longer than wide, lacking postocular lobes, disc usually not tuberculate; metepisternal suture absent; body length 1.9-3.5 mm *Macrostyphlus* Kirsch 3
- Body vestiture of setalike scales; pronotum transverse, wider than long, with postocular lobes, disc tuberculate; metepisternal
- suture present; body length > 4.0 mm ...
2. Elytra ovate, with anteapical tubercle and lacking declivital carina; female hemisternites lacking baculi; body length 4.0-6.8 mm
Acrorius Kirsch (*A. sisyphus* Morrone)
Elytra subquadangular, lacking anteapical tubercle and with declivital carina; female hemisternites with baculi; body length 5.3-6.6 mm *Rupanius*, new genus (*R. carinatus*, new species)
3. Pronotum with apical impression; tibiae lacking mucro *M. frodo* Morrone
Pronotum lacking apical impression; tibiae mucronate 4
4. Antennae with funicular articles 3-6 transverse; pronotum with median carina *M. sturmi* Morrone
Antennae with funicular articles 3-6 moniliform; pronotum lacking median carina 5
5. Pronotum with two subcircular, lateral impressions; scutellum not prominent; elytra with small tubercles; tibiae lacking spurs *M. gandalf* Morrone

Pronotum lacking impressions; scutellum prominent; elytra lacking tubercles; tibiae with two spurs. *M. gualcalae* Kirsch

Rupanius Morrone, new genus

TYPE SPECIES. *R. carinatus* Morrone, new species

DIAGNOSIS. *Rupanius* is easily recognized by the conspicuous declivital carina on the elytra. In addition, it is separated from other Rhytirrhinini by the combination of the following characters: vestiture of setalike scales, suprascrobal keel absent, epistome slightly prominent, pronotal disc tuberculate, metepisternal suture present, elytra subquadangular and flat, and hemisternites with baculi.

DESCRIPTION. Habitus (figs. 1, 2). Medium-sized (body length 5.3-6.6 mm); body vestiture of setalike scales and setae. Frons with fovea. Eyes ovate, medium-sized, flat. Rostrum slightly curved, shorter than pronotum, with three dorsal keels. Scrobes shallow,

lateral, directed toward, but not reaching the eyes; ventral keel lacking teeth. Pterygia well developed. Epistome slightly protruding. Mandibles robust, external face with two setae. Antennae subapically inserted; scape exceeding hind margin of eye when resting in scrobe; funicular article 1 longer than 2, articles 4-7 monilliform; club inflated. Pronotum (fig. 1) slightly transverse, sides expanded in anterior third; postocular lobes narrow. Metepisternal suture present. Scutellum visible. Elytra (fig. 1) subquadangular, wider than pronotum, flat; humeri rounded; disc tuberculate; anteapical tubercle absent; apical declivity with conspicuous carina. Legs with robust femora; tibiae slightly mucronate, with one spur; tarsomere 3 bilobed.

MALE GENITALIA. Aedeagus (figs. 3, 4) symmetrical, sclerotized, robust in lateral view, sides subparallel. Tegmen lacking parameres.

FEMALE GENITALIA. Sternum 8 (figs. 5) subpentagonal, with two sclerotized arms; apical margin with long setae; apodeme straight. Hemisternites (figs. 6) short, with baculi; styli absent. Spermatheca (figs. 7) with nodulus and ramus developed.


REMARKS. *Rupanius* is superficially similar to *Acrorius* Kirsch (MORRONE, 1994a) and *Puranus* Germain (MORRONE, 1994b). Its proper placement, however, cannot be properly assessed until a cladistic analysis of its related genera within the Rhytirrhinini is undertaken.

DISTRIBUTION. The single species was found at high altitudes (1800-3300 m) in Colombia. Its distribution broadly corresponds to the Paramo biogeographic province of Cabrera and WILLINK (1973).


ETYMOLOGY. The generic name is an anagram of *Puranus*. Gender masculine.

Rupanius carinatus Morrone, new species
figs. 1-7.

TYPE MATERIAL. Holotype female (labels separated by square brackets, with a slash to



Figuras. 1, 2. *Rupanius carinatus*, female holotype: 1, dorsal view; 2, lateral view.



Figuras. 3-7. *Rupanius carinatus*, male and female genitalia: 3, aedeagus, dorsal view; 4, aedeagus, lateral view; 5, female sternum 8, ventral view; 6, hemisternite; 7, spermatheca.

separate each line): COLOM., 20 km E/ Silvia, Cauca, VII./ 16.1970, 11,000'/ J. M. Campbell [CANADIAN NATIONAL/ COLLECTION- AGRICULTURE/ CANADA, OTTAWA] [*Rupanius carinatus*/ Morrone/ holotype female]; one paratype male: [COLOM., 20 km W/ Silvia, Cauca, VII./ 17.1970, 6,000'/ J. M. Campbell] [CANADIAN NATIONAL/ COLLECTION- AGRICULTURE/ CANADA, OTTAWA] [*Rupanius carinatus*/ Morrone/ paratype male]. Both types deposited in the Canadian National Collection of Insects, Center for Land and Biological Resources Research, Biological Research Division, Agriculture Canada, Ottawa, Canada.

DESCRIPTION. Holotype female. Habitus (figs. 1, 2). Integument dark brown. Seta-like scales whitish. Rostrum 1.6 times as long as wide, 0.7 times as long as pronotum. Antennae with article 1 of funicle 1.5 times longer than 2. Pronotum (fig. 1) 1.2 times wider than long, hind margin sinuate; disc strongly tuberculate.

Elytra (fig. 1) 1.2 longer than wide; striae inconspicuous, intervals flat; humeri not prominent; disc with declivital carina with tubercles. Sternum 8 (fig. 5) with apodemes 1.7 times longer than plate. Hemisternites (fig. 6). Spermatheca (fig. 7). Length (pronotum + elytra) 6.6 mm.

MALE. Aedeagus (figs. 3, 4) with rounded apex, slightly curved in apical third, apodemes as long as aedeagal body. Length (pronotum + elytra) 5.3 mm.

ETYMOLOGY. The name of this species refers to the elytral apical carina.

Acknowledgments

I thank BOB ANDERSON for loaning the specimens to me and FEDERICO OCAMPO for taking the habitus photographs. This study was supported by grant 4662-91 of the National Geographic Society. Support of the Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Argentina, to which I belong, is acknowledged.

Literature cited

- CABRERA, A. L. AND A. WILLINK. 1973. *Biogeografía de América Latina. Monografías de la OEA, serie de Biología*, (13).
- MORRONE, J. J. 1993. Revisión sistemática de un nuevo género de *Rhytiirrhinini* (Coleóptero: Curculionidae), con un análisis biogeográfico del dominio subantártico. *Bol. Soc. Biol. Concepción*. 64:121-145.
- . 1994a. Systematics of the Andean genus *Acrorius* Kirsch (Coleoptera: Curculionidae). *Coleo. Bull.* 48(2):101-114.
- . 1994b. Systematics, cladistics, and biogeography of the Andean weevil genera *Macrostyphlus*, *Adioristidius*, *Puranius*, and *Amathyinetoides*, new genus (Coleoptera: Curculionidae). *Am. Mus. Novit.* (3104):1-63.