

MORPHOMETRICS, MOLT, CLOACAL TEMPERATURES AND ECTOPARASITES IN COLOMBIAN BIRDS

YOSHIKA ONIKI
EDWIN O. WILLIS

Departamento de Zoología, UNESP, Caixa Postal 199 13.500 Rio Claro, S.P., Brasil.

Resumen

Se presenta los datos morfométricos, muda, temperaturas cloacales y ectoparásitos de 123 individuos de 57 especies de aves colombianas.

Abstract

Morphometric data, molt, cloacal temperatures and ectoparasites are presented for 123 individuos of 57 species of Colombian birds.

Weights and other morphological measurements of birds are essential to studies in a wide variety of disciplines. Karr, Wilson & Moriarty (1978) pointed out that weights of tropical birds could be used for testing hypotheses on a variety of ecological and evolutionary phenomena. Dunning (1984) listed recent studies in which bird weights have been used in avian physiology, ecology, morphology, community structure and theoretical modeling. A comprehensive review of the uses and potential of weights appeared in Clark (1979). Unfortunately these data are not always available from museum specimens.

This account reports the live weights, measurements and cloacal temperatures of 123 birds of 57 species in 1989 in Colombia. During examination of birds, molt and brood patches were recorded to ascertain breeding overlap with molt, as reported for some tropical species (Foster 1974, 1975).

Ectoparasites were also collected for further study.

Methods and study area

Birds were caught in mist nets, weighed with Pesola scales to the nearest 0.2 g, and cloacal temperatures were taken with a quick measuring Schultheis thermometer. Field work was conducted on 5-6 February at San Nicolas 2900 m at the Parque Nacional Puracé, Cauca, between 11 February and 14 March at the Reserva Natural La Planada, Nariño, and on 13 and 24 March

at the Estación de Bombeo Guamués, 700 m near Orito, Putumayo.

At Puracé, birds were netted in woods and in trails parallel to the main road; at La Planada in several locations off the main trails and at Orito at the petroleum pumping station, where birds gathered to feed on insects attracted by the light left on all night.

Culmens were measured from the feathers to tip, tarsi from tibiotarsal joint to above the first toe, tails between the central feathers, and wings as the unflattened chord.

In Table 1, taxonomic sequence follows Meyer de Schauensee (1970).

Results

The number of birds per net was not calculated because nets were open and closed irregularly to avoid bird mortality due to constant rains, especially at La Planada. Here, even though average air temperature throughout the day was 18°C, (n= 16) some birds died easily due to the rains.

At Puracé, 21 birds ("P" in Table 1) were caught in 1.5 days of mist netting, while at La Planada 90 birds were captured, and 10 birds ("G" in Table 1) were caught at Guamués. At La Planada, the 2 *Semnornis ramphastinus* were captive birds that had died; they could have been under-

weight, and their tail measurements are probably short since the tail feathers were worn and battered.

MOLT AND BREEDING - Table 1 shows molt of body, tail and wing as well as presence or absence of brood patch. One *Grallaricula flavivestris* had in its oviduct an egg formed and ready to be laid.

ECTOPARASITES - Ectoparasites for these Colombian birds are indicated in Table 2; descriptions of new species and other details will appear in the future.

Single fleas (Order Siphonaptera) were identified by R. Traub as *Dasyptyllus gallinulae*, said to be the most widely ranging species known and infesting a variety of avian hosts. On each bird there were at least 2 fleas, but they moved through the feathers so rapidly that it was difficult to catch them with forceps. According to R. Traub (in litt.), fleas from South American birds are relatively little-known. They are found mainly on birds of highland, cold areas such as these Andean ones; *D. gallinulae* is northern, and the nominate form occurs in Europe.

Due to low temperatures, ticks (Acari) attached preferentially at the edge of the outer ear where they were protected by the feathers. Compared to birds from warmer climates in Brasil, few (in *Diglossa* and *Anisognathus*) were exposed just above the eyes.

Mites (Order Acari) were mostly found on body, wing and tail feathers, except for nose mites in hummingbirds (hitching rides between flowers) and for chiggers.

Lice (Suborder Mallophaga) eggs were mainly found on chin, lores, above eyes and top of the head, where difficult for the bird to preen.

Discussion

According to Table 1, most birds presented body molt and many had finished molting the wing and tail feathers. Some species, mostly suboscines, still had brood patches, i. e., they were

breeding or ending breeding in Feb. -March. Songbirds mostly lacked brood patches. The lack of brood patches in songbirds may indicate that they have short breeding seasons, ending well before March, in contrast to long breeding seasons in the forest suboscines. Willis (1990) suggested that songbirds regularly occupy seasonal habitats and, for this reason, have short reproductive periods that require much song and prevent males from incubating eggs. Hummingbirds may keep their brood patches all year long, so we are not certain they were breeding in February and March.

Adult lice and eggs were found both on breeding and nonbreeding birds. However, we were only able to study the birds early in the year rather than throughout the year or at the July-August peak of nesting (at La Planada, according to G. Cantillo). Thus, we could not verify ectoparasite relationships with molt and breeding seasons, which are likely to vary widely in an equatorial country like Colombia.

Acknowledgements

We appreciate a grant from the National Geographic Society for field work in Colombia. A. Negret and G. Cantillo helped with logistics. K. Schuchmann, G. Corredor, C. I. Acevedo and C. Restrepo helped with field work. Drs. D. Scott Wood and Kenneth Parkes commented on the manuscript and helped with logistics. Dr. R. Traub identified the fleas. We also appreciate the help of the Empresa Colombiana de Petróleos (ECOPETROL, Distrito Sur) with transportation and permission to work at Guamués, of INDERENA for permission to visit and quarters at the National Park of Puracé, and of G. Cantillo and J. Orejuela for hospitality at La Planada. Writing was completed while Oniki was on an International Program at the Carnegie Museum of Natural History in Pittsburgh.

Literature cited

- CLARK, G.A. 1979. Body weights of birds: a review. *Cóndor* 81: 193-202.
- DUNNING Jr., J.B. 1984. Body weights of 686 species of North American birds. *Western Bird Banding Assoc., Monog.* No. 1: 38 p.
- FOSTER, M.S. 1974. A model to explain molt-breeding overlap and clutch size in some tropical birds. *Evolution* 28: 182-190.
- 1975. The overlap of molting and breeding in some tropical birds. *Cóndor* 77: 304-314.

KARR, J.R., M.F. WILLSON & D.J. MORIARTY. 1978. Weights of some Central American birds. *Brenesia* 14-15: 249-257.

MEYER DE SCHAUENSEE, R. 1970. *A Guide to the Birds of South America*. Wynnewood, PA., Livingston Publ. Co.

WILLIS, E.O. 1990. Are songbird success and non-incubating males related to seasonal habitats? *Acta XX Congr. Intern. Ornithol.*, Christchurch, New Zealand Suppl. p. 424.

Table 1. Measurements, molt, and cloacal temperatures of Colombian birds

	Sex	Wt.	Molt' b t w	Brood Patch	Bill	Tarsus	Tail	Wing	Cloacal Temp.
TROCHILIDAE									
<i>Doryfera ludoviciae</i>	M	7.1			3.6	0.5	3.5	6.1	
	M	5.9	+-	-	3.8	0.3	3.7	6.2	
<i>Phaethornis syrmatophorus</i>		6.1	+++	+	3.9	0.5	7.0	6.3	
		6.0	+		3.8	0.4	6.8	5.8	
	M	6.2	-	-	3.8	0.3	7.0	6.0	
<i>Colibri coruscans</i> P		8.5	+++	+	2.6	0.6	5.2	8.0	
<i>Urochroa bougueri</i>	M	12.5		-	3.2	0.6	5.2	8.5	41.4
	M	11.2	+++	-	3.1	0.5	5.1	8.1	39.7
<i>Lafresnaya lafresnayi</i> P		6.1	+++	+	2.8	0.5	4.0	6.3	
<i>Coeligena wilsoni</i>		6.7	--	+	2.8	0.5	4.3	6.6	37.2
	F	6.5	+++		3.5	0.6	4.1	7.1	
	M	7.0	+++		3.6	0.2	4.3	7.0	
		6.9	++		3.6	0.5	3.8	6.6	
<i>Heliangelus exortis</i> P		4.8	+++		1.7	0.6	4.0	6.7	
<i>Eriocnemis vestitus</i> P	M	4.1	+++	+	1.8	0.6	4.2	6.1	
<i>Haplophaedia lugens</i>		6.0	+++	-	1.8	0.6	4.1	6.3	40.6
	F	6.9	+++	-	1.8	0.7	4.2	6.8	38.3
	F	8.8	+	+	1.9	0.5	4.1	6.1	
<i>Metallura tyrianthina</i> P	F	3.1	+++	+	1.2	0.6	3.2	5.1	
P	M	3.6	+++	+	1.2	0.6	4.0	5.9	
<i>Agelaiocercus coelestis</i>	M	5.0	+++	-	1.6	0.5	7.4	6.0	
	F?	5.0	+	-	1.4	0.5	4.0	5.6	
	M	4.7	+	-	1.5	0.5	4.1	5.3	
	F	4.7	+	-	1.6	0.5	4.2	5.6	
CAPITONIDAE									
<i>Semnornis ramphastinus</i>	M	99.5			2.3	3.2	6.5	9.4	
	M	73.0			2.0	2.5	10.1		
DENDROCOLAPTIDAE									
<i>Xiphocolaptes promeropyrhynchus</i>		123.8	+++	+	4.6	3.4	10.8	13.7	34.8
FURNARIIDAE									
<i>Synallaxis azarae</i>	M	16.0	+	-	1.2	2.4	8.8	5.4	38.8
<i>Premnoplex brunnescens</i>	F	15.4	+++	+	1.6	2.0	5.5	5.9	40.0
<i>Premnornis guttuligera</i>		15.1	+++	-	1.4	2.0	6.3	6.6	42.0
<i>Thripadectes virgaticeps</i>		65.0	+		2.8	3.2	9.2	10.3	36.6
		66.5	+		2.4	3.2	8.6	10.1	
<i>Sclerurus mexicanus</i>		24.0	+++	-	2.5	2.4	5.6	7.7	40.7
FORMICARIIDAE									
<i>Grallaria flavotincta</i>		63.0	+++	-	2.5	5.0	4.5	9.6	39.0
<i>Grallaricula flavirostris</i>	F	18.0	+++	-	1.3	2.3	2.5	6.7	
	M	19.5	+	+	1.5	2.5	2.8	7.2	
	F	21.5	+++	+	1.4	2.3	2.7	6.7	

COTINGIDAE

<i>Pipreola riefferii</i>		52.0	+	+	1.3	2.6	7.0	9.3	38.2
		44.9	+	-	1.4	2.4	7.1	9.1	40.0
		51.0	--	+	1.2	2.8	7.1	9.6	
<i>Pachyramphus versicolor</i>	F	15.4		--+	1.1	1.8	4.6	6.3	41.0

PIPRIDAE

<i>Masius chrysopterus</i>	F	12.0	--	-	0.8	1.8	4.0	5.9	38.6
		12.0	---	-	0.8	2.0	4.2	6.1	37.6
	F	12.8	+	-	0.8	1.7	4.1	5.8	40.2
		7.0	+	+	0.9	1.8	4.1	5.9	39.5

TYRANNIDAE

<i>Myiophobus flavicans</i>	M	12.9	+--	-	1.2	1.8	5.5	6.8	41.1
	F	11.4	+		1.2	1.8	5.5	6.4	
<i>Platyrinchus mystaceus</i>	F	9.0	+--	-	1.1	1.3	2.9	5.6	38.4
<i>Rhynchocyclus fulvipectus</i>	M	11.4		+	1.4	1.8	6.6	7.7	39.0
<i>Tyranniscus cinereiceps</i>		9.5	---	-	0.9	1.6	5.2	6.1	37.6
<i>Mionectes striaticollis</i>	M	14.5		+	1.2	1.9	4.8	7.0	

HIRUNDINIDAE

<i>Notiochelidon murina</i> P		12.1	+--	-	0.6	2.0	5.7	11.5	
-------------------------------	--	------	-----	---	-----	-----	-----	------	--

TROGLODYTIDAE

<i>Troglodytes aedon</i> G		15.0	+--	-	1.3	1.9	4.2	5.6	40.2
	G	17.5	+--	+	1.2	1.9	4.4	5.8	40.1
	G	14.5	---		1.2	2.0	3.7	5.4	40.8
<i>Henicorhina leucophrys</i>		18.2			1.5	2.5	2.8	5.7	
		15.5	+--	-	1.5	2.7	2.5	5.8	37.3
		18.2	--	-	1.5	2.5	2.6	5.5	38.4

TURDIDAE

<i>Myadestes ralloides</i>		32.0	+--	-					38.4
		31.0		-	1.1	2.1	7.2	8.5	37.8
		28.0	--	-	1.1	2.4	7.2	8.8	37.2
		27.0							
<i>Catharus ustulatus</i>		28.0	--	-	1.2	3.1	6.8	10.2	41.6
<i>Turdus fuscater</i> P		130.5	+--	-	2.9	4.8	14.0	15.2	
<i>Turdus ignobilis</i> G	F	68.0	+--	+	2.0	2.6	8.1	11.2	41.8
	M?	63.0		-	1.9	2.8	8.7	11.8	42.3

VIREONIDAE

<i>Vireo leucophrys</i>	F	11.3		-	1.0	1.7	4.6	6.7	37.2
	F	12.0		-	1.2	1.9	4.5	6.5	37.7
		11.5	---	-	1.0	1.8	4.8	6.8	38.4

PARULIDAE

<i>Dendroica fusca</i>	F	9.8	+	-	1.0	1.9	4.5	6.6	41.6
	F?	9.0	+--	-	1.0	1.7	4.5	6.4	41.2
<i>Basileuterus tristriatus</i>	M	12.3	+--	-	1.2	2.2	5.3	6.1	36.9
	M	13.0	+--	-	1.2	2.2	5.5	6.3	37.0
	M	12.5	--	-	1.2	2.1	5.7	6.9	37.8

COEREBIDAE

<i>Diglossa albilatera</i> P		11.0	++	-	1.0	2.0	4.9	6.1	
		10.2	+--	-	1.0	1.9	4.8	5.8	
	M	8.7	---	-	0.9	2.0	4.9	6.0	37.4
		9.5	--	-	0.9	2.0	4.9	6.0	39.4
<i>Diglossa cyanea</i> P		17.2	+--	-	1.6	2.9	5.9	7.2	
		16.5	+--	-	1.4	2.3	5.6	7.1	

THRAUPIDAE

<i>Euphonia xanthogaster</i>	M	13.7	++-	-	0.8	1.7	3.4	6.2	37.2
		14.9	---	-	0.9	1.6	3.6	6.6	38.9
		14.0		-	0.8	1.5	3.2	6.0	41.5
	F	13.7	+	-	0.8	1.1	3.2	6.0	40.7
<i>Cholorochrysa phoenicotis</i>	M	21.5			1.2	2.2	4.2	7.3	40.8
<i>Tangara arthus</i>	M	20.0			1.1	1.8	5.1	7.8	40.0
	F	19.1	+++	+	1.0	1.9	4.7	7.1	41.8
	F	19.0	+	-	1.1	1.9	5.8	7.7	40.5
		18.0	+++	-	1.1	1.3	4.6	7.0	40.3
		21.5	+++		1.0	1.8	4.7	7.7	39.6
<i>Tangara vassorii</i> P		19.0	+++	-	0.8	2.0	5.2	7.9	
<i>Tangara labradorides</i>	M	14.1		-	0.9	1.8	4.9	6.6	39.0
	M	14.5			1.0	1.8	4.5	6.5	39.8
		12.9	+	-	1.0	1.8	4.5	6.2	40.8
<i>Tangara nigroviridis</i>	F	16.1			0.8	1.8	4.5	6.6	41.4
<i>Iridisornis rufivertex</i> P		25.0	+++	-	1.0	2.4	7.8	8.5	
<i>Anisognathus lacrymosus</i> P		43.0	+-	-	1.3	2.7	7.8	9.8	
<i>Anisognathus flavinucha</i>		43.0		-	1.6	2.4	7.0	9.1	42.0
		50.0	-		1.9	1.5	7.0	9.2	
		41.4	+	-	1.7	2.4	6.8	8.8	41.0
<i>Thraupis episcopus</i> G	F	35.0	++	-	1.4	2.1	6.6	9.2	42.5
<i>Thraupis cyanocephala</i>	F	34.0	++	-	1.4	2.4	6.8	8.6	40.8
	M	32.0	---	-	1.3	2.5	6.9	8.7	41.0
<i>Creurgops verticalis</i>	F	22.1	+	+	1.4	2.1	6.5	7.2	42.4
FRINGILLIDAE									
<i>Tiaris olivacea</i>	F	9.0	+++	-	0.9	1.9	3.8	5.1	41.0
<i>Atlapetes pallidinucha</i> P		34.0	+++		1.2	3.1	7.8	8.0	
		37.0	+	-	1.3	3.7	8.3	8.3	
<i>Atlapetes schistaceus</i> P		27.0	+++	-	1.2	2.9	8.6	8.0	
<i>Myospiza aurifrons</i> G	F	18.4	+++	-	1.2	2.3	4.5	6.2	40.6
	M	19.0	+++	-	1.2	2.2	4.4	6.2	38.0
	M	18.0	+++	-	1.2	2.2	4.5	6.1	40.4
	M	19.0	+++	-	1.2	2.3	4.4	6.2	40.5
<i>Zonotrichia capensis</i> P		26.0	++	-	1.3	2.7		6.7	
	lm	21.0	+++	-	1.2	2.4	5.1	6.6	
		21.5	++-	-	1.3	2.4	5.0	6.7	
		28.0	---	-	1.2	2.2	5.5	6.5	37.4
		22.0	+	-	1.2	2.3	5.9	6.7	40.9
		19.5	---	-	1.3	2.3	5.4	6.8	41.4
		20.5	---	-	1.2	2.4	5.3	6.1	39.0
		19.5		-	1.3	2.2	5.3	6.3	40.2
		23.0	+++	-	1.3	2.4	5.4	6.6	42.0
		22.5	+++	-	1.3	2.4	5.5	6.7	38.6
		22.1	+++	-	1.4	2.3	5.6	6.6	41.5

1. b = body, t = tail, w = wing.

Weight in grams; measurements in centimeters; cloacal temperatures in degrees centigrades.

Presence of molt denoted by "+" and explicit absence by "-".

P= Parque Nacional Puracé; G= Estación de Bombeo Guamués; rest are from Reserva La Planada.

Table 2. Ectoparasites on Colombian birds.

Bird	Flea	Tick ¹	Chigger	Nose Mite	Mite ²	Lice ²
<i>Phaethornis symratorphorus</i>				+		
<i>Colibri coruscans</i>					+	
<i>Lafresnaya lafresnayi</i>				+	+	
<i>Coeligena wilsoni</i>						E
<i>Heliangulus exortis</i>				+		
<i>Haplophaedia lugens</i>				+	E	
<i>Metallura tyrianthina</i>				+	+	
<i>Aglaiocercus coelestis</i>						EN
<i>Semnormis ramphastinus</i>					+	
<i>Xiphocolactes promeropirhynchus</i>					E	
<i>Synallaxis azarae</i>						A
<i>Thripadectes virgaticeps</i>						AE
<i>Premnormis guttuligera</i>		R				
<i>Grallaria flavotincta</i>	+			+		
<i>Grallaricula flaviviridis</i>						AE
<i>Platyrinchus mystaceus</i>						A
<i>Mionectes olivaceus</i>					+	E
<i>Notiochelidon murina</i>						+
<i>Myadestes ralloides</i>					+	
<i>Catharus ustulatus</i>					+	A
<i>Turdus ignobilis</i>			+			E
<i>Vireo leucophrys</i>						A
<i>Dendroica fusca</i>					+	E
<i>Basileuterus tristriatus</i>						AE
<i>Diglossa albilatera</i>		L				E
<i>Diglossa cyanea</i>					+	
<i>Euphonia xanthogaster</i>	+					
<i>Tangara arthus</i>					+	AE
<i>Tangara vassorii</i>					E	
<i>Tangara nigroviridis</i>						E
<i>Iridosornis rufivertex</i>						E
<i>Anisognathus lachrymosus</i>		EL				
<i>Anisognathus flavinuchus</i>					AE	
<i>Thraupis cyanocephala</i>					AE	
<i>Atlapetes pallidinucha</i>	E					
<i>Atlapetes schistaceus</i>	E					
<i>Zonotrichia capensis</i>			+	+		

1_R = rump; L = eyelid;

E = ear

2_A = adult; N = nymph;

E = egg