

**TYPE STUDIES ON *FRULLANIA* SUBGENUS
METEORIOPSIS (HEPATICAE). II. ON *F. APOLLINARII*
AND *F. PARANENSIS***

**Estudios en ejemplares tipo del género *Frullania* subgénero
Meteoriopsis (Hepaticae). II. Sobre *F. apollinarii* y *F. paranensis***

JAIME URIBE-M.

*Instituto de Ciencias Naturales, Facultad de Ciencias, Universidad Nacional de Colombia.
Apartado 7495, Bogotá, D. C., Colombia. juribem@unal.edu.co.*

ABSTRACT

As a result of a redefinition of subgenus *Meteoriopsis* and the study of type material, *Frullania apollinarii* Steph. is transferred from this subgenus to subgenus *Thyopsiella* and *F. paranensis* to subgenus *Chonanthelia*.

Key words. Hepaticae, *Frullania* subgenus *Thyopsiella*, *Frullania* subgenus *Meteoriopsis*.

RESUMEN

Con base en los caracteres que definen al subgénero *Meteoriopsis* y el estudio de los ejemplares tipo se concluye que *Frullania apollinarii* Steph. debe ser transferida al subgénero *Thyopsiella* y *F. paranensis* al subgénero *Chonanthelia*.

Palabras clave. Hepaticae, *Frullania* subgénero *Thyopsiella*, *Frullania* subgénero *Meteoriopsis*.

INTRODUCTION

Stephani (1911) assigned a number of species of *Frullania* to subgen. *Meteoriopsis* based solely on the possession of a pendant habit; later, Uribe & Gradstein (2003) pointed out that the diagnostic characters of subgen. *Meteoriopsis* are the presence of the long-cylindrical lobule and auriculated leaf-lobe base, with auricles both antical and postical. Accordingly, several species previously assigned to subgen. *Meteoriopsis* were transferred to subgen. *Thyopsiella* by Uribe & Gradstein (2003). In this paper, *F. apollinarii* is transferred to subgen. *Thyopsiella* and *F. paranensis* to subgen. *Chonanthelia*.

Frullania apollinarii Steph., Spec. Hepat. 4: 602. 1911. Type. Colombia. Cordillera de Bogotá, 2700 m, 1905, *Apollinaire s. n.* (holotype, G 20609) Fig. 1

Frullania villosa Steph., Spec. Hepat. 4: 608. 1911. Type. Ecuador, 1896, *Mathan s. n.* (holotype, G 20627), **syn. nov.**

Plants of medium to large size, up to 15 cm long and 1.8 mm wide including leaves, dark reddish to black, irregularly pinnate. **Branches** frequent, *Frullania*-type, to 1 cm long. **Stems** to 220 µm wide in cross section. **Leaf-lobes** imbricate, convoluted around the stem when dry, spreading when wet, ovate, rounded,

asymmetrical, concave, 950-1475 x 1000-1525 μm , apex rounded, blunt, margin entire, dorsal base auriculate, arching over the stem, ventral base not auriculate. **Leaf-lobe cells:** apical cells 12-17 x 10-12 mm, median cells 20-27 x 10-15 mm, basal cells 30-40 x 20-25 mm, walls thick, sinuous, with trigones inconspicuous and intermediate thickenings. **Lobules** cylindrical to long cylindrical to clavate, contiguous and parallel to the stem, 320-420 x 120-170 mm. **Stylus** filiform, three cells long, with a terminal slime papilla. **Underleaves** ovate, oblong to obovate, contiguous, 1025-1075 x 720-800 mm, up to 4x wider than the stem, margin entire, undulate, recurved near the apex, bifid to 1/5 of their length, segments acute or blunt, base auriculate, auricles rather long, undulate, insertion line straight. **Branch appendages:** first branch underleaf divided to the base into a triangular, entire ventral segment and a saccate dorsal segment. **Androecia** lateral on short branches, capitate, 752 mm long, bracts in 4-5 series. **Gynoeceia** terminal on the stem, the bracts and bracteoles in three series, bracts bifid, and margin toothed. **Perianth** 1/3 exerted, oblong-elliptic, ca. 2000 μm long, smooth, terete, with an apical short beak.

Distribution. Colombia and Ecuador. Known only from the type localities.

Frullania apollinarii was erroneously included in subgenus *Meteoriopsis*, section *Acutifolia*. The study of the type specimen, however, revealed that this species has a rounded not acute leaf-lobe apex, (fig. 1C). This species is a genuine member of subgenus *Thyopsiella*, because of the presence of auricle only antical in base leaf-lobe.

Frullania apollinarii can be recognized by its obovate, rounded, semicordate stem leaf-lobes and auriculate underleaves, with long and undulate auricles. This species is related

to *F. paradoxa* Lehm. & Lindenb. but in the latter the underleaves lack auricles.

Frullania paranensis Steph., Spec. Hepat. 4: 607. 1911. Type: Brazil, Paraná, "Serrinha", 1904, *Dusén* 4395 (holotype, G 20627, isotype, NY) Fig. 2

Plants of medium to large size, 6-15 cm long, 1.5 mm wide including leaves, yellow, reddish-yellow to green-yellowish, projecting growth (not pendent). **Stems** 175-190 μm wide in cross section. **Branches** infrequent, *Frullania*-type, very large, to 10 cm long. **Leaf-lobes** distant to subimbricate, convoluted around the stem when dry, obliquely spreading when wet, ovate, symmetric, plane, 900-1350 x 1350-1600 μm , apex rounded, margin entire, base auriculate, cordate. **Leaf cells:** apical cells 7-10 x 7-10 mm, median cells 12-17 x 30-35 mm, and basal cells 22-25 x 27-35 mm; cells with walls thick, sinuous, with cordate trigones and without intermediate thickenings, basal cells with bulging trigones. **Lobules** explanate or laminate, 450-525 mm, contiguous to the stem. **Stylus** not seen. **Underleaves** ovate to subquadrate, distant, to 475-575 μm x 575-700 μm , up to 3x wider than the stem, margin toothed, with one or two lateral teeth, bifid to 1/2 of their length, segments blunt to acute, base without auricles, insertion line curved. **Branch appendages:** first branch underleaf divided to the base into a triangular, entire ventral segment and an explanate or canaliculate dorsal segment; first branch leaf transformed into two canaliculate segments. **Androecia** not seen. **Gynoeceia** on short lateral branches, bracts and bracteoles in two or three series, acute, entire. **Perianth** oblong, 5-8-keeled, smooth, narrowed to a short beak, 1 mm long, 850 μm wide.

Distribution. Known only from the type locality.




Fig. 1. *Frullania apollinari* Steph. A. Part of shoot, ventral view; B-C. Stem leaves; D-E. Underleaves; F. Ventral lobule; G. Androecium; H. Gynoecium with perianth. (A-G from type of *F. villosa* Steph.; H from type of *F. apollinari* Steph.)




Fig. 2. *Frullania paranensis* Steph. A. Part of shoot, ventral view; B-C. Stem leaves; D-E and F. Underleaves; G-H. Ventral lobules, canaliculate; I. Perianth. (All from the type of *F. paranensis* Steph.)

Frullania paranensis is a poorly known species; the absence of saccate lobules, an important character for placement in a particular subgenus, renders identification difficult; I consider, however, this species a genuine member of subgenus *Chonantheria*, based on the keeled perianth and the presence of canaliculate lobules. *F. paranensis* is probably conspecific with *F. lindmannii* Steph. both species has been reported from southern Brazil, Paraná and Rio Grande do Sul (Gradstein & Pinheiro da Costa, 2003).

ACKNOWLEDGMENTS

This paper is part of the doctoral dissertation "Monograph of *Frullania* subgenus *Meteoriopsis*", prepared under the direction of the Prof. Dr. S. R. Gradstein (Göttingen) in the Ph.D. program in Biology at the National University of Colombia. I thank the curators of the herbaria G and NY, for the loan of the type specimens. I would like to thank the National University of Colombia supporting my Ph.D. studies, to the Deutsche Akademischer Austauschdienst (DAAD) for the scholarship to work during the first half of 2002 in the Department of Systematic Botany

of the University of Göttingen. Special thanks are due to S. R. Gradstein for valuable discussions and comments on the manuscript, and to E. Forero and F. González for comments on the manuscript.

REFERENCES

- GRADSTEIN, S. R. & PINHEIRO DA COSTA, D. 2003. The Hepaticae and Anthocerotae of Brazil. Memoirs of New York Botanical Garden 87: 1-318.
- URIBE-M., J. & GRADSTEIN, S. R. 2003. Type studies on *Frullania* subgenus *Meteoriopsis* (Hepaticae). I. The lectotypification of the genus *Frullania*, *F.* subgen. *Thyopsiella* and *F.* subgen. *Meteoriopsis*, and some species transferred from subgen. *Meteoriopsis* to subgen. *Thyopsiella*. Cryptogamie, Bryologie 24(3): 193-207.
- STEPHANI, F. 1911. Species Hepaticarum, vol. IV. Acrogynae, (pars tertia). Georg & Cie., Geneva, Basel.

Recibido: 10/09/2003

Aceptado: 21/04/2004

