

Two records of anurophagy in *Leptodactylus macrosternum* from Cerrado, Brazil, with a review list of cases for the genus *Leptodactylus* (Anura: Leptodactylidae)

Dos nuevos registros de anurofagia para *Leptodactylus macrosternum* en el Cerrado, Brasil, con una lista de revisión de los casos para el género *Leptodactylus* (Anura: Leptodactylidae)

Diego G. Cavalheri ¹, Juan F. C. Carrillo ², Victoria dos Santos Souza ³, Diego J. Santana ⁴

- Received: 13/Jul/2021
- Accepted: 21/Jun/2022
- Online Publishing: 03/Oct/2022

Citation: Cavalheri DG, Carrillo JFC, Souza VS, Santana DJ. 2023. Two records of anurophagy in *Leptodactylus macrosternum* from Cerrado, Brazil, with a review list of cases for the genus *Leptodactylus* (Anura: Leptodactylidae). *Caldasia* 45(1):136-150. doi: <https://doi.org/10.15446/caldasia.v45n1.96789>

ABSTRACT

Anurophagy represents a small percentage of anurans' diet, however, the observation of this behavior is frequent and well-documented. Herein we reported two events of anurophagy of *Leptodactylus macrosternum* preying a newly metamorphosed *Physalaemus nattereri* and an adult *Boana punctata*, both in the municipality of Dois Irmãos do Buriti, Mato Grosso do Sul state, Western Brazil. Additionally, we summarized literature records of anurophagy in *Leptodactylus*. We found 99 records of anurophagy in different stages of development. *Physalaemus nattereri* and *Boana punctata* have never been reported as prey for *Leptodactylus* and this predation might be due to a combination of abundance and the high call activity during the breeding season. Anurophagy in *Leptodactylus* has more records for large-sized species, even in the larvae stage. Anurophagy occurrence in leptodactylids may be related to the size of the predator, habitat, and area diversity since more complex and diverse environments tend to present more anurophagy events.

Keywords: Batracophagy, diet, predation.

¹ Universidade Estadual Paulista, Instituto de Biociências, Letras e Ciências Exatas. Rua Cristóvão Colombo - de 1897/1898 ao fim, Jardim Nazareth, C. P. 15054000. São José do Rio Preto, Brazil. diego.cavalheri@unesp.br

² Universidade Federal de Mato Grosso do Sul, Maringá—Laboratório de Sistemática e Biogeografia de Anfíbios e Répteis, Instituto de Biociências. Av. Costa e Silva, sn, Universitário, C. P. 79070900. Campo Grande, Brazil. jfcuestas@gmail.com

³ Universidade Federal de Mato Grosso do Sul, Maringá—Laboratório de Sistemática e Biogeografia de Anfíbios e Répteis, Instituto de Biociências. Av. Costa e Silva, sn, Universitário, C. P. 79070900. Campo Grande, Brazil. vic.ssouza07@gmail.com

⁴ Universidade Federal de Mato Grosso do Sul, Maringá—Laboratório de Sistemática e Biogeografia de Anfíbios e Répteis, Instituto de Biociências. Av. Costa e Silva, sn, Universitário, C. P. 79070900. Campo Grande, Brazil. jose.santana@ufms.br

* Corresponding author



RESUMEN

La anurofagia representa un pequeño porcentaje de la dieta de los anuros, sin embargo, es un comportamiento común y bien documentado. Reportamos aquí dos eventos de anurofagia de *Leptodactylus macrosternum* depredando un juvenil metamorfo de *Physalaemus nattereri* y un adulto de *Boana punctata*, ambos eventos en el municipio de Dois Irmãos do Buriti, estado de Mato Grosso del Sur, Brasil. Adicionalmente, presentamos una revisión de los eventos de anurofagia del género *Leptodactylus*. Encontramos en total 99 registros de diferentes estados de desarrollo. *Physalaemus nattereri* y *Boana punctata* no habían sido reportadas como presas de *Leptodactylus*. Estas depredaciones pueden deberse a una combinación de la abundancia y la alta actividad de canto durante la reproducción. La anurofagia en leptodactílicos ha sido registrada mayormente en especies de gran tamaño, incluyendo en estados larvales. La ocurrencia de la anurofagia puede ser predicha por el tamaño del depredador, el hábitat y la diversidad del área, siendo que ambientes más complejos y de mayor diversidad tienden a presentar más eventos de anurofagia.

Palabras clave. Batracofagia, dieta, depredación.

Anurans play an important role in food webs in different habitats (Duellman and Trueb 1994). In the Neotropical region, they feed mainly on arthropods, and they are predated by fish, mammals, birds, reptiles, and invertebrates (Andrade et al. 2012, Aoki and Landgref-Filho 2013, Oliveira et al. 2014, Cortés-Gomez et al. 2015, Ferreira et al. 2017, Folt and Lapinski 2017, Ceron et al. 2019, Landgref-Filho et al. 2019, Cruz-Sáenz et al. 2020). Eventually, anurans can eat other frogs (i.e., anurophagy), and although this behavior represents a small percentage of anurans' diet, these events are well-documented (Toledo et al. 2007, Loebmann 2013, Oda et al. 2019, Caicedo-Martínez et al. 2021). Most anurans are known by their generalist diet, and their prey sizes may be related to the tongue and the mouth shape (Emerson 1985). For those reasons not only, the size is related to anurophagy, but also the skull shape and the habitat, especially in environments with high anuran abundance (Emerson 1985, Scott and Aquino 2005, Measey et al. 2015). Among Neotropical anurans, reports on anurophagy events in *Leptodactylus* are frequent (Sanabria et al. 2005, Toledo et al. 2007, Ávila-López et al. 2012, Sierra-Rueda and Acosta-Ortiz 2020). *Leptodactylus macrosternum* Miranda-Ribeiro, 1926, commonly called butter frog, is a large species (48.7–98.9 mm snout-vent length [SVL]) belonging to the *L. latrans* group (Magalhães et al. 2020). *Leptodactylus macrosternum* has a wide distribution in South America, from Santa Fé, Argentina, to northern Brazil and the Guiana Shield; also, it has been recorded in Trinidad and Tobago (Magalhães et al. 2020, Frost 2021).

Leptodactylus macrosternum reproductive activity occurs mainly during the rainy season (Prado et al. 2000, Camurugi et al. 2017). It is a generalist species with a foraging strategy between sit-and-wait and active search; its diet includes mainly coleopterans (Schaefer et al. 2006, Camurugi et al. 2017). Although anurophagy is well known for *Leptodactylus* species, most of the records are solely based on anecdotal observation, with the preyed anurans identified only to the order level (i.e., Anura) (e.g., Gallardo 1964, Maneyro et al. 2004, Solé et al. 2009, Camurugi et al. 2017).

Prey identification is fundamental to understanding trophic interactions (Ceron et al. 2020a), the role of anurans as energy and protein sources to other animals (Cortés-Gomes et al. 2015), the role of predators as regulators, and the possible drivers of anurophagy (Measey et al. 2015). Given the lack of specific records, herein we report two new anurophagy events of *Leptodactylus macrosternum* and provide a list of anurophagy cases for the genus *Leptodactylus*.

Both new records occurred at the edge of a permanent pond at Estância Crioula Farm, located at Dois Irmãos do Buriti municipality, Mato Grosso do Sul state, Western Brazil (20° 32' 07" South, 55° 33' 00" West). All specimens were collected under SISBIO license (SISBIO 49080-5), killed with lidocaine (5 %), fixed in 10 % formalin, preserved in 70 % Ethanol, and housed at Coleção Zoológica da Uni-

versidade Federal de Mato Grosso do Sul (ZUFMS-AMP). Additionally, to recover a list of anurophagy records by *Leptodactylus* species as predators from literature, we used the following keywords: “*Leptodactylus*”, “Anurophagy”, “Diet”, “Predation” and “Cannibalism” in “Google Scholar”. As a complement, we checked specialized journals such as *Herpetological Review* and *Herpetology Notes*, in the “Natural History” section. We included studies of diet, even with unidentified prey. The size category of species follows Haddad *et al.* (2013), who considered species as small (< 3 cm), medium (3 – 10 cm), and large (> 10 cm). We included reports of predation attempts or regurgitations and exclude those that not occurred in nature (e.g., Regnet and Loebmann 2017). When information about the locality and/or photographs allowed a more precise prey and predator identification, we included it (e.g., Faivovich *et al.* 2005, Heyer 2005, Frost *et al.* 2006, Narvaes and Rodrigues 2009, Duellman *et al.* 2016, Magalhães *et al.* 2020; Gazoni *et al.* 2021). Details about species names originally reported by authors are in [Supplementary file 1](#).

Our first report occurred on 11 Nov 2020, around 20:00h, when we collected a female of *L. macrosternum* (SVL=77.10 mm, ZUFMS-AMP13951, [Fig. 1a](#)) with an anuran leg out of its mouth. After dissection, we identified that the prey was a juvenile of *Physalaemus nattereri* (Steindachner, 1863) (SVL=13.62, ZUFMS-AMP13952, [Fig. 1b](#)). That night we recorded at the same pond the species *Boana raniceps* (Cope, 1862) in calling activity and saw individuals of *Leptodactylus podicipinus* (Cope, 1862) and *L. macrosternum*. The second report occurred at the edge of the same pond. This report occurred on 3 Mar 2021, around 20:00h. During that night *Boana punctata* (Schneider, 1799), *B. raniceps*, *Dendropsophus minutus* (Peters, 1872), and *Pithecopus azureus* (Cope, 1862) were calling; *L. macrosternum* and *Leptodactylus podicipinus* (Cope, 1862) were also observed although no calling behavior was recorded in these species. We observed and handled a female of *L. macrosternum* (SVL=79.21mm, ZUFMS-AMP14659, [Fig. 1c](#)) with an adult of *Boana punctata* (SVL=35.29 mm, ZUFMS-AMP14649, [Fig. 1d](#)) in its mouth. After we stored the *L. macrosternum* in a plastic bag, the *B. punctata* was regurgitated.

We found 66 references reporting events of anurophagy for a total of 99 records (101 including both records described here) ([Table 1](#)). Large-size species represent 64.35 % of the total records, Medium-size species, represent the others 35.65 % and we were not able to find anurophagy in Small-size leptodactylids.

The four-eyed frog, *P. nattereri*, has inguinal macroglands that can be used against predators during deimatic display (Lenzi-Mattos *et al.* 2005). This species has already been reported as prey for other species of *Leptodactylus* (Vaz-Silva *et al.* 2003, França *et al.* 2004), however, in two events, with *L. luctator* (Hudson, 1892), the toxin secreted by *P. nattereri* was efficient to avoid predation (Rodrigues and Filho 2004). In the present report, the preyed individual was a juvenile, newly metamorphosed, which may suggest that this is a less-effective defensive method at this life stage such as occur in *Rhinella arenarum* (Hensel, 1867) (Regueira *et al.* 2016). During the night of the second event of anurophagy reported here, many *B. punctata* were exhibiting calling behavior. This species selects different vegetation strata to call and exhibits territorial behavior, moving 100 cm around the calling site (Brunetti *et al.* 2014). The combination of the high abundance and the increase in calling activity and mobility during the breeding season of these species might result in occasional encounters and subsequent predation of *L. macrosternum* upon *B. punctata*. This is the first report of *Leptodactylus* preying on *B. punctata*.

Although some large frogs, such as bufonids, are negligible predators of anurans (Measey *et al.* 2015), in leptodactylids the majority of anurophagy events are displayed by large-size frogs ([Table 1](#)). In medium size species, *L. macrosternum* represents half of the anurophagy records and it is the largest species of its category (Maximum SVL=9.89cm). Trophic ecology studies are paramount to follow temporal variations and the diversity of interactions (Ceron *et al.* 2020a, 2020b). For that purpose, we need proper identification of both, predator, and prey species. Our literature review on anurophagy by *Leptodactylus* leaps further in the anuran trophic ecology, which includes new records.




Figure 1. Individuals from the two anurophagy events reported. **a.** *Leptodactylus macrosternum*, **b.** juvenile of *Physalaemus nattereri* preyed on by the specimen “a”, **c.** *L. macrosternum*, **d.** *Boana punctata* found inside the mouth of the specimen “c”. Scale bar= 20mm.

Table 1. Events of anurophagy of *Leptodactylus* species and their preys. AP = Amapá, BA = Bahia, CE = Ceará, ES = Espírito Santo, GO = Goiás, MG = Minas Gerais, MS = Mato Grosso do Sul, MT = Mato Grosso, PE = Pernambuco, PI = Piauí, PR = Paraná, RJ = Rio de Janeiro, RN = Rio Grande do Norte, RS = Rio Grande do Sul, SC = Santa Catarina, SE = Sergipe, SP = São Paulo.

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
<i>Leptodactylus cunicularius</i> (0.99)	Medium	Leptodactylidae	<i>Leptodactylus cunicularius</i> Sazima and Bokermann, 1978	Ouro Preto - MG	Brazil	Pirani et al. 2010
<i>Leptodactylus furnarius</i> (0.99)	Medium	Hylidae	<i>Scinax fuscomarginatus</i> (Lutz, 1925)	Botucatu - SP	Brazil	Rolim et al. 2009
<i>Leptodactylus fuscus</i> (2.97)	Medium	Hylidae	<i>Boana</i> sp.	Villavicencio - Meta	Colombia	Sierra-Rueda and Acosta-Ortiz 2020
		Leptodactylidae	<i>Leptodactylus caatingae</i> Heyer and Juncá, 2003	Jequié - BA	Brazil	Novaes-e-Fagundes and Zina 2017
		-	Unidentified Anura	southeastern Brazil	Brazil	Araújo et al. 2007
<i>Leptodactylus labyrinthicus</i> (14.85)	Large	Hylidae	<i>Boana albopunctata</i>	-	-	Toledo et al. 2007
			<i>Boana faber</i> (Wied-Neuwied, 1821)	-	-	Toledo et al. 2007
			<i>Dendropsophus minutus</i>	Uberlândia - MG	Brazil	França et al. 2004
			<i>Dendropsophus minutus</i>	Uberlândia - MG	Brazil	Silva et al. 2005
			<i>Scinax</i> cf. <i>similis</i>	Uberlândia - MG	Brazil	França et al. 2004
			<i>Scinax</i> sp.	Uberlândia - MG	Brazil	França et al. 2004
			<i>Trachycephalus typhonius</i> (Linnaeus, 1758)	Corumbá - MS	Brazil	Landgraf-Filho et al. 2012
		Leptodactylidae	<i>Leptodactylus labyrinthicus</i>	Uberlândia - MG	Brazil	França et al. 2004

(Continúa)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
			<i>Leptodactylus labyrinthicus</i>	Uberlândia - MG	Brazil	Silva et al. 2005
			<i>Physalaemus cuvieri</i> Fitzinger, 1826	Águas de Santa Bárbara - SP	Brazil	Diaz-Ricaurte and Marques 2021
			<i>Physalaemus cuvieri</i>	Uberlândia - MG	Brazil	Silva et al. 2005
			<i>Physalaemus marmoratus</i> (Reinhardt and Lütken, 1862)	Uberlândia - MG	Brazil	Silva et al. 2005
			<i>Physalaemus nattereri</i>	Caldas Novas - GO	Brazil	Vaz-Silva et al. 2003
			<i>Physalaemus nattereri</i>	Uberlândia - MG	Brazil	França et al. 2004
			<i>Pseudopaludicola saltica</i> (Cope, 1887)	Uberlândia - MG	Brazil	França et al. 2004
<i>Leptodactylus latrans</i> (13.86)	Large	Bufonidae	<i>Rhinella crucifer</i> (Wied-Neuwied, 1821)	Uruçuca - BA	Brazil	Rojas-Padilha et al. 2018
			<i>Rhinella ornata</i> (Spix, 1824)	Ubatuba - SP	Brazil	Bovo et al. 2014
		Hylidae	<i>Boana albomarginata</i> (Spix, 1824)	Anchieta - ES	Brazil	Teixeira and Vrcibradic 2003
			<i>Boana pardalis</i> (Spix, 1824)	Espera Feliz - MG	Brazil	Heitor et al. 2012
			<i>Boana raniceps</i>	Capela - SE	Brazil	Ferreira et al. 2011
			<i>Scinax fuscovarius</i> (Lutz, 1925)	Vassouras - RJ	Brazil	Bezerra et al. 2015
			<i>Scinax x-signatus</i> (Spix, 1824)	São Lourenço da Mata - PE	Brazil	Santos 2009
			<i>Trachycephalus mesophaeus</i> (Hensel, 1867)	Igrapiúma - BA	Brazil	Mendes et al. 2012

(Continua)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
		Leptodactylidae	<i>Leptodactylus latrans</i>	Anchieta - ES	Brazil	Teixeira and Vrcibradic 2003
			<i>Physalaemus crombiei</i> Heyer and Wolf, 1989	Anchieta - ES	Brazil	Teixeira and Vrcibradic 2003
			<i>Physalaemus cuvieri</i>	São Lourenço da Mata - PE	Brazil	Santos 2009
		Ranidae	<i>Lithobates catesbeianus</i> (Shaw, 1802)	Vieira - MG	Brazil	Silva and Filho 2009
		-	Unidentified Anura	Ilhéus - BA	Brazil	Solé <i>et al.</i> 2009
		-	Unidentified Anura	Anchieta - ES	Brazil	Teixeira and Vrcibradic 2003
<i>Leptodactylus luctator</i> (15.84)	Large	Bufo	<i>Rhinella mirandariibeiroi</i> (Gallardo, 1965)	Uberlândia - MG	Brazil	França <i>et al.</i> 2004
			<i>Rhinella ornata</i>	São Paulo - SP	Brazil	Capela <i>et al.</i> 2020
			<i>Rhinella</i> sp.	Uberlândia - MG	Brazil	França <i>et al.</i> 2004
			Unidentified Bufonidae	San Juan - Dep. Zonda	Argentina	Sanabria <i>et al.</i> 2005
		Hylidae	<i>Boana faber</i>	Apiaí - SP	Brazil	Pombal Jr., 2007
			<i>Boana faber</i>	Serra do Japi - SP	Brazil	Haddad and Sazima 1992
			Unidentified Hylidae	Uberlândia - MG	Brazil	França <i>et al.</i> 2004
		Leptodactylidae	<i>Leptodactylus furnarius</i>	Uberlândia - MG	Brazil	França <i>et al.</i> 2004
			<i>Leptodactylus luctator</i>	Unai - MG	Brazil	Kokubum and Rodrigues 2005

(Continúa)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
			<i>Physalaemus biligonigerus</i> (Cope, 1861)	Tucumán	Argentina	Gallardo 1964
			<i>Physalaemus centralis</i> Bokermann, 1962	Araguari - MG	Brazil	Heyer and Giaretta 2009
			<i>Physalaemus marmoratus</i>	Paraúna - GO	Brazil	Santos et al. 2021
			<i>Physalaemus nattereri</i>	Uberlândia - MG	Brazil	Rodrigues and Filho 2004
		-	Unidentified Anura	Apiá - SP	Brazil	Pombal Jr 2007
		-	Unidentified Anura	Caçapava do Sul - RS	Brazil	Pazinato et al. 2011
		-	Unidentified Anura	Arroyo Espinas - Maldonado	Uruguay	Maneyro et al. 2004
<i>Leptodactylus macrosternum</i> (18.81)	Medium	Bufo	<i>Rhinella major</i> (Müller and Hellmich, 1936)	Santana - AP	Brazil	Sousa et al. 2016
		Hylidae	<i>Boana albomarginata</i>	Macaíba - RN	Brazil	Baracho et al. 2013
			<i>Boana punctata</i>	Aquidauana - MS	Brazil	This study
			<i>Dendropsophus nanus</i> (Boulenger, 1889)	Corguinho - MS	Brazil	Carrillo et al. 2019
			<i>Pseudis platensis</i> Gallardo, 1961	Diamante do Norte - PR	Brazil	Oda et al. 2016
			<i>Scinax nasicus</i> Cope, 1862	Selvíria - MS	Brazil	Queiroz et al. 2019
		Leptodactylidae	<i>Leptodactylus elenae</i> Heyer, 1978	Estirão Comprido - MT	Brazil	Moreira and Smaniotto 2017
			<i>Leptodactylus fuscus</i>	Macapá - AP	Brazil	Silva et al. 2018
			<i>Leptodactylus fuscus</i>	Acreúna - GO	Brazil	Costa and Trevelin 2020

(Continúa)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
			<i>Leptodactylus macrosternum</i>	Santana - AP	Brazil	Sousa et al. 2016
			<i>Leptodactylus podicipinus</i>	Corumbá - MS	Brazil	Carrillo et al. 2019
			<i>Leptodactylus troglodytes</i>	Serra Negra do Norte - RN	Brazil	Sales et al. 2015
			<i>Physalaemus nattereri</i>	Aquidauana - MS	Brazil	This study
			<i>Physalaemus centralis</i>	Jardim - MS	Brazil	Costa-Pereira et al. 2015
			<i>Physalaemus cuvieri</i>	Selvíria - MS	Brazil	Queiroz et al. 2019
		-	Unidentified Anura	Corrientes - Corrientes	Argentina	Dure 1999
		-	Unidentified Anura	Macaíba - RN	Brazil	Camurugi et al. 2017
		-	Unidentified Anura	Miranda - MS	Brazil	Piatti and Souza 2011
<i>Leptodactylus melanotus</i> (2.97)	Medium	Hylidae	<i>Dendropsophus ebraccatus</i> (Cope, 1874)	San Luis - Limón	Costa Rica	Hoffmann 2006
			<i>Tlalocohyla smithii</i> (Boulenger, 1902)	Minatitlán - Colima	Mexico	Ávila-Lopez et al. 2012
			<i>Lithobates vaillanti</i> (Brocchi, 1877)	Barra del Colorado	Costa Rica	Lewis et al. 2008
<i>Leptodactylus mystaceus</i> (0.99)	Medium	Bufoidea	<i>Atelopus hoogmoedi</i> Lescure, 1974	Laranjal do Jarí - AP	Brazil	Pinto and Costa-Campos 2015
<i>Leptodactylus natalensis</i> (3.96)	Medium	Hylidae	<i>Dendropsophus aff. decipiens</i>	Guarapari - ES	Brazil	Ferreira et al. 2007
			<i>Scinax argyreornatus</i> (Miranda-Ribeiro, 1926)	Guarapari - ES	Brazil	Ferreira et al. 2007
		Leptodactylidae	<i>Physalaemus</i> sp.	Guarapari - ES	Brazil	Ferreira et al. 2007

(Continua)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
		-	Unidentified Anura	São Lourenço da Mata - PE	Brazil	Santos et al. 2004
<i>Leptodactylus paranaru</i> (0.99)	Large	Hylidae	<i>Boana bischoffi</i> (Boulenger, 1887)	Azambuja - SC	Brazil	Peres et al. 2017
<i>Leptodactylus payaya</i> (0.99)	Large	Hylidae	<i>Boana ranceps</i>	Barbalha - CE	Brazil	Chaves et al. 2012
<i>Leptodactylus pentadactylus</i> (1.98)	Large	Bufo	<i>Rhinella margaritifera</i> (Laurenti, 1768)	Serra do Návio - AP	Brazil	Santana et al. 2018
		Hylidae	<i>Boana rosenbergi</i> (Boulenger, 1898)	-	-	Kluge 1981
<i>Leptodactylus podicipinus</i> (2.97)	Medium	Bufo	<i>Rhinella diptycha</i> (Cope, 1862)	Corumbá - MS	Brazil	Ceron et al. 2018
			<i>Rhinella major</i>	Macapá - AP	Brazil	Santos et al. 2018
<i>Leptodactylus rhodonotus</i> (0.99)		Bufo	<i>Rhinella marina</i>	Mascoitania reserve - Peru	Peru	Thomas et al. 2017
			<i>Rhinella mirandaribeiroi</i>	Pontalina - GO	Brazil	Guimarães et al. 2004
<i>Leptodactylus savagei</i> (0.99)	Large	Leptodactylidae	<i>Engystomops pustulosus</i> (Cope, 1864)	Norcasia - Caldas	Colombia	González-Duran and Gutiérrez-Cárdenas 2013
<i>Leptodactylus troglodytes</i> (0.99)	Medium	Leptodactylidae	<i>Leptodactylus troglodytes</i>	Natal - RN	Brazil	Jorge and Freire 2011
<i>Leptodactylus vastus</i> (8.91)	Large	Bufo	<i>Rhinella mirandaribeiroi</i>	Parque Nacional de Sete Cidades - PI	Brazil	Cardoso and Sazima 1977
		Hylidae	<i>Boana albomarginata</i>	Capela - SE	Brazil	Santana et al. 2012
			<i>Boana albopunctata</i>	Parque Nacional de Sete Cidades - PI	Brazil	Cardoso and Sazima 1977
			<i>Boana faber</i>	Capela - SE	Brazil	Neto et al. 2015
		Leptodactylidae	<i>Leptodactylus latrans</i>	Capela - SE	Brazil	Santana et al. 2012

(Continúa)

Predator (% of observations)	Size	Family of Prey	Prey	Locality	Country	Reference
			<i>Leptodactylus natalensis</i>	São Lourenço da Mata - PE	Brazil	Santos 2009
			<i>Leptodactylus vastus</i>	Mata de São João - BA	Brazil	Guimarães et al. 2015
			<i>Physalaemus albifrons</i> (Spix, 1824)	Aiuaba - CE	Brazil	Teles et al. 2015
			<i>Physalaemus cuvieri</i>	São Lourenço da Mata - PE	Brazil	Santos 2009
<i>Leptodactylus</i> sp. (gr. <i>latrans</i>) (5,94)	Large	Bufonidae	<i>Rhinella crucifer</i>	-	Brazil and Argentina	Gallardo 1964
			<i>Rhinella</i> sp. (gr. <i>granulosa</i>)	-	Brazil and Argentina	Gallardo 1964
		Hylidae	<i>Boana albomarginata</i>	-	-	Toledo et al. 2007
			<i>Pseudis minuta</i> Günther, 1858	-	Brazil and Argentina	Gallardo 1964
			<i>Scinax squallirostris</i> (Lutz, 1925)	-	Brazil and Argentina	Gallardo 1964
			<i>Boana</i> sp. (gr. <i>pulchella</i>)	-	Brazil and Argentina	Gallardo 1964

AUTHOR CONTRIBUTIONS

J.F.C. Carrillo and D. G. Cavalheri designed the research structure and D.G. Cavalheri and V.S. Souza collected and drafted the manuscript. J.F.C. Carrillo and D. J. Santana substantially edited and improved the manuscript.

ACKNOWLEDGEMENTS

We thank Vanessa Stavis from Estância Crioula for all the logistic support. DGC and JFCC thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES) for the scholarship (Funding code 001). DJS thanks Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq processes 404239/2021-8, 402519/2016-7) and Fundação de Apoio ao Desenvolvimento do Ensino, Ciência e Tecnologia do Estado de Mato Grosso do Sul (FUNDECT process 054/2017) for financial support. DJS also thanks CNPq for his research fellowship (CNPq 309420/2020-2).

CONFLICT OF INTEREST

The authors do not report any potential conflict of interest.

LITERATURE CITED

- Andrade EB, Lima Júnior TB, Leite Júnior JMA, Leite JRSA. 2012. Predation by native fish and feeding by crab species on *Leptodactylus macrosternum* Miranda-Ribeiro, 1926 (Anura: Leptodactylidae) in northeastern, Brazil. *Herpetol. Notes* 5:173–175.
- Aoki C, Landgraf-Filho P. 2013. Predation on *Leptodactylus chaquensis* (Anura: Leptodactylidae) by the whistling heron *Syrigma sibilatrix* (Ciconiiformes: Ardeidae) in Central Brazil. *Herpetol. Notes* 6:261–262.
- Araújo MS, Reis SF, Giaretta AA, Machado G, Bolnick DI. 2007. Intrapopulation diet variation in four frogs (Leptodactylidae) of the Brazilian Savannah. *Copeia* 2007(4):855–865. doi: [https://doi.org/10.1643/0045-8511\(2007\)7\[855:IDVIF2\]2.0.CO;2](https://doi.org/10.1643/0045-8511(2007)7[855:IDVIF2]2.0.CO;2)
- Ávila-López OA, Reyes-Velasco J, Hermsillo-López A. 2012. *Leptodactylus melanotus* (Sabinal Frog). *Prey. Herpetol. Rev.* 43(1):122–123.
- Baracho EBO, Queiroz MHC, Mângia S. 2013. Predation of *Hypsiboas albomarginatus* (Spix, 1824) (Anura: Hylidae) by Miranda's white-lipped frog *Leptodactylus macrosternum* (Miranda-Ribeiro, 1926) (Anura: Leptodactylidae). *Herpetol. Notes* 6:599–601.
- Bezerra AM, Ruggeri J, Carvalho-e-Silva SP. 2015. *Scinax fuscovarius* (Snouted Treefrog). *Predation. Herpetol. Rev.* 46(2):239.
- Bovo RP, Bandeira LN, Condez TH. 2014. *Rhinella ornata* (Mexican Spadefoot). Predation by *Leptodactylus latrans*. *Herpetol. Rev.* 45(1):115.
- Brunetti AE, Taboada C, Faivovich J. 2014. The reproductive biology of *Hypsiboas punctatus* (Anura: Hylidae): male territoriality and the possible role of different signals during female choice. *Salamandra* 50(4):215–224.
- Caicedo-Martínez LS, Escobar-Lasso S, Zuluaga-Isaza JC, Londoño-Quiceno C, Orrego-Meza JG, Rivera-Pérez JM. 2021. Review of post-metamorphic frog-eat-frog predation, with a description of a new cases of anurophagy. *Food Webs* 27:1–2. doi: <https://doi.org/10.1016/j.fooweb.2021.e00191>
- Camurugi F, Magalhães FM, De Queiroz MHC, Pereira TCSO, Tavares-Bastos L, Lopes-Marinho ES, Neves JMM, Garda AA. 2017. Reproduction, sexual dimorphism, and diet of *Leptodactylus chaquensis* (Anura, Leptodactylidae) in Northeastern Brazil. *Herpetol. Conserv. Bio.* 12(2):498–508.
- Capela DJV, Struett MM, Leivas PT. 2020. Predation attempt of *Rhinella ornata* (Spix, 1824) (Anura, Bufonidae) by *Leptodactylus cf. latrans* (Anura, Leptodactylidae) in the Atlantic Forest, Brazil. *Herpetol. Notes* 13:11–13.
- Cardoso AJ, Sazima I. 1977. Batracofagia na fase adulta e larvária da rã pimenta, *Leptodactylus labyrinthicus* (Spix, 1824) - Anura, Leptodactylidae. *Cien. Cult.* 29(10):1130–1132.
- Carrillo JFC, Ferreira VG, Santana DJ. 2019. Batrachophagy by *Leptodactylus chaquensis* (Anura: Leptodactylidae) in the Brazilian Cerrado and Pantanal. *Herpetol. Notes* 12:261–263.
- Ceron K, Moroti MDT, Benício RA, Balboa P, Marçola Y, Pereira LB, Santana DJ. 2018. Diet and first report of batrachophagy in *Leptodactylus podicipinus* (Anura: Leptodactylidae). *Neotrop. Biodivers.* 4(1):70–74. doi: <https://doi.org/10.1080/23766808.2018.1467173>
- Ceron K, Oliveira-Santos LGR, Souza CS, Mesquita DO, Caldas FLS, Araujo AC, Santana DJ. 2019. Global patterns in anuran-prey networks: structure mediated by latitude. *Oikos* 128(11):1537–1548. doi: <https://doi.org/10.1111/oik.06621>
- Ceron K, Santana DJ, Lucas EM, Zocche JJ, Provete DB. 2020a. Climatic variables influence the temporal dynamics of an anuran metacommunity in a non-stationary way. *Ecol. Evol.* 10(11):4630–4639. doi: <https://doi.org/10.1002/ece3.6217>
- Ceron K, Santana DJ, Valente-Neto F. 2020b. Seasonal patterns of ecological uniqueness of anuran metacommunities along different ecoregions in Western Brazil. *PLoS ONE.* 15: doi: <https://doi.org/10.1371/journal.pone.0239874>
- Chaves LS, Veja ESF, Moura RCM, Muniz SLS, Moura GJB. 2012. *Leptodactylus latrans* (Caçote Frog). *Predation. Herpetol. Rev.* 43(3):464–465.
- Cortés-Gomez AM, Ruiz-Agudelo CA, Valencia-Aguilar A, Ladle RJ. 2015. Ecological functions of neotropical amphibians and reptiles: A review. *Univ. Sci.* 20(2): 229–245. doi: <https://doi.org/10.11144/Javeriana.SC20-2.efna>

- Costa WP, Trevelin CC. 2020. Congeneric predation of *Leptodactylus fuscus* (Schneider, 1799) by *Leptodactylus chaquensis* Cei, 1950 (Anura, Leptodactylidae). *Herpetol. Notes* 13:109–111.
- Costa-Pereira R, Sugai JLMM, Duleba S, Sugai LSM, Terra JS, Souza FL. 2015. Predation on *Physalaemus centralis* by the Chaco Frog *Leptodactylus chaquensis*. *Herpetol. Notes* 8:345–346.
- Cruz-Sáenz D, Lazcano D, Ríos-Martínez JO, García-Salas JA, Rodríguez-López A, Wilson LD. 2020. Notes on the Herpetofauna of Western Mexico 24: Predation on *Lithobates neovolcanicus* (Hillis & Frost, 1985) by *Quiscalus mexicanus* (Gmelin, 1788) in the Gardens of Guadalajara Zoo, Jalisco, Mexico. *Bull. Chic. Herpetol. Soc.* 55(8):153–157.
- Diaz-Ricaurte JC, Marques PR. 2021. *Leptodactylus labyrinthicus* (Pepper Frog). *Batrachophagy*. *Herpetol. Rev.* 52(1):117.
- Duellman WE, Marion AB, Hedges SB. 2016. Phylogenetics, classification, and biogeography of the treefrogs (Amphibia: Anura: Arboranae). *Zootaxa* 4104(1):1–109. doi: <https://doi.org/10.11646/zootaxa.4104.1.1>
- Duellman WE, Trueb L. 1994. *Biology of Amphibians*. Baltimore and London: The Johns Hopkins University Press.
- Dure MI. 1999. *Leptodactylus chaquensis* (NCN). *Diet*. *Herpetol. Rev.* 30(2):92.
- Ersparmer V, Glasser A. 1960. The pharmacological actions of (m-hydroxyphenethyl) trimethylammonium (leptodactiline). *Br. J. Pharmacol.* 15(14):14–22. doi: <https://doi.org/10.1111/j.1476-5381.1960.tb01204.x>
- Emerson SB. 1985. Skull shape in frogs – correlation with diet. *Herpetologica* 41(12):177–188.
- Faivovich J, Haddad CFB, Garcia PCA, Frost DR, Campbell JA, Wheeler WC. 2005. Systematic review of the frog family Hylidae, with special reference to Hylinae: Phylogenetic analysis and taxonomic revision. *Bull. Am. Mus. Nat. Hist.* (294):1–227. doi: [https://doi.org/10.1206/0003-0090\(2005\)294\[0001:SR0TFF\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2005)294[0001:SR0TFF]2.0.CO;2)
- Ferreira RB, Dantas RB, Teixeira RL. 2007. Reproduction and ontogenetic diet shifts in *Leptodactylus natalensis* (Anura, Leptodactylidae) from southeastern Brazil. *Bol. Mus. Biol. Mello Leitão* 22:45–55.
- Ferreira AS, Faria RG, Silva IRS, Schettino SC. 2011. *Hypsiboas raniceps* (Chaco Treefrog). *Predation*. *Herpetol. Rev.* 42(2):86–87.
- Ferreira VL, Leuzinger L, Soares P, Leão TRF, Himmelstein J. 2017. Predation of *Leptodactylus chaquensis* Cei, 1950 (Amphibia: Anura) by the Neotropical Otter, *Lontra longicaudis* (Olfers, 1818) (Carnivora: Mustelidae), in the Southern Pantanal of Brazil. *Herpetol. Notes* 10:111–114.
- Folt B, Lapinski W. 2017. New observations of frog and lizard predation by wandering and orb-weaver spiders in Costa Rica. *Phyllomedusa* 16(2):269–277. <https://doi.org/10.11606/issn.2316-9079.v16i2p269-277>
- França LF, Facure KG, Giaretta AA. 2004. Trophic and Spatial Niches of two Large-sized Species of *Leptodactylus* (Anura) in Southeastern Brazil. *Stud. Neotrop. Fauna Environ.* 39(3):243–248. doi: <https://doi.org/10.1080/01650520400007330>
- Frost DR. 2021. *Amphibian Species of the World: an Online Reference*. Version 6.1 [last accessed: 01 Sep 2022]. <https://amphibiansoftheworld.amnh.org/>
- Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, de Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler P, Drewes RC, Nussbaum RA, Lynch JD, Green DM, Wheeler WC. 2006. The Amphibian Tree of Life. *Bull. Am. Mus. Nat. Hist.* 2006(297):1–291. doi: [https://doi.org/10.1206/0003-0090\(2006\)297\[0001:TATOL\]2.0.CO;2](https://doi.org/10.1206/0003-0090(2006)297[0001:TATOL]2.0.CO;2)
- Gallardo JM. 1964. Consideraciones sobre *Leptodactylus ocellatus* (L.) (Amphibia, Anura) y especies aliadas. *Physis* (68):373–384.
- Gazoni T, Lyra ML, Ron SR, Strüssmann C, Baldo D, Narimatsu H, Pansonato A, Schneider RG, Giaretta AA, Haddad CFB, Parise-Maltempi PP, Carvalho TR. 2021. Revisiting the systematics of the *Leptodactylus melanonotus* group (Anura: Leptodactylidae): Redescription of *L. petersii* and revalidation of its junior synonyms. *Zool. Anz.* 290: 117–134.
- González-Duran G, Gutiérrez-Cárdenas PDA. 2013. *Leptodactylus savagei* (Savage's Thin-toed Frog). *Prey*. *Herpetol. Rev.* 44(2):294–295.
- Guimarães LD, Pinto RM, Juliano RF. 2004. *Bufo granulatus* (NCN). *Predation*. *Herpetol. Rev.* 35(3):259.
- Guimarães FPBB, Marques R, Tinôco MS. 2015. *Leptodactylus vastus* (Northeastern Pepper Frog). *Cannibalism*. *Herpetol. Rev.* 46(1):74–75.
- Haddad CFB, Sazima I. 1992. Anfíbios anuros da Serra do Japi. En: Morellato LPC, editor. *História natural da Serra do Japi: Ecologia e Preservação de uma área florestal no sudeste do Brasil*. Campinas: Unicamp and Fapesp. p. 188–211.
- Haddad CFB, Toledo LF, Prado CPA, Loebmann D, Gasparini JL, Sazima I. 2013. *Guia dos anfíbios da Mata Atlântica: Diversidade e Biologia*. São Paulo: Anolisbooks.
- Heitor RC, Lacerda JVA, Silva ET, Peixoto MA, Eloi RG. 2012. Predation of *Hypsiboas pardalis* (Anura, Hylidae) by the butterfly frog *Leptodactylus cf. latrans* (Anura, Leptodactylidae), in municipality of Espera Feliz, State of Minas Gerais, southeastern Brazil. *Herpetol. Notes* 5:23–25.
- Heyer WR. 2005. Variation and taxonomic clarification of the large species of the *Leptodactylus pentadactylus* species group (Amphibia: Leptodactylidae) from Middle America, Northern South America, and Amazonia. *Arq. Zool.* 37(3):269–348. doi: <https://doi.org/10.11606/issn.2176-7793.v37i3p269-348>
- Heyer WR, Giaretta AA. 2009. Advertisement calls, notes on natural history, and distribution of *Leptodactylus chaquensis* (Amphibia: Anura: Leptodactylidae) in Brazil. *Proc. Biol. Soc. Wash.* 122(3):292–305. doi: <https://doi.org/10.2988/08-42.1>

- Hoffmann H. 2006. Observations on behaviour and parental care of *Leptodactylus melanonotus* (Hallowell) in Costa Rica. *Salamandra* 42:109–116.
- Jorge JS, Freire EMX. 2011. *Leptodactylus troglodytes* (Caçote). Larval cannibalism. *Herpetol. Rev.* 42(4):585.
- Kluge AG. 1981. The Life History, Social Organization, and Parental Behavior of *Hyla rosenbergi* Boulenger, a Nest-Building Gladiator Frog. *Misc. Publ. Mus. Zool. Univ. Mich.* (160):1–170.
- Kokubum MNC, Rodrigues AP. 2005. *Leptodactylus ocellatus* (Rã-Manteiga). Cannibalism. *Herpetol. Rev.* 36(3):303.
- Landgraf-Filho P, Aoki C, Sousa DLK, Souza EO, Brandão RB, Ávila RW, Oda FH. 2019. Escape or be preyed: New records and current knowledge on predators of pseudinae frogs (Anura: Hylidae) in South America. *Acta Biol. Colomb.* 24(2):397–402. doi: <https://doi.org/10.15446/abc.v24n2.74650>
- Landgraf-Filho P, Uetanabaro M, Mori IK, Hokama DL, Aoki C. 2012. *Trachycephalus typhonius* (Milky Treefrog). Predation. *Herpetol. Rev.* 43(3):472.
- Lenzi-Mattos R, Antoniazzi MM, Haddad CFB, Tambourgi DV, Rodrigues MT, Jared C. 2005. The inguinal macroglands of the frog *Physalaemus nattereri* (Leptodactylidae): Structure, toxic secretion and relationship with deimatic behaviour. *J. Zool.* 266(4):385–394. doi: <https://doi.org/10.1017/S095283690500703X>
- Lewis TR, Ryall C, Laduke TC. 2008. *Leptodactylus melanonotus* (Sabinal Frog). Diet. *Herpetol. Rev.* 39(1):79.
- Loebmann D. 2013. A record of *Trachycephalus typhonius* (Linnaeus, 1758) preying upon *Dendropsophus soaresi* (Caramaschi & Jim, 1983) (Anura, Hylidae). *Herpetol. Notes* 6:275–276.
- Magalhães FM, Lyra ML, Carvalho TR, Baldo D, Bruschetti F, Burella P, Colli GR, Gehara MC, Giaretta AA, Haddad CFB, Langone JA, López JA, Napoli MF, Santana DJ, De Sá RO, Garda AA. 2020. Taxonomic review of South American Butter Frogs: Phylogeny, Biogeographic Patterns, and species delimitation in the *Leptodactylus latrans* Species Group (Anura: Leptodactylidae). *Herpetol. Monogr.* 34(1):131–177. doi: <https://doi.org/10.1655/0733-1347-31.4.131>
- Maneyro R, Naya DE, Rosa I, Canavero A, Camargo A. 2004. Diet of the South American frog *Leptodactylus ocellatus* (Anura, Leptodactylidae) in Uruguay. *Iheringia, Ser. Zool.* 94(1):57–61. doi: <https://doi.org/10.1590/S0073-47212004000100010>
- Measey GJ, Vimercati G, De Villiers FA, Mokhatla MM, Davies SJ, Edwards S, Altwegg R. 2015. Frog eat frog: Exploring variables influencing anurophagy. *PeerJ.* 2015:1–16. doi: <https://doi.org/10.7717/peerj.1204>
- Mendes CVM, Ruas DS, Solé M. 2012. Predation attempt of *Trachycephalus mesophaeus* (Hylidae) by *Leptodactylus* cf. *latrans* (Leptodactylidae). *Herpetol. Notes* 5:163–164.
- Moreira LFB, Smaniotto NP. 2017. *Leptodactylus elenae*. Predation. *Herpetol. Rev.* 48(1):163–164.
- Narvaes P, Rogrigues MT. 2009. Taxonomic revision of *Rhinella granulosa* species group (Amphibia, Anura, Bufonidae), with a description of a new species. *Arq. Zool.* 40(1):1–73. doi: <https://doi.org/10.11606/issn.2176-7793.v40i1p1-73>
- Neto AGS, Barbosa LP, Santos DP, Santos-Silva CR. 2015. Predação de *Hypsiboas faber* (Anura: Hylidae) por *Leptodactylus vastus* em uma lagoa temporária na mata atlântica de Capela, Sergipe. *Nat. On Line.* 13(3):126–127.
- Novaes-e-Fagundes G, Zina J. 2017. *Leptodactylus caatingae*. Predation. *Herpetol. Rev.* 48(3):607.
- Oda FH, Gambale PG, Batista VG. 2016. *Leptodactylus chaquensis* (Anura, Leptodactylidae) as a predator of *Pseudis platensis* (Anura, Hylidae) in southern Brazil. *Lundiana* 12(1):45–48.
- Oda FH, Guerra V, Grou E, Lima LD, Proença HC, Gambale PG, Takemoto RM, Teixeira CP, Campião KM, Ortega JCG. 2019. Native anuran species as prey of invasive American bullfrog *Lithobates catesbeianus* in Brazil: a review with new predation records. *Amphib. Reptile. Conserv.* 13(2):217–226.
- Oliveira RH, Silva MC, Ávila RW. 2014. Predation of *Leptodactylus macrosternum* Miranda-Ribeiro, 1926 (Anura: Leptodactylidae) by *Lygophis dilepis* Cope, 1862 (Squamata: Dipsadidae). *Herpetol. Notes* 7:357–358.
- Pazinato DMM, Trindade ADO, Oliveira SV, Capellarri LH. 2011. Dieta de *Leptodactylus latrans* (Steffen, 1815) na Serra do Sudeste, Rio Grande do Sul, Brasil. *Biotemas* 24(4):147–151. doi: <https://doi.org/10.5007/2175-7925.2011v24n4p147>
- Peres PB, Souza KS, Zocche JJ, Mendonça RA. 2017. *Leptodactylus latrans* (Butter Frog). Diet. *Herpetol. Rev.* 48(1):164.
- Piatti L, Souza FL. 2011. Diet and resource partitioning among anurans in irrigated rice fields in Pantanal, Brazil. *Braz. J. Biol.* 71(3):653–661. doi: <https://doi.org/10.1590/S1519-69842011000400009>
- Pinto RO, Costa-Campos CE. 2015. *Atelopus hoogmoedi* (Harlequin Toad). Predation. *Herpetol. Rev.* 46(4):612.
- Pirani RM, Silva ET, Feio RN. 2010. Tadpole cannibalism in *Leptodactylus cunicularius* Sazima & Bokermann, 1978 (Anura, Leptodactylidae) at a temporary stream in south-eastern Brazil. *Herpetol. Notes* 3:359–360.
- Pombal Jr JP. 2007. Notas sobre predação em uma taxocenose de anfíbios anuros no sudeste do Brasil. *Rev. Bras. Zool.* 24(3):841–843. doi: <https://doi.org/10.1590/S0101-817520070003000034>
- Prado CPA, Uetanabaro M, Lopes FS. 2000. Reproductive strategies of *Leptodactylus chaquensis* and *L. podicipinus* in the Pantanal, Brazil. *J. Herpetol.* 34(1):135–139. doi: <https://doi.org/10.2307/1565249>
- Queiroz MS, Couto RMP, Miguel MC. 2019. *Leptodactylus chaquensis* - the voracious generalist frog as a predator of two different anuran species. *Herpetol. Notes* 12:483–486.
- Regnet RA, Loebmann D. 2017. *Leptodactylus latrans* tadpoles preying the eggs and tadpoles of sympatric anurans. *Herpetol. Bull.* 140:13–15.

- Regueira E, Dávila C, Hermida GN. 2016. Morphological Changes in Skin Glands During Development in *Rhinella arenarum* (Anura: Bufonidae). *Anat. Rec.* 299(1):141–156. doi: <https://doi.org/10.1002/ar.23284>
- Rodrigues AP, Filho JCO. 2004. *Physalaemus nattereri* (NCN). Attempted predation. *Herpetol. Rev.* 35(2):160–161.
- Rojas-Padilla O, Menezes VQ, Rios CHV, Pendu YL, Mira-Mendes CV. 2018. Predation attempt on *Rhinella crucifer* (Wied-Neuwied, 1821) (Anura, Bufonidae) by *Leptodactylus cf. latrans* (Steffen, 1815) (Anura, Leptodactylidae) in southern Bahia, Brazil. *Herpetol. Notes* 11:831–834.
- Rolim DC, Almeida SC, Jim J. 2009. *Scinax fuscomarginatus* (NCN). Predation. *Herpetol. Rev.* 40(1):73.
- Sales RFD, Jorge JS, Kokubum MNC, Freire EMX. 2015. Predation of *Leptodactylus troglodytes* by *Leptodactylus macrosternum* (Anura: Leptodactylidae) in the Brazilian Caatinga. *Herpetol. Notes* 8:421–423.
- Sanabria EA, Quiroga LB, Acosta JC. 2005. Dieta de *Leptodactylus ocellatus* (Linnaeus, 1758) (Anura: Leptodactylidae) en un humedal del oeste de Argentina. *Rev. Peru. Biol.* 12(3):473–477. doi: <https://doi.org/10.15381/rpb.v12i3.2424>
- Santana DO, Rocha SM, Silva IRS, Faria RG. 2012. Predation of *Leptodactylus latrans* (Anura, Leptodactylidae) and *Hypsiobas albomarginatus* (Anura, Hylidae) by *Leptodactylus vastus* (Anura, Leptodactylidae) in north-eastern Brazil. *Herpetol. Notes* 5:449–450.
- Santana MMS, Guedes-Lima PH, Figueiredo VAMB, Reis TS, Cardoso NMC, Costa-Campos CE. 2018. *Leptodactylus pentadactylus* (South American Bull-Frog). Diet. *Herpetol. Rev.* 49(2):305.
- Santos EM, Almeida AV, Vasconcelos SD. 2004. Feeding habits of six anuran (Amphibia: Anura) species in a rainforest fragment in Northeastern Brazil. *Iheringia. Ser. Zool.* 94(4):433–438. doi: <https://doi.org/10.1590/S0073-47212004000400014>
- Santos EM. 2009. Notas sobre predação de anuros em uma poça temporária no nordeste do Brasil. *Bol. Mus. Biol. Mello Leitão.* 25:77–82.
- Santos FP, Sanches PR, Souza JC, Costa-Campos CE. 2018. *Rhinella major* (granulated Toad). Predation. *Herpetol. Ver.* 49(2):309–310.
- Santos CGA, Assis RA, Borges RE, Benvido-Souza M, Santos LRS. 2021. Predation attempt of *Physalaemus marmoratus* (Reinhardt and Lütken, 1862) (Anura: Leptodactylidae) by *Leptodactylus latrans* (Steffen, 1815) (Anura: Leptodactylidae) at Rio Verde, Goiás State, Brazil. *Herpetol. Notes* 14:177–180.
- Schaefer EF, Hamann MI, Kehr AI, González CE, Duré MI. 2006. Trophic, reproductive and parasitological aspects of the ecology of *Leptodactylus chaquensis* (Anura: Leptodactylidae) in Argentina. *Herpetol. J.* 16(4):387–394.
- Scott NJr, Aquino AL. 2005. It's a frog-eat-frog world in the Paraguayan Chaco: Food habits, anatomy, and behavior of the frog-eating anurans. En: Donnelly MA, Crother BI, Guyer C, Wake MH, White ME, editors. *Ecology and evolution in the tropics. A herpetological perspective.* Chicago: University of Chicago Press. p. 243–259.
- Sierra-Rueda AS, Acosta-Ortiz JM. 2020. Depredación de un metamorfo de *Boana* sp. (Anura: Hylidae) por un adulto de *Leptodactylus fuscus* (Anura: Leptodactylidae). *Bol. Asoc. Herpetol. Esp.* 31(1):44–46.
- Silva WR, Giaretta AA, Facure KG. 2005. On the natural history of the South American Pepper Frog, *Leptodactylus labyrinthicus* (Spix, 1824) (Anura: Leptodactylidae). *J. Nat. Hist.* 39(7):555–566. doi: <https://doi.org/10.1080/00222930410001671273>
- Silva ET, Filho OPR. 2009. Predation on juveniles of the invasive American Bullfrog *Lithobates catesbeianus* (Anura, Ranidae) by native frog and snake species in South-eastern Brazil. *Herpetol. Notes* 2:215–218.
- Silva PHE, Santos FP, Costa LFST, Avelar YKS, Santana MMS, Costa-Campos CE. 2018. *Leptodactylus fuscus* (Rufous Frog). Predation. *Herpetol. Ver.* 49(2):303–304.
- Solé M, Dias IR, Rodrigues EAS, Marciano-Jr E, Branco SMJ, Cavalcante KP, Rödder D. 2009. Diet of *Leptodactylus ocellatus* (Anura: Leptodactylidae) from a cacao plantation in southern Bahia, Brazil. *Herpetol. Notes* 2:9–15.
- Sousa JC, Baía RRJ, Costa-Campos CE. 2016. *Rhinella major* (Anura: Bufonidae) and *Leptodactylus macrosternum* (Anura: Leptodactylidae): predation and cannibalism by *Leptodactylus macrosternum*. *Cuad. Herpetol.* 30(1):25–27. doi: <https://doi.org/10.31017/6503>
- Teixeira RL, Vrcibradic D. 2003. Diet of *Leptodactylus ocellatus* (Anura: Leptodactylidae) from coastal lagoons of Southeastern Brazil. *Cuad. Herpetol.* 17(1-2):111–118.
- Teles DA, Rodrigues JK, Teixeira AAM, Ribeiro SC. 2015. *Leptodactylus vastus* (Northeastern Pepper Frog). Diet. *Herpetol. Rev.* 46(1):75.
- Thomas M, Beirne C, Bailey E, Whitworth A. 2017. Attempted predation of the toad *Rhinella marina* (Linnaeus, 1758) (Amphibia: Bufonidae) by *Leptodactylus rhodonotus* (Günther, 1868) (Amphibia: Leptodactylidae) in southeast Peru. *Herpetol. Notes* 10:533–534.
- Toledo LF, Ribeiro RS, Haddad CFB. 2007. Anurans as prey: An exploratory analysis and size relationships between predators and their prey. *J. Zool.* 271(2):170–177. doi: <https://doi.org/10.1111/j.1469-7998.2006.00195.x>
- Vaz-Silva W, Silva HLR, Silva Jr NJ. 2003. *Leptodactylus labyrinthicus* (Labyrinth Frog). Diet. *Herpetol. Rev.* 34(4):359.