



<https://doi.org/10.15446/cr.v5n2.72317>

IMAGENOLOGICAL FINDINGS OF EXTERNAL SNAPPING HIP SYNDROME. CASE REPORT

Keywords: Hip Injuries; Femur; Ultrasonography; Diagnostic Imaging; Snapping Hip.

Palabras clave: Lesiones de la cadera; Fémur; Ultrasonido; Imágenes diagnósticas; Cadera en resorte.

Ingrid Carolina Donoso-Donoso

Hospital Universitario Nacional de Colombia -
Department of Radiology
- Bogotá D.C. - Colombia.

Enrique Calvo-Páramo

Hospital Universitario Nacional de Colombia -
Department of Radiology
- Bogotá D.C. - Colombia.
Universidad Nacional de Colombia
- Bogotá Campus - Faculty of Medicine
- Department of Diagnostic Imaging -
Bogotá D.C. - Colombia.

Roger David Medina-Ramírez

Universidad Nacional de Colombia
- Bogotá Campus - Faculty of Medicine
- Department of Diagnostic Imaging -
Bogotá D.C. - Colombia.

Corresponding author

Roger David Medina-Ramírez.
Department of Diagnostic Imaging,
Faculty of Medicine,
Universidad Nacional de Colombia.
Bogotá D.C. Colombia.
Email: rdmedinar@gmail.com

RESUMEN

Introducción. El síndrome de cadera en resorte externa es una entidad en la cual hay una sensación de dolor acompañada de un sonido palpable durante el movimiento de la cadera. Esta es una condición ampliamente conocida por los ortopedistas, pero aún es necesario que los hallazgos imagenológicos sean reconocidos por todos los radiólogos con el fin de brindar mayor información que permita un adecuado manejo multidisciplinario. La Z-plastia de la banda iliotibial es la técnica de tratamiento más reconocida y con mejores resultados.

Presentación del caso. Paciente femenino con síndrome de cadera en resorte externo bilateral sintomático en el lado derecho, quien fue manejada de forma conservadora sin adecuada respuesta y requirió manejo quirúrgico por vía artroscópica. Las técnicas utilizadas en la paciente no fueron exitosas y se presentó recurrencia de los síntomas.

Discusión. El diagnóstico del síndrome de cadera en resorte es principalmente clínico. Sin embargo, el aporte de las imágenes diagnósticas es importante para caracterizar las estructuras involucradas en este proceso nosológico, para realizar el planeamiento terapéutico y para hacer el seguimiento.

Conclusión. Conocer los hallazgos imagenológicos en ultrasonido y resonancia magnética del síndrome en cadera en resorte externa permite a los radiólogos identificarlo y hacer aportes al manejo de esta patología en forma oportuna.

ABSTRACT

Introduction: External snapping hip syndrome is characterized by a painful sensation accompanied by an audible snapping noise in the hip when moving. Even though orthopedists are widely aware of this condition, imaging findings still need to be recognized by all radiologists in order to provide more information that allows for the best multidisciplinary treatment. Z-plasty of the iliotibial band is the most used treatment with the best results.

Case presentation: Female patient with bilateral external hip snapping syndrome on the right side, who was treated initially in a conservative manner without adequate response; hence, she required surgical management with arthroscopy. All treatment options used for this patient were not successful, and symptoms recurred.

Discussion: The diagnosis of snapping hip syndrome is mainly clinical. However, the contribution of diagnostic imaging is important to characterize the structures involved in this nosological process, in order to develop the therapeutic planning and do the follow-up.

Conclusion: Knowledge on ultrasound and magnetic resonance findings related to this pathology allows radiologists to identify this syndrome and contribute to a timely treatment.

INTRODUCTION

The iliotibial band (ITB) is a flat morphological structure that is made up of connective tissue and is part of the tensor fasciae latae muscle. It extends the fibers of the anterior tensor fasciae latae, and involves the gluteal aponeurosis and the posterior fibers of the gluteus maximus

muscle (1) (Figure 1). It originates at the iliac crest, runs parallel to the diaphysis of the femur, inserts at the Gerdy's tubercle—which is located in the anterior and lateral corner of the proximal tibia—and joins the lateral femoral condyle and the tibia. Its functions are to extend, abduct, and laterally rotate the hip, and it is also involved in maintaining posture and lateral knee stabilization.




Figure 1. Magnetic resonance and ultrasound (cross-sectional plane) of the right hip of an asymptomatic patient, where the relationship between the iliotibial band, the tendon of the gluteus maximus and the greater trochanter is evident.

Source: Document obtained during the study.

Snapping hip syndrome is a complex condition in which there is an audible noise in the hip when moving (flexion or extension), usually accompanied by pain. (2) This condition was first described by Nunziata & Blumenfeld (3) in 1951; however, in 1995, Allen introduced the term “coxa saltans” to differentiate the causes. (2) At present, three types are described: internal, external (included by some authors in extra-articular) (1,2,4) and intra-articular. The first is caused by the IBT sliding over the femoral head or acetabular ridge, the second by the IBT sliding over the greater trochanter

(2), and the third by different causes, mainly degenerative, such as labrum tears, damage to the ligamentum teres or loose bodies in the hip. (1,2,4).

The external subtype is the most common type of “coxa saltans” and refers to the snapping sensation caused by a sudden motion of the IBT, whose posterior portion and, to a lesser extent, the distal and anterior portion of the gluteus maximus thicken during hip movements, most frequently during flexion, external rotation and abduction. (1,2,4) During hip extension, the IBT slides over the greater trochanter and returns

to the anterior portion during flexion. (2) Pain may or may not be observed during movement, and if it occurs, it coincides with the noise.

External snapping hip syndrome is associated with repetitive physical activities or overuse of anatomical structures related to IBT movements. Some intrinsic risk factors have been identified, such as decreased cervico-diaphyseal angle (coxa vara), the narrowing of the bi-iliac distance, increased distance between major trochanters, prominence of the major trochanter, and fibrosis of the iliotibial band secondary to multiple intramuscular injections. (5-7)

The incidence and prevalence of this syndrome in the general population is not clear (2), although it is usually found in athletes and dancers, with a reported frequency of up to 62% in young women (8), and also in an asymptomatic manner in between 5% to 10% of the population. (9) Affected people visit their doctor because they have difficulty performing common activities such as running, climbing stairs, lifting heavy objects, and even, in some cases, walking. The physical examination is characterized by the reproduction of the noise or palpation of the dislocation during triggering maneuvers: hip flexion and extension.

CASE PRESENTATION

A 29-year-old female patient from Bogotá D.C., with a master's degree in education, white, who works as a psychologist, attended consultation due to symptoms with a year of evolution that began with a sensation of "snap" or prominence in the right greater trochanter, associated with pain, which occurred during walking and when flexing the hip to sit.

Medical history included ligamentous laxity (under study to confirm Ehlers-Danlos syndrome), flexible flatfoot treated with insoles with internal wedge in the hindfoot, length asymmetry in the

lower limbs, severe lumbar hyperlordosis, scoliosis from infancy in management with orthosis and Raynaud's phenomenon under study with final diagnosis of vasculitis.

The physical examination showed height of 1.52m, weight of 45k and hypotonic posture caused by lumbar hyperlordosis and thoracic hyperkifosis. The evaluation in standing position revealed tendency to genu valgum with flat foot that compensates with the retraction of the tensor fasciae latae muscle; asymmetry of 4mm in the comparative length of the lower limbs, being greater on the right side; myofascial snapping of the gluteus medius with external extension and flexion hip rotation, without severe hamstring or tensor fasciae latae retraction due to its superimposed hyperlaxity; and significant weakness of the buttocks, predominantly on the right side. Hip impingement and pain were observed when internally rotating the hip bilaterally in a flexed position, especially on the right side, with localized pain in the greater trochanter.

The patient underwent multiple analgesic (until being prescribed with hydrocodone + acetaminophen, which she takes irregularly) and anti-inflammatory treatments, and received multiple physiotherapy sessions without improvement; in fact, her symptoms increased progressively. After assessment by Orthopedics, she was diagnosed with bilateral snapping hip syndrome, with retraction of the tensor fasciae latae. An ultrasound of the hip was performed on both sides, finding thickening of the right tensor fasciae latae (4mm), alteration of the fibrillary pattern and tendon protrusion over the greater trochanter during dynamic assessment, accompanied by an audible and palpable "snap" (Figure 2) (Video 1). Magnetic resonance imaging (MRI) of the hip showed bilateral edema of the ITB and slight thickening of the ITB on the right side (2.3mm) (Figure 3).




Figure 2. Dynamic ultrasound of the iliotibial band over the greater trochanter. A) neutral position of the right hip; B) hip internal rotation; C) thickening of the iliotibial band measured in the transverse plane associated with heterogeneous echogenicity of the tendon.

Source: Document obtained during the study.



Video 1. Dynamic ultrasound of the iliotibial band over the greater trochanter before the patient underwent the surgical procedure.

Source: Document obtained during the study.

Note: Dear reader in order to watch the video you need to download the PDF file and open it with Acrobat Reader (having previously installed the Adobe Flash Player add-on) and click on the video once.




Figure 3. Magnetic resonance imaging. A) Enhanced axial T1-image showing slight thickening and undulation of the right iliotibial band (black arrow); B) axial plane with Short tau inversion recovery sequences, where an increase in signal intensity of the iliotibial band is identified on both sides (white arrows).

Source: Document obtained during the study.

Two years after the onset of the symptoms, the patient required an arthroscopy, so a T-shaped incision of the ITB was made in the right hip using radiofrequency. During a second surgery, an X-shaped incision was made in the fasciae latae muscle on the left side, also by radiofrequency, with

subsequent formation of a rhombus and fixation. The patient presented recurrence of predominant symptoms in the right hip. A control ultrasound was performed again, showing greater tendon thickening and peritendinous fluid, probably of post-surgical origin (Figure 4) (Video 2).




Figure 4. Ultrasound of the patient taken during post-surgical control 1 month after iliotibial band release. Persistence of symptoms. A) Increase in the thickness of the right iliotibial band (7.5 mm) with respect to the initial ultrasound. B) Dislocation of the iliotibial band during hip external rotation.

Source: Document obtained during the study.



Video 2. Ultrasound of the patient taken during post-surgical control 1 month after iliotibial band release.

Source: Document obtained during the study.

Note: Dear reader in order to watch the video you need to download the PDF file and open it with Acrobat Reader (having previously installed the Adobe Flash Player add-on) and click on the video once.

One year after the first surgery, the woman underwent a new surgical procedure on the right hip, consisting of ITB opening, partial synovectomy and tenotomy of the hip at two levels, and proximal and distal release. Severe tendon fibrosis secondary to previous surgical procedures was reported as an additional finding. At the time of completion of this case report, the patient had shown partial improvement of the symptoms and had not undergone any new imaging controls.

DISCUSSION

The main structure involved in external snapping hip syndrome is ITB; however, to a lesser extent, there may also be involvement of the tendon of the gluteus maximus. (10) Although the diagnosis is mainly clinical, diagnostic images allow ruling out other causes of hip pain such as bone or soft tissue alterations, bursa, labrum tear, among others. The most widely used and useful modality is dynamic ultrasound. Simple x-rays are usually normal, but allow ruling out intra-articular causes of this syndrome. There is still no consensus to determine diagnostic parameters. (2,11)

Dynamic ultrasound shows thickening of the iliotibial band and heterogeneous echogenicity of the fibers, as well as secondary findings such as distension of the greater trochanteric bursa. This dynamic evaluation is useful since it verifies movement of the ITB or the tendon of the gluteus maximus on the greater trochanter, and allows perceiving the dislocation or listening to the “snap”, thus leading to the definitive diagnosis of this syndrome. (2,4)

The appropriate tool to perform the dynamic ultrasound evaluation is a high frequency linear transducer (12 MHz or higher), with the patient in supine or standing position. Based on the radiologist’s preference, the transducer is placed transversely to the greater trochanter,

and the exploration is performed with active and passive flexion and external rotation maneuvers of the hip. The abrupt displacement of the ITB or the gluteus maximus muscle is related to the painful “snap” perceived by the patient. (4,7)

According to the literature, the MRI findings include thickening of the iliotibial band (>2mm) and of the anterior edge of the tendon of the gluteus maximus (>2mm), with undulation of the contours of both structures (7); the signal is hypointense in T1 and hyperintense in T2. Inflammatory changes of the greater trochanter bursa, visualized as a hyperintense T2 collection, and peritendinous enhancement following administration of intravenous contrast medium can also be seen; however, these findings are nonspecific and should always be analyzed bearing in mind the patient’s clinical features. (1) Retrospectively, ipsilateral gluteus maximus atrophy has been identified as an additional finding of this syndrome. It is important to enlarge the field of vision in order to fully visualize the region of interest: this includes the entire pelvis and its soft tissues.

Computerized tomography permit a better evaluation of bone structures, while the multiplanar reconstructions that can be carried out allow better characterizing the findings that predispose to the snapping hip syndrome, and ruling out intra-articular causes. This study should not be the first choice as the resolution of the soft tissues is not good. (8)

Taking into account the findings of the present case, a dynamic ultrasound was the diagnostic modality that best allowed evaluating the patient and corroborating what was identified during the physical examination. Besides identifying the key finding, which was the subluxation of the ITB over the greater trochanter accompanied by pain, changes in the soft tissues (in the second ultrasound) were found that could explain the persistence of symptoms.

Treatment in these patients includes rest, physical means (local heat), stretching exercises, release of loads during daily activities, and analgesics and non-steroidal anti-inflammatory drugs. Conducting anesthetic or corticosteroid infiltration of the trochanteric bursa or iliotibial band has been proposed if there is no response. (8)

When there is no response to conservative treatment, surgery should be done in order to eliminate tendon protrusion. Possible surgical procedures include resection of the major trochanteric bursa, trochanter osteotomy and, in refractory cases, arthroscopic release of the ITB or lengthening of the ITB by Z-plasty. (1,2)

The failure of surgery in the reported case is still cause for study. Specialists believe that the recurrence of symptoms may be associated to a marked fibrotic process following the two surgical procedures; however, that baseline hyperlaxity may be associated with recurrence of symptoms is considered a possibility.

CONCLUSIONS

This case report describes the ultrasound and MRI imaging findings of external snapping hip syndrome, which allow radiologists to recognize and contribute to the management of this pathology in a timely manner.

Dynamic ultrasound is the most useful technique for the diagnosis of snapping hip syndrome, as it allows confirming the main finding, which is the protrusion of the ITB or the tendon of the gluteus maximus over the greater trochanter. Other techniques, such as x-ray and MRI, provide secondary information on possible differential diagnoses of painful hip or help rule out the intra-articular variant of this syndrome.

CONFLICT OF INTEREST

None stated by the authors.

FUNDING

None stated by the authors.

ACKNOWLEDGEMENTS

None stated by the authors.

REFERENCES

1. **Fiato R, Passanante GJ, Skalski MR, Patel DB, White EA, Matcuk GR Jr.** The iliotibial tract: imaging, anatomy, injuries and other pathology. *Skeletal Radiol.* 2017;46(5):605-22. <http://doi.org/c7f9>.
2. **Potavilo G, Bugiantella W.** Snapping hip syndrome: systematic review of surgical treatment. *Hip Int.* 2017;27(2):111-21. <http://doi.org/f933zf>.
3. **Nunziata A, Blumenfeld I.** Cadera a resorte: a propósito de una variedad. *Prensa Med Argent.* 1951;38832):1997-2001.
4. **Choi YS, Lee SM, Song BY, Paik SH, Yoon YK.** Dynamic Sonography of External Snapping Hip Syndrome. *J Ultrasound Med.* 2002;21(7):753-8. <http://doi.org/c7gb>.
5. **Krishnamurthy G, Connolly BL, Narayanan U, Babyn PS.** Imaging findings in external snapping hip syndrome. *Pediatr Radiol.* 2007;37(12):1272-4. <http://doi.org/dxv9zj>.
6. **Johnston CA, Wiley JP, Lindsay DM, Wiseman DA.** Iliopsoas Bursitis and Tendinitis. *A review.* *Sports Med.* 1998;25(4):271-83. <http://doi.org/dcwkn5>.
7. **Choi JE, Sung MS, Lee KH, Lee BY, Park JM, Kim JY, et al.** External snapping hip syndrome: emphasis on the MR imaging. *J Korean Soc Radiol.* 2010;62:185-90. <http://doi.org/c7gc>.
8. **Lee KS, Rosas HG, Phancao JP.** Snapping Hip: imaging and treatment. *Semin Musculoskelet Radiol.* 2013;17(3):286-94. <http://doi.org/f43vws>.
9. **Lewis C.** Extra-articular Snapping Hip: A literature Review. *Sports Health.* 2010;2(3) 186-90. <http://doi.org/b43jtb>.

- 10. Chang CY, Kreher J, Torriani M.** Dynamic sonography of snapping hip due to gluteus maximus subluxation over greater trochanter. *Skeletal radiol.* 2016;45(3):409-12. <http://doi.org/f8bj7z>.
- 11. Teitz CC, Garrett WE Jr, Miniaci A, Lee MH, Mann RA.** Tendon problems in athletic individuals. *Instr Course Lect.* 1997;46:569-82.