

Performance of broilers submitted to different intensities and duration of thermal stress

Lucas Henrique Pedrozo Abreu ^a, Tadayuki Yanagi Junior ^a, Marcelo Bahuti ^a, Renato Ribeiro de Lima ^a, Dian Lourençoni ^b & Édison José Fassani ^a

^a Universidad Federal de Lavras, Lavras, Brasil. lucas.abreu@ufla.br, yanagi@ufla.br, marcelo_bahuti@hotmail.com, rrlima@ufla.br, fassani@ufla.br

^b Universidad Federal del Vale do São Francisco, Juazeiro, Brasil. dian.lourenconi@univasf.edu.br

Received: May 2nd, 2019. Received in revised form: August 30th, 2019. Accepted: October 2nd, 2019.

Abstract

The performance of broilers submitted to thermal discomfort in the first weeks of life may be impaired, since the thermoregulatory system of the bird is not fully developed. The objective of this study was to analyze the effects of different thermal challenges on the performance of broilers. Two hundred and forty broilers were submitted to different desired dry-bulb temperatures (t_{db} : 24, 27, 30 and 33°C) and stress duration (SD: 1, 2, 3 and 4 days) in the second week of life. Feed intake (FI), weight gain (WG) and feed conversion (FC) were evaluated at 21 days of age. Although differences were verified for FI ($p < 0.05$), none were observed for WG ($p > 0.05$). Reductions of FC for the 24, 27 and 33°C t_{db} were 14.8, 8.6 and 10.1%, respectively. The smallest FC was observed at 30°C, indicating comfort condition.

Keywords: animal environment; poultry farming; productive responses; thermal comfort.

Desempeño de pollos de engorde sometidos a diferentes intensidades y duraciones de estrés

Resumen

El desempeño de pollos de engorde sometidos a estrés térmico en las primeras semanas de vida puede ser perjudicado, una vez que el sistema termorregulador del ave no está completamente desarrollado. El objetivo de este trabajo fue analizar los efectos de diferentes desafíos térmicos sobre el desempeño de pollos de engorde. Doscientos cuarenta pollos de engorde fueron sometidos a diferentes temperaturas de bulbo seco del aire deseada (t_{bs} : 24, 27, 30 y 33°C) y a duraciones de estrés (DE: 1, 2, 3 y 4 días) en la segunda semana de vida. El consumo de ración (CR), ganancias de peso (GP) y conversión alimenticia (CA) fueron evaluadas a los 21 días de edad de las aves. A pesar de que las diferencias fueron verificadas para CR ($p < 0,05$), ninguna diferencia fue observada para GP ($p > 0,05$). Reducciones de CA para las t_{bs} de 24, 27 y 33°C fueron de 14,8, 8,6, y 10,1%, respectivamente. El CA más pequeño se observó a 30°C, lo cual se caracterizó como condición de confort.

Palabras clave: ambiente animal; avicultura; respuestas productivas; confort térmico.

1. Introduction

To obtain maximum productivity, broilers need to be reared in an appropriate thermal environment, with adequate control of air dry-bulb temperature (t_{db}), relative humidity (RH), air speed and radiation [1,2]. Thus, it is possible to maintain the productivity as well as attend to new policies and exigences of the market related to animal well-being.

Therefore, aiming to provide ideal conditions throughout production phases, the thermal control in the breeding environment is necessary [3,4], mainly in the first days of life [5], once the bird's thermoregulation system is not completely developed and there is some limitation to the bird produces heat.

When subjected to unfavorable thermal conditions, the birds are subjected to thermal stress [6], thus modifying their

How to cite: Abreu, L.H.P., Yanagi Junior, T., Bahuti, M., Lima, R.R., Lourençoni, D. and Fassani, E.J., Performance of broilers submitted to different intensities and duration of thermal stress. DYNA, 86(211), pp. 131-137, October - December, 2019.

feeding behavior, decreasing or increasing feed intake (FI) [7]. This modification in eating behavior helps in physiological changes to adapt to stress [8]. However, the energy expenditure that should be used to provide adequate weight gain (WG) will be allocated to the process of body thermoregulation [9,10], harming productive performance [11].

The evaluation of the broilers productive responses according to the thermal conditions of the facilities is performed through the analysis of zootechnical parameters, such as FI, WG and feed conversion (FC) [12].

Several studies have shown that the productive responses of broilers are closely related to the thermal conditions of the facilities [13-15]. However, most researches only vary the intensity of the thermal stress, not varying the its duration [15-18].

Given the importance of the thermal environment and its influence on broilers productive responses in the initial stages, the objective of this study was to evaluate the performance of broilers submitted to different intensities and duration of thermal stress in the second week of life.


2. Material and methods

The research was carried out in air-conditioned wind tunnels, and the procedures performed during the experiment were approved by the Ethics Committee for Animal Use (Comissão de Ética para Uso de Animais - CEUA) of the Universidade Federal de Lavras - (UFLA), Minas Gerais, protocol No. 008/12.

Four air-conditioned wind tunnels (0.8 x 5.0 m) were used, which were built in steel sheets and PVC pipes (Fig. 1a), each tunnel being equipped with two electric heaters and two humidifiers, in two stages of operation with capacity of 600 W and 300 mL_{water} h⁻¹ per stage, respectively.



a)



b)

Figure 1. a) Overview of air-conditioned wind tunnels. b) Top view of the cages placed inside the tunnels. Source: The Authors.

The ventilation inside each tunnel was performed by means of an exhaust fan, 40 cm in diameter, and the speed adjusted by potentiometer. Inside the tunnels, cages of 0.24 m² (0.40 x 0.60 m) were allocated, in which they were divided into three equal parts and equipped with independent feeders and drinkers in each division, thus characterizing three replicates (Fig. 1b). To condition the temperature inside the room, two air conditioning systems with 5,272.2 W (18,000 BTUs) of power were used.

The control of the thermal environment within the wind tunnels was performed through a datalogger (CR1000, Campbell Scientific), a relay controller (SDM-CD16AC, Campbell Scientific), a channel multiplexer (AM16/32B, Campbell Scientific), and t_{db} and RH sensors (HMP45c, Vaisala, accuracy of ± 0.3°C for t_{db} and ± 2 % for RH).

During the experimental period, 240 chicks of the Cobb 500® males and females were used. The chicks were from the same hatchery and vaccinated against Marek, Gumboro and Bouda aviary diseases. The birds arrived at the experiment shortly after birth and remained until they were twenty-two days old. During this period, water and feed *ad libitum* were supplied to the birds to meet their nutritional requirements according to [19]. The feed used was the same for all the chicks throughout the experimental period, with no variation of their formulation. The broiler chicks received a corn and soybean meal mash diet, formulated according to the nutritional recommendations described by [19], for chickens from 1 to 21 days old, containing 3000 kcal of metabolizable energy per kg of diet, 21.4% crude protein, 0.90% Ca, 0.43% P disponible, 0.21% Na, 1.20% lysine digestible, 0.89% methionine + cystine digestible, 0.80% threonine digestible and 0.21% tryptophan digestible.

The RH was maintained at 60 ± 1% and the air velocity at 0.2 ± 0.1 m s⁻¹, characterized as comfort for the broilers [20]. The luminosity was adjusted via a luxmeter (LDR-380, accuracy of ± 3%), with intensities of 25 and 10 lux for the first and second week of life, respectively [21].

Table 1.

Desired air dry-bulb temperature (t_{db}) and thermal stress duration (in bold) applied to broilers from 1 to 21 days of age during the second week of life.

Days of life		Air dry-bulb temperature (°C)					Week 3 15-21
		Week 1	Week 2			Week 3	
		1-7	8	9	10		
Stage	Tunnel						
1	A		33	30	30	30	
	B	33	33	33	30	30	27
	C		33	33	33	30	
	D		33	33	33	33	
A	27		30	30	30		
2	B	33	27	27	30	30	27
	C		27	27	27	30	
	D		27	27	27	27	
	A		24	30	30	30	
3	B	33	24	24	30	30	27
	C		24	24	24	30	
	D		24	24	24	24	
	A, B, C and D		33			30	

Source: The Authors.

The experiment was carried out in four stages of 21 days each (Table 1), using sixty birds in each stage.

In each division of the cage were allocated five birds, characterizing fifteen birds per air-conditioned wind tunnel in the first week of life. In order to maintain optimal comfort density, in the second and third week of life, four and three birds, respectively, were maintained in each replicate, according to recommendations of the lineage guide [21]. The cleaning of the breeding environment was carried out daily to avoid the formation of gases and to provide an adequate environment for the development of the broilers. The time that handling opening of each air-conditioned wind tunnel was maintained opened was very short to avoid changes in the thermal environment desired.

The birds were submitted to thermal challenge only in the second week of life, from the 8th day of life. During the first and third weeks of life, the t_{db} were maintained in the thermoneutral zone, i.e. 33°C and 27°C, respectively [22]. In the second week, the differences between the treatments were established between the intensities and duration of the thermal challenge, and the stress intensities were 24, 27 and 33°C for each stage, in addition to t_{db} of 30°C, considered as of comfort for the second week of life [22]. These desired t_{db} values were maintained throughout each treatment.

In Table 1, it can be observed that the thermal stress was applied in four duration (1, 2, 3 and 4 days) in the first four days that compose the second week (8th, 9th, 10th and 11th days of life). Soon after this period the t_{db} returned to the recommended value as characteristic of the thermal comfort zone (30°C) [5,23].

The experimental stage submitted to t_{db} of 30°C is considered as control (comfort), and the others were maintained in order to provide thermal stresses both by low (24 and 27°C) and high (33°C) temperatures [22,24].

The values of FI, WG and FC accumulated in 21 days were analyzed, adopting the completely randomized design (DIC) in split plots in time scheme. The plots were set according to the 4x4 factorial arrangement, that is, four t_{db} in the second week of the birds' life (33°C, 30°C, 27°C and 24°C) and four stress duration (1, 2, 3 and 4 days), according to the model specified in eq. (1). The variables were analyzed by the statistical program SISVAR 5.3 [25] submitted to analysis of variance, and the means were compared using the Scott-Knott test and regression at the 5% probability level.

$$Y_{ijkl} = \mu + \alpha_i + \beta_j + \alpha\beta_{ij} + \epsilon_{ijk} + \tau_l + \alpha\tau_{il} + \beta\tau_{jl} + \alpha\beta\tau_{ijl} + e_{ijkl} \quad (1)$$

in which, Y_{ijkl} is the observed value of the replicate k of the i-th temperature by j-th duration at time l; μ is a constant; α_i is the effect of i-th temperature; β_j is the effect of the j-th duration of the thermal stress; $\alpha\beta_{ij}$ is the interaction between i-th temperature and j-th duration; ϵ_{ijk} is the whole-plot error; τ_l is the effect of the time; $\alpha\tau_{il}$ is the interaction between i-th temperature and l-th time; $\beta\tau_{jl}$ is the interaction between j-th duration and l-th time; $\alpha\beta\tau_{ijl}$ is the interaction between i-th temperature, j-th duration and l-th time, and e_{ijkl} is the subplot error.

3. Results and discussion

The thermal environments inside of the air-conditioned wind tunnels were monitored during the experimental period to verify the t_{db} and RH values to which the birds were subjected. The averages and standard deviations (in parentheses) of the observed values are listed in Table 2.

In the first week of life the t_{db} and RH desired were 33°C and 60%, respectively, and the observed value (Table 2) was very close at all stages, with the highest standard deviations for t_{db} and RH were $\pm 0.3^\circ\text{C}$ and $\pm 1.4\%$, respectively. During the second week of life, period which the birds were submitted to different intensities of t_{db} , the highest standard deviations found were $\pm 0.3^\circ\text{C}$ for t_{db} and $\pm 1.7\%$ for RH. To the t_{db} of third week of life, in which the birds were submitted to t_{db} of 27°C, considered of comfort [22], it is observed that the averages of $t_{db,obs}$ are close to desired. The small variations demonstrate the efficiency of the air-conditioned wind tunnels for the maintenance of the desired thermal variables, corroborating with [26].

The interaction between t_{db} and thermal stress duration was significant ($p < 0.05$, F test) for the FI variable (Table 3). When analyzing the t_{db} within each stress duration (Table 3), it was generally observed that the highest FI averages occurred at t_{db} of 24°C ($p < 0.05$, Scott-Knott test), except for the duration of 4 days that no significant difference was verified ($p > 0.05$, Scott-Knott test). The increased in FI in low temperature stress conditions is due to the bird's need to obtain more energy from its diet [27].

On the other hand, the lowest FI for the different stress duration studied was verified for t_{db} of 30°C ($p < 0.05$, Scott-Knott test), except for the stress duration of 4 days, in which there was no statistical difference between the various t_{db} evaluated ($p < 0.05$, Scott-Knott test).

Table 2. Air dry-bulb temperature values desired ($t_{db,desired}$, °C), observed means ($t_{db,obs}$ (°C) and RH_{obs} (%)) and standard deviations (in brackets) inside the air-conditioned wind tunnels.

Stage	Thermal Variables	Week 1				Week 2			Week 3
		1 day				2 days	3 days	4 days	
1	$t_{db,desired}$	33.0				24.0			27.0
	$t_{db,obs}$	33.1(0.3)				24.3 (0.3)	24.3 (0.3)	24.3 (0.3)	27.2(0.3)
	RH _{obs}	60.3(0.6)				60.4 (0.7)	60.4 (0.8)	59.9 (1.0)	60.8 (0.6)
2	$t_{db,desired}$	33.0				27.0			27.0
	$t_{db,obs}$	33.1(0.3)				27.2 (0.3)	27.2 (0.2)	27.1 (0.2)	27.2 (0.3)
	RH _{obs}	60.3(0.9)				60.0 (1.6)	60.3 (1.0)	60.0 (0.9)	60.4 (0.9)
3	$t_{db,desired}$	33.0				33.0			27.0
	$t_{db,obs}$	33.1(0.3)				33.0 (0.3)	33.0 (0.3)	33.0 (0.3)	27.2(0.3)
	RH _{obs}	60.2(1.4)				60.0 (1.7)	60.1 (1.3)	59.9 (1.4)	60.5 (1.3)
4	$t_{db,desired}$	33.0				30.0			27.0
	$t_{db,obs}$	33.2 (0.3)				30.2(0.2)			27.2 (0.2)
	RH _{obs}	60.1 (1.1)				60.3(0.7)			60.3 (0.8)

$t_{db,desired}$ = Desired air dry-bulb temperature; $t_{db,obs}$ = Observed air dry-bulb temperature; RH_{obs} = Observed air relative humidity.
Source: The Authors.

Table 3.

Average values and standard deviation (in brackets) of the mean accumulated feed intake (FI, g) of broilers at 21 days of age, submitted to different intensities and thermal stress duration.

Air dry-bulb temperature (°C)	Thermal stress duration (days)			
	1	2	3	4
24	941.33 (80.89) c A	901.93 (47.78) b A	902.00 (75.49) b A	867.86 (75.98) a A
27	824.43 (31.30) b A	811.36 (59.47) a A	888.53 (68.68) b A	830.03 (36.55) a A
30	717.86 (37.45) a A	758.96 (16.04) a A	776.26 (28.09) a A	797.63 (50.27) a A
33	911.86 (28.16) c A	826.83 (46.76) a B	783.70 (36.34) a B	808.00 (26.17) a B
CV (%)	6.05			

Averages with different letters (uppercase in the row and lowercase in the column) indicate statistical differences ($p < 0.05$) by the Scott-Knott test.

CV = Coefficient of variation.

Source: The Authors.

For t_{db} of 33°C, as the duration of thermal stress increases, it was verified the reduction in the effect of t_{db} in FI at 21 days of the birds' age, indicating the occurrence of adaptation of the broilers to the stress environment. It is also verified that t_{db} of 33°C applied to thermal stress duration of 2, 3 and 4 days, does not yet represent a condition of significant discomfort ($p > 0.05$, Scott-Knott test) for broilers when compared to the condition recommended in the literature as of comfort ($t_{db} = 30^\circ\text{C}$) [5,23].

However, considering the stress duration of 1 day, there was statistical difference between the t_{db} of 33°C and 30°C. In conditions of high t_{db} , the reduction of FI is expected to decrease the metabolic increment [28], thus facilitating the homeothermic control [29]. In this study, it was verified that FI on the first day of stress was also higher at 33°C, this probably occurred due to this t_{db} not fully characterize discomfort by high temperature. In Fig. 2, the exploratory analysis is illustrated in bar format, the days of thermal stress (1, 2, 3 and 4 days) for the FI associated with each t_{db} .

The interaction of t_{db} and the thermal stress duration was not significant ($p > 0.05$, F test) for WG. Analyzing the separate effect of the different t_{db} and thermal stress duration (Table 4), there was no significant difference ($p > 0.05$, F test). Thus, it is verified that even with the variation of FI, WG was the same in all treatments tested, thus characterizing as compensatory gain reported in several studies [30-32]. This occurs when broilers are housed in a breeding system in which t_{db} is outside the comfort zone, since the birds need to redirect the energy destined for production to the thermoregulatory process [9,10]. Thus, the birds possibly used the energy supplied by the feed to maintain body temperature [33].

According to the analyzes presented, the highest FI values are associated with the thermal stress of (24 and 33°C), the average being 926.60g and for thermal comfort the average was 717.86g, demonstrating that the birds consumed an average of 208.73g more to obtain the same WG at 21 days. As the feed represents about 60% of the total costs for broiler breeding [34], any extra consumption represents financial

risks for the segment, increasing the production cost, thus reducing the profit margin. In modern poultry farming, in order to keep the activity even more attractive and competitive on the world scenario, lower production costs should be obtained [35], thus, it is important to control the thermal variables inside the facilities.

The interaction ($p > 0.05$, F test) between the factors assessed (t_{db} and thermal stress duration) for the FC (Table 5) was not observed. The best (1.28) and worst (1.47) FC values were observed for t_{db} of 30°C and 24°C, respectively, which are statically different ($p < 0.05$, Scott-Knott test). On the other hand, the t_{db} of 27°C and 33°C presented intermediate FC values, and similar ($p > 0.05$, Scott-Knott test).

The lowest FC value associated with t_{db} of 30°C is in agreement with literature, given that this temperature is considered as thermal comfort for broilers in the second week of life [5,23].

On the other hand, the poor performance of the broilers is associated with the stress of low temperatures (24°C), because when subjected to cold stress, the growing birds modify their behavior, in which affects their physiological and metabolic functions, decreasing their productivity [10]. For t_{db} of 33°C, the FC rise in relation to the thermal comfort condition (30°C) was verified in the first day of stress duration. It was due to the use of energy intake from the feed for helping the biophysical and chemical adjustments of broilers to dissipate body heat. There was no significant difference ($p > 0.05$, F test) between FC values for the various durations of thermal stress. In Fig. 3, a bar chart showing the FC as a function of t_{db} , comparing them with the performance of broilers, raised in comfort conditions, in the second week of life, according to the breed's Guide [36], with the reference value of 1.32 g g^{-1} .

When submitted to t_{db} 24, 27 and 33°C, the birds' performance was worse in 14.8, 8.6 and 10.1%, respectively, when compared with the comfort t_{db} (30°C). [29] observed a reduction of this variable in 9.4 and 5.7% when subjected to stress by low (24°C) and high temperatures (33°C), respectively, in the second week of life. Therefore, the results found in this study, as that obtained by [29], indicating that broilers respond in different ways when subject to low or high temperatures. Theses results corroborate with [37] that reported the deleterious effect of thermal stress in promote a fall in the final performance of the lot, damaging the productivity and profitability of the activity.




Figure 2. Analysis of feed intake (FI, g) as a function of air dry-bulb temperatures (t_{db}) and thermal stress duration evaluated.

Source: The Authors.

Table 4.

Mean weight gain (WG) and standard deviation (in brackets) accumulated in the broilers' third week of life submitted to different intensities and duration of thermal stress.

Air dry-bulb temperature (°C)	Weight gain (g)	Thermal stress duration (days)	Weight gain (g)
24	613.77 (19.74) a	1	605.33 (28.80) a
27	599.22 (21.08) a	2	598.88 (11.41) a
30	592.68 (21.29) a	3	612.73 (41.25) a
33	590.13 (44.33) a	4	578.87 (13.92) a
CV (%)	5.96		

Means followed by different letters differ from one another by the Scott Knott test (p <0.05).

CV = Coefficient of variation.

Source: The Authors.

Table 5.

Mean and standard deviation (in brackets) of the accumulated feed conversion (FC) in the broilers' third week submitted to different intensities and duration of thermal stress.

Air dry-bulb temperature (°C)	Feed conversion (g g ⁻¹)	Length of stress (days)	Feed conversion (g g ⁻¹)
24	1.47 (0.01) c	1	1.38 (0.08) a
27	1.39 (0.03) b	2	1.37 (0.10) a
30	1.28 (0.06) a	3	1.38 (0.08) a
33	1.41 (0.02) b	4	1.42 (0.04) a
CV (%)	5.16		

Means followed by different letters differ from one another by the Scott Knott test (p <0.05).

CV = Coefficient of variation.

Source: The Authors.




Figure 3. Feed conversion (FC, g g⁻¹) of broilers subjected to different air dry-bulb temperatures (t_{db}), compared to the performance of broilers kept in thermal comfort according to the lineage manual.

Source: The Authors.

4. Conclusions

The performance of the broilers was affected as a function of the thermal stress intensities, and the best performance based on FC variable was observed for t_{db} of 30°C, confirmed as thermal comfort for the birds' second week of life. The reduction of feed conversion to the t_{db} of 24, 27 and 33°C were 14.8, 8.6 and 10.1%, respectively.

Acknowledgments

The authors would like to thanks CAPES - Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brazil (Financing Code 001), the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq) (Process 307746/2014-3) and the Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) for the financial support to this research.

References

- [1] Garcia, R.G., Almeida Paz, I.C.L., Caldara, F.R., Nääs, I.A, Pereira, D.F. and Ferreira, V.M.O.S., Selecting the most adequate bedding material for broiler production in Brazil. *Brazilian Journal of Poultry Sciences*, 14(2), pp. 71-158, 2012. DOI: 10.1590/S1516-635X2012000200006
- [2] Paulino, M.T.F., Oliveira, G.D. and Toledo, J.B., Criação de frangos de corte e acondicionamento térmico em suas instalações: Revisão. *PUBVET*, 13(2), pp. 1-14, 2019. DOI: 10.31533/pubvet.v13n3a280.1-14
- [3] Nascimento, G.R., Nääs, I.A., Baracho, M.S., Pereira, D.F. and Neves, D.P., Termografia infravermelho na estimativa de conforto térmico de frangos de corte. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 18(6), pp. 658-663, 2014. DOI: 10.1590/S1415-43662014000600014
- [4] Santos, M.P., Vale, M.M., Branco, T., Klein, D.R. and Santos, J.P.A., Heat stress in broilers and the need of climatization systems/estresse por calor em frangos de corte e a necessidade de sistemas de climatização. *Revista Brasileira de Engenharia de Biossistemas*, 11(3), pp. 265-272, 2017. DOI: 10.18011/bioeng2017v11n3p265-272
- [5] Menegali, L., Tinoco, I.F.F., Carvalho, C.C.S., Souza, C.F. and Martins, J.H., Comportamento de variáveis climáticas em sistemas de ventilação mínima para produção de pintos de corte. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 17(1), pp. 106-113, 2013. DOI: 10.1590/S1415-43662013000100015
- [6] Souza, V.L.F., Buranelo, G.S., Gasparino, E., Cardozo, R.M. and Barbosa, M.J.B., Efeito da automatização nas diferentes estações do ano sobre os parâmetros de desempenho, rendimento e qualidade da carne de frangos de corte. *Acta Scientiarum. Animal Sciences*, 32(2), pp. 175-181, 2010. DOI: 10.4025/actascianimsci.v32i2.7241
- [7] Cordeiro, M.B., Freitas, H.J., Aquino, E.O. and Sousa, E.M., Avaliação do estresse térmico em frangos caipiras criados em condições climáticas do estado do Acre. *Revista Enciclopédia Biosfera*. 10(19), pp. 358-365, 2014.
- [8] Schiassi, L., Yanagi Junior, T., Reis, G.M., Abreu, L.H.P., Campos, A.T. and Castro, J.O., Modelagem fuzzy aplicada na avaliação do desempenho de frangos de corte. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 19(2), pp. 140-146, 2015. DOI: 10.1590/1807-1929/agriambi.v19n2p140-146.
- [9] Nazareno, A.C., Pandorfi, H., Almeida, G.L.P., Giongo, P.R., Pedrosa, E.M.R. and Guiselini, C., Avaliação do conforto térmico e desempenho de frangos de corte sob regime de criação diferenciado. *Revista Brasileira de Engenharia Agrícola Ambiental*, 13(6), pp. 802-808, 2009. DOI: 10.1590/S1415-43662009000600020
- [10] Cassuce, D.C., Determinação das faixas de conforto térmico para frangos de corte de diferentes idades criados no Brasil, PhD. dissertation Dr., Universidade Federal de Viçosa, Brazil, 2011, 91 P.
- [11] Baracho, M.S., Cassiano, J.A., Nääs, I.A, Tonon, G.S., Garcia, R.G., Royer, A.F.B., Moura, D.J. and Santana, M.R., Ambiente interno em galpões de frango de corte com cama nova e reutilizada. *Revista Agrarian*, 6(22), pp. 473-478, 2013. DOI: 10.30612/agrarian.v6i22.1974
- [12] Rocha, H.P., Furtado, D.A., Nascimento, J.W.B. and Silva, J.H.V., Índices bioclimáticos e produtivos em diferentes galpões avícolas no semiárido paraibano. *Revista Brasileira de Engenharia Agrícola e Ambiental*, 14(12), pp. 1330-1336, 2010. DOI: 10.1590/S1415-43662010001200012
- [13] Cordeiro, M.B., Tinoco, I.F.F., Silva, J.N., Vigoderis, R.B., Pinto,

- F.A.C. and Cecon, P.R., Conforto térmico e desempenho de pintos de corte submetidos a diferentes sistemas de aquecimento no período de inverno. *Revista Brasileira de Zootecnia*, 39(1), pp. 217-224, 2010. DOI: 10.1590/S1516-35982010000100029
- [14] Arcila, P.J.C., Tinoco, I.F.F., Saraz, J.A.O., Rocha, K.S.O. and Candido, M.G.L., Zootechnical and physiological performance of broilers in the final stage of growth subjected to different levels of heat stress. *Revista Facultad Nacional de Agronomía Medellín*, 71(2), pp. 8469-8476, 2018. DOI: 10.15446/rfna.v71n2.71927
- [15] Cândido, M.G., Tinôco, I.F.F., Pinto, F.A.C., Santos, N.T. and Roberti, R.P., Determination of thermal comfort zone for early-stage broilers. *Engenharia Agrícola*, 36(5), pp. 760-767, 2016. DOI: 10.1590/1809-4430-Eng.Agric.v36n5p760-767/2016
- [16] Zhang, S.S., Su, H.G., Zhou, Y., Li, X.M., Feng, J.H. and Zhang, M.H., Effects of sustained cold and heat stress on energy intake, growth and mitochondrial function of broiler chickens. *Journal of Integrative Agriculture*, 15(10), pp. 2336-2342, 2016. DOI: 10.1016/S2095-3119(15)61314-4
- [17] Cassuce, D.C., Tinôco, I.F.F., Baêta, F.C., Zolnier, S., Cecon, P.R. and Vieira, M.D.F., Thermal comfort temperature update for broiler chickens up to 21 days of age. *Engenharia Agrícola*, 33(1), pp. 28-36, 2013. DOI: 10.1590/S0100-69162013000100004
- [18] Al-Zghoul, M.B., Dalab, A.E.S., Yahya, I.E., Althnaian, T.A., Al-ramadan, S.Y., Ali, A.M., Albokhadaim, I.F., El-Bahr, S.M., Al-Busadah, K.A. and Hannon, K.M., Thermal manipulation during broiler chicken embryogenesis: Effect on mRNA expressions of Hsp108, Hsp70, Hsp47 and Hsf-3 during subsequent post-hatch thermal challenge. *Research in veterinary science*, 103, pp. 211-217, 2015. DOI: 10.1016/j.rvsc.2015.10.015
- [19] Rostagno, H.S., Albino, L.F.T., Donzele, J.L., Gomes, P.C., Oliveira, R.F., Lopes, D.C., Ferreira, A.S., Barreto, S.L.T. and Euclides, R.F., Tabelas brasileiras para aves e suínos: composição de alimentos e exigências nutricionais, Universidade Federal de Viçosa, 3ª ed. Viçosa, Brasil, 2011, 252 P.
- [20] Medeiros, C.M., Baêta, F.D.C., Oliveira, R.D., Tinôco, I.D.F.F., Albino, L.F.T. and Cecon, P.R., Efeitos da temperatura, umidade relativa e velocidade do ar em frangos de corte. *Engenharia na Agricultura*, 13(4), pp. 277-286, 2005.
- [21] Vantress, C., Cobb broiler management guide, Siloam Springs, Arkansas, USA, 2013, 73 P.
- [22] Cony, A.V. and Zocche, A.T., Manejo de frangos de corte, in: Mendes, A., Naas, I.A. and Macari, M., Produção de frangos de corte. Facta Ed., Campinas, Brasil, 2004, pp. 118-136.
- [23] Lourençoni, D., Yanagi-Junior, T., Abreu, P.G.D., Campos, A.T. and Yanagi, S.D.N., Productive responses from broiler chickens raised in different commercial production systems-part i: fuzzy modeling. *Engenharia Agrícola*, 39(1), pp. 1-10, 2019. DOI: 10.1590/1809-4430-Eng.Agric.v39n1p1-10/2019
- [24] Curtis, S.E., Environmental management in animal agriculture. The Iowa State University, Ames, USA, 1983, 410 P.
- [25] Ferreira, D.F., Programa de análises estatísticas (Statistical Analysis Software) e planejamento de Experimentos-SISVAR 5.3. Universidade Federal de Lavras, Brasil, 2010.
- [26] Schiassi, L., Yanagi-Junior, T., Damasceno, F.A., Saraz, J.A.O. and Amaral, A.G., Thermal-acoustic comfort index for workers of poultry houses using fuzzy modeling. *International Journal of Engineering Research and Applications (IJERA)*, 4(9), pp. 60-64, 2014.
- [27] Silva, V.K., Silva, J.D.T., Gravena, R.A., Marques, R.H., Hada, F.H. and Moraes, V.M.B., Desempenho de frangos de corte de 1 a 21 dias de idade alimentados com rações contendo extrato de leveduras e prebiótico e criados em diferentes temperaturas. *Revista Brasileira de Zootecnia*, 38(4), pp. 690-696, 2009. DOI: 10.1590/S1516-35982009000400015
- [28] Oba, A., Lopes, P.C.F., Boiago, M.M., Silva, A.M.S., Montassier, H.J. and Souza, P.A., Características produtivas e imunológicas de frangos de corte submetidos a dietas suplementadas com cromo, criados sob diferentes condições de ambiente. *Revista Brasileira de Zootecnia*, 41(5), pp. 1186-1192, 2012. DOI: 10.1590/S1516-35982012000500016
- [29] Faria Filho, D.E., Efeito de dietas com baixo teor protéico, formuladas usando o conceito de proteína ideal, para frangos de corte criados em temperaturas fria, termoneutra e quente, MSc. Thesis, Faculdade de Ciências Agrárias e Veterinárias, Universidade Estadual Paulista, Brazil, 2003, 85 P.
- [30] Koiyama, N.T.G., Rosa, A.P., Padilha, M.T.S., Boemo, L.S., Scher, A., Melo, A.M.S. and Fernandes, M.O., Desempenho e rendimento de carcaça de frangos de corte alimentados com mistura de aditivos fitogênicos na dieta. *Pesquisa Agropecuária Brasileira*, 49(3), pp. 225-231, 2014. DOI: 10.1590/S0100-204X2014000300009
- [31] Matias, C.F.Q., Rocha, J.S.R., Pompeu, M.A., Baião, R.C., Baião, N.C., Lara, L.J.C., Climaco, W.L.S., Pereira, L.F.P., Caldas, E.O., Teixeira, M.P.F. and Cardeal, P.C., Efeito da protease sobre o coeficiente de metabolizabilidade dos nutrientes em frangos de corte. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 67(2), pp. 492-498, 2015. DOI: 10.1590/1678-7883
- [32] Mendes, A.S., Paixão, S.J., Marostega, J., Restelatto, R., Oliveira, P.A.V. and Possenti, J.C., Mensuração de problemas locomotores e de lesões no coxim plantar em frangos de corte. *Archivos de Zootecnia*, 61(234), pp. 217-228, 2012. DOI: 10.4321/S0004-05922012000200006
- [33] Boiago, M.M., Barba, H., Souza, P.A., Scatolini, A.M., Ferrari, F.B. and Giampietro-Ganeco, A., Desempenho de frangos de corte, alimentados com dietas contendo diferentes fontes de selênio, zinco e manganês, criados sob condições de estresse térmico. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 65(1), pp. 241-247, 2013. DOI: 10.1590/S0102-09352013000100034
- [34] Innocentini, R.C.P., Análise dos custos de produção de frangos de corte nos sistemas integrado e independente-comunicação. *Veterinária Notícias*, 15(2), pp. 9-18, 2009.
- [35] Fernandes, R.T.V., Vasconcelos, N.V.B., Lopes, F.F., Arruda, A.M.V. and Pinto, A.R.M., Aspectos gerais sobre alimentos alternativos na nutrição de aves. *Revista Verde de Agroecologia e Desenvolvimento Sustentável*, 7(5), pp. 67-72, 2012.
- [36] Vantress, C. Suplemento de nutrição e desempenho do frango de corte Cobb 500, 2015, 14 P.
- [37] Carvalho, L.S.S., Nutrição de poedeiras em clima quente. *Revista Científica Eletrônica de Medicina Veterinária*, 18, pp. 1-15, 2012.

L.H.P. Abreu, has a BSc. in Agricultural Engineering in 2012, from the Universidade Federal de Lavras - UFLA, Brazil, a MSc. in 2014 and a Dr. in 2015, in Rural Constructions and Environment by the same institution. PhD in Engineering of Work Safety in 2014, by the Centro Universitário de Lavras - Unilavras, Brazil. He is currently professor at the Universidade Federal de Lavras, where he develops activities in the field of Agricultural Engineering and Civil Engineering with emphasis on rural constructions and work safety.
ORCID: 0000-0002-5295-0177

T. Yanagi Junior, he BSc. in Agricultural Engineering in 1992 from the Universidade Federal de Lavras - UFLA, Brazil, where he holds a MSc. in 1995 from the same intuition, and a PhD in Agricultural Engineering in 2002 from the Universidade Federal de Viçosa, Brazil, with sandwich training at Iowa State University of Science and Technology, USA, where he served as a visiting researcher. He is currently professor at the Universidade Federal de Lavras, where he has experience in Agricultural Engineering and Systems Engineering, especially in the area of construction and ambience, in which he works to guide undergraduate and graduate students in the development of basic and applied research, involving evaluation and mathematical modeling of biosystems.
ORCID: 0000-0001-9653-205X

M. Bahuti, is a BSc. in Agricultural Engineering in 2018, from the Universidade Federal de Lavras - UFLA, Brazil. He currently holds a MSc. in Agricultural Engineering for the same intuition with a concentration area in Construction and Environment. His research interest includes: mathematical modeling, performance evaluation and analysis of the breeding environment for animals.
ORCID: 0000-0002-1542-5058

R.R. Lima, he BSc. in Zootechnics in 1993, from the Universidade Federal de Lavras - UFLA, Brazil, MSc. in Genetics and Improvement in 1997, by the Universidade Federal de Viçosa, PhD in Agronomy / Statistics and

Agronomic Experimentation in 2005, by the Escola Superior de Agricultura Luiz de Queiroz, Brazil, with sandwich training at the University of Kent at Canterbury, England, and PhD fellow at the University of Wisconsin, Madison, WI, United States, in Genomic Statistics. He is currently professor of the Universidade Federal de Lavras, Brazil, where he develops activities in the area of probability and statistics, with emphasis on probability and applied statistics, working mainly on the following topics: spatial statistics, agricultural experimentation and genomic statistics.
ORCID:0000-0003-4607-4964

D. Lourençoni, he is a BSc. in Agricultural Engineering in 2011, from the Universidade Federal de Lavras - UFLA, Brazil. He holds a MSc. in 2013 and a PhD. In 2017, in Agricultural Engineering from the same institution in the area of Construction and Environment. He is currently a professor at the Universidade Federal do Vale do São Francisco - UNIVASF, Brazil, participating in the postgraduate program in Agricultural Engineering, developing research on rural structures and buildings, unconventional materials, biosystems environment, artificial intelligence and thermal efficiency.
ORCID:0000-0003-1173-2381

E.J. Fassani, he BSc. in Zootechnics in 1994, from the Universidade Federal de Lavras - UFLA, Brazil. He holds a MSc. in 1998 and PhD in 2003 in Zootechnics, by the same institution, in the area of Animal Nutrition and Feeding. He is currently a professor at the Universidade Federal de Lavras where he develops activities in the area of Zotechnology, with emphasis on nutrition of monogastric and poultry, working mainly with nutrition (evaluation of food and determination of nutritional requirements) in poultry (laying hens, broilers and coturniculture).
ORCID: 0000-0002-7101-3998



UNIVERSIDAD NACIONAL DE COLOMBIA

SEDE MEDELLÍN
FACULTAD DE MINAS

Área Curricular de Recursos Minerales

Oferta de Posgrados

Maestría en Ingeniería - Recursos Minerales
Especialización en Recursos Minerales
Especialización en Gestión del Negocio Minero

Mayor información:

E-mail: acremin_med@unal.edu.co
Teléfono: (57-4) 425 53 68