

A TRIANGLE INEQUALITY FOR ANGLES IN A HILBERT SPACE

by

D. K. RAO

Let x, y, z be unit vectors in a Hilbert space, and define the angle θ_{xy} by $\cos \theta_{xy} = \Re e(x, y)$, $0 \leq \theta_{xy} \leq \pi$. The object this note is to give a proof of the following inequality

$$(1) \quad \theta_{xz} \leq \theta_{xy} + \theta_{yz} .$$

This result was mentioned without proof in [1] and is important in inequalities for operator cosines introduced in [2].

In order to prove (1) we need the following lemma :

LEMMA. Let x, y, z be unit vectors in a Hilbert space, and $(x, y) = a_1 + i b_1$, $(y, z) = a_2 + i b_2$, $(x, z) = a_3 + i b_3$. Then

$$(2) \quad \cos \theta_{xz} \geq \cos (\theta_{xy} + \theta_{yz}) .$$

Proof. By Schwarz inequality we have $|a_j|^2 + |b_j|^2 \leq 1$, $j=1, 2, 3$. On the other hand, (2) is equivalent to

$$(1-a_1^2)^{\frac{1}{2}} (1-a_2^2)^{\frac{1}{2}} \geq a_1 a_2 - a_3 .$$

This result is obvious if $a_1 a_2 - a_3 \leq 0$. Otherwise we need to prove that

$$1 - a_1^2 - a_2^2 - a_3^2 + 2a_1 a_2 a_3 \geq 0.$$

(It is interesting to note that the above expression does not contain b_1, b_2 or b_3 .) For this let $f(p, q, r) = 1 - p^2 - q^2 - r^2 + 2pqr$, so that

$$p \frac{\partial f}{\partial p} + q \frac{\partial f}{\partial q} + r \frac{\partial f}{\partial r} = -2[p^2 + q^2 + r^2 - 3pqr].$$

In the cube $E = \{ |p| \leq 1, |q| \leq 1, |r| \leq 1 \}$, we have $|pqr| \leq |pq|$, etc, and hence

$$2[p^2 + q^2 + r^2 - 3pqr] \geq (|p| - |q|)^2 + (|q| - |r|)^2 + (|p| - |r|)^2 \geq 0.$$

Therefore

$$p \frac{\partial f}{\partial p} + q \frac{\partial f}{\partial q} + r \frac{\partial f}{\partial r} \leq 0 \text{ in } E.$$

The above inequalities show that for any rectangle V contained in E , f attains its minimum value on the surface of V . In particular, this is true for the cube E . Now consider the Gramian

$$\begin{vmatrix} (x,x) & (x,y) & (x,z) \\ (y,x) & (y,y) & (y,z) \\ (z,x) & (z,y) & (z,z) \end{vmatrix} \geq 0,$$

or equivalently,

$$1 - a_1^2 - a_2^2 - a_3^2 + 2a_1 a_2 a_3 \geq b_1^2 + b_2^2 + b_3^2 + 2b_1(b_2 a_3 - b_3 a_2) - 2a_1 b_2 b_3.$$

If we take (a_1, a_2, a_3) on the surface of E and assume, for example, that $|a_1| = 1$, then $b_1 = 0$ and we have

$$1 - a_1^2 - a_2^2 - a_3^2 + 2a_1 a_2 a_3 \geq b_2^2 + b_3^2 - 2a_1 b_2 b_3 \geq b_1^2 + b_2^2 - 2|b_2||b_3| = (|b_2| - |b_3|)^2 \geq 0,$$

that is to say the desired inequality.

Q.E.D.

Proof of inequality (1). If $\theta_{xy} + \theta_{yz} \geq \pi$, there is nothing to prove. If $\theta_{xy} + \theta_{yz} < \pi$, it follows from (2) and the fact that cosine is a non-increasing function on $[0, \pi)$ that inequality (1) holds.

References

1. M. Krein : *Angular localization of a multiplicative integral in a Hilbert space*, *Funct. Anal. Applic.* 3 (1969).
2. K. Gustafson : *The angle of an operator and positive operator products*, *Bull. Amer. Math. Soc.* 74 (1968), 188-192.

Departamento de Matemáticas
Universidad del Valle
Apartado Aéreo 2188
Cali, Colombia, S. A.

(Recibido en abril de 1976).