

# Global well-posedness for two dimensional semilinear wave equations

GERMÁN E. FONSECA

Universidad Nacional de Colombia, Bogotá

**ABSTRACT.** We consider the initial value problem (IVP) for certain semilinear wave equations in two dimensions. It is shown that global well-posedness holds in spaces of lower regularity than that suggested by the energy space  $\dot{H}^1 \times L_x^2$ . The technique to be used is adapted from a general scheme originally introduced by J. Bourgain to establish global well posedness of the cubic nonlinear Schrödinger equation.

**Key words and phrases.** Nonlinear wave equations, global solutions, initial value problems.

**1991 Mathematics Subject Classification.** Primary 35L15. Secondary 35A05.

## 1. Introduction

We consider the initial value problem (IVP) for the semilinear wave equation

$$\begin{cases} \partial_t^2 u - \Delta_x u = -|u|^{k-1}u, & t \in \mathbb{R}, x \in \mathbb{R}^2, \\ u(0, x) = f(x), \\ u_t(0, x) = g(x), \end{cases} \quad (1.1)$$

where  $k \geq 3$ .

Our purpose in this paper is to establish the global well-posedness of this IVP in spaces of lower regularity than that dictated by the respective conservation law

$$\int_{\mathbb{R}^2} (u_t)^2 + (\nabla u)^2 + \frac{2}{k+1} |u|^{k+1} = \text{constant} \quad (1.2)$$

and that, in addition with the existent local theory, guarantee global well-posedness of this IVP in the energy space.

The notion of local well-posedness for IVP (1.1) in a function space  $X$  includes local existence, uniqueness, continuous dependence upon the data, and persistence; i.e., that for  $u_0 \in X$ , the corresponding solution describes a continuous curve in  $X$ . The global well-posedness requires, in addition, that the notion above holds for every time interval. In this regard we have the following local well-posedness result.

**Theorem 1.1.** *Let  $s \in (s(k), 1]$  with  $s(k)$  given by*

$$s(k) = \begin{cases} \frac{3k-7}{4(k-1)}, & \text{if } 3 \leq k \leq 5, \\ \frac{k-3}{k-1}, & \text{if } k \geq 5. \end{cases} \quad (1.3)$$

*Then IVP (1.1) is locally well-posed in  $\dot{H}^s(\mathbb{R}^2) \times \dot{H}^{s-1}(\mathbb{R}^2)$ . Furthermore, the time  $T$  of existence of the solution depends on the size of the initial data; i.e.,  $T = T(\|f\|_{\dot{H}^s(\mathbb{R}^2)}, \|g\|_{\dot{H}^{s-1}(\mathbb{R}^2)})$ .*

The proof of this theorem is carried out in [8], where three distinguished cases are considered: the conformal case  $k = 5$  and the superconformal and subconformal cases  $k > 5$  and  $k < 5$ , respectively. It is worth mentioning that this result is sharp in the sense that for initial data with lower regularity than that in (1.3), IVP (1.1) is ill posed (see [7], [8]). Notice that in the conformal and superconformal cases, the required regularity for the initial data to obtain local well posedness is the one suggested by the scaling argument: if  $u(x, t)$  solves IVP (1.1) then  $\lambda^{\frac{2}{k-1}} u(\lambda x, \lambda t)$  also solves the equation in (1.1), and there is invariance of the  $\dot{H}^s \times \dot{H}^{s-1}$  norm of the initial data exactly for  $s = \frac{k-3}{k-1}$ .

It follows from this theorem and from the conservation law in (1.2) that IVP (1.1) is globally well-posed in  $\dot{H}^1 \cap L_x^{k+1} \times L^2$ .

Our main result can now be stated:

**Theorem 1.2.** *Let  $s \in (s(k), 1]$  with  $s(k)$  given by*

$$s(k) = \begin{cases} \frac{k-2}{k-1}, & \text{if } 3 \leq k \leq 5, \\ \frac{k-1}{k}, & \text{if } k > 5. \end{cases} \quad (1.4)$$

Then, for any  $(u_0, u_1) \in \dot{H}^s(\mathbb{R}^2) \cap L_x^{k+1} \times \dot{H}^{s-1}(\mathbb{R}^2)$  IVP (1.1) has a unique global solution  $u(t)$ . For a given time interval  $[0, T]$ ,  $u$  can be expressed as

$$u(t) = \partial_t W(t)u_0 + W(t)u_1 + I(t), \quad (1.5)$$

with  $I$  more regular than the initial data, more precisely,  $I(t) \in \dot{H}^1(\mathbb{R}^2)$  and satisfies the polynomial growth condition

$$\sup_{t \in [0, T]} \|I(t)\|_{\dot{H}^1} \leq c T^{\frac{1-s}{(k-1)s-(k-2)}+}, \quad (1.6)$$

and  $W(t)$  is defined below.

We restrict our attention to two dimensions, but the same problem can be studied in other dimensions as well. In fact, Kenig, Ponce and Vega [5] obtain similar results for the three dimensional case, but a much more restrictive range for the nonlinearity has to be imposed, namely,  $2 \leq k < 5$ . This is a required condition in the local theory for such IVP's in three dimensions.

Our proof follows ideas recently introduced by Bourgain [1], where he obtained that the IVP associated with the two dimensional cubic Schrödinger is globally well-posed in Sobolev spaces of indexes in between those where the conservation laws hold. He also applied it for the PBVP for the Klein-Gordon equation (see [2]). This method has been extensively applied to many other equations as well (see [3], [4], [5]).

The sketch of the proof is the following: we split the initial data following the ideas in [1], that is,  $u_0(x) = u_{0,1} + u_{0,2}$  and  $u_1(x) = u_{1,1} + u_{1,2}$ , where  $(u_{0,1}, u_{1,1}) \in \dot{H}^1 \cap L_x^{k+1} \times L_x^2$  and  $(u_{0,2}, u_{1,2}) \in \dot{H}^s \cap L_x^{k+1} \times \dot{H}^{s-1}$ . Then we consider IVP (1.1) with data  $(u_{0,1}, u_{1,1})$ , for which the global theory allows to obtain a solution,  $v(t)$ , defined in an interval  $[0, \Delta T]$  with  $\Delta T$  an appropriate time. Next, we consider an IVP with variable coefficients for  $z(t) = u(t) - v(t)$  whose solution,  $z(t)$ , is also defined in  $[0, \Delta T]$  and can additionally be expressed, via Duhamel's principle, as  $z(t) = y(t) + I(t)$ , with  $I(t)$  more regular than the initial data, more precisely,  $(\nabla I, \partial_t I, I) \in L_x^2 \times L_x^2 \times L_x^{k+1}$ . Then the argument is re-applied in the interval  $[\Delta T, 2\Delta T]$  having initial data  $(v(\Delta T) + I(\Delta T), \partial_t v(\Delta T) + \partial_t I(\Delta T))$  and  $(y(\Delta T), \partial_t y(\Delta T))$  for their respective IVP's. The process is continued until reaching any given time  $T \gg 1$ . Since at each step the Sobolev norms involved grow, we have to obtain some estimates to keep the process uniform.

## Notations.

- $D$  denotes the homogeneous derivative  $\sqrt{-\Delta_x}$ .
- $W(t)$  denotes the operator  $W(t)g = \frac{\sin(Dt)}{D}g$ , which solves the linear wave equation with initial data  $(0, g)$ .
- $\|\cdot\|$  denotes the  $L_x^2$  norm.

- The mixed space-time norm is denoted by

$$\|f\|_{L_T^q L_x^p} = \left( \int_0^T \left( \int_{-\infty}^{\infty} |f(x, t)|^p dx \right)^{q/p} dt \right)^{1/q}.$$

In case that  $T = \infty$ , we will use  $\|\cdot\|_{L_t^q L_x^p}$ .

- The letter  $c$  denotes a constant that may change from line to line.

## 2. Proof of Theorem 1.2

Before we proceed with the proof of our main result, we state some Strichartz estimates for solutions of the linear wave IVP in two dimensions given by

$$\begin{cases} \partial_t^2 u - \Delta_x u = 0, & t \in \mathbb{R}, x \in \mathbb{R}^2, \\ u(0, x) = u_0(x), & u_t(0, x) = u_1(x). \end{cases} \quad (2.1)$$

**Theorem 2.1.** *Let  $\theta \in [0, 3/4)$ ,  $\gamma \in \mathbb{R}$ . Then, for any pair of functions  $(f, g) \in \dot{H}^\gamma(\mathbb{R}^2) \times \dot{H}^{\gamma-1}(\mathbb{R}^2)$ , the solution of IVP (2.1) satisfies*

$$\|D^{\gamma-\theta} u\|_{L_t^{\frac{3}{\theta}} L_x^{\frac{6}{3-4\theta}}} \leq c_\theta (\|D^\gamma u_0\| + \|D^{\gamma-1} u_1\|). \quad (2.2)$$

The proof of this estimate and of its versions in higher dimensions can be found in [9].

Let us start with the proof of our Theorem 1.2. We split the initial data  $(u_0, u_1) \in \dot{H}^s \cap L_x^{k+1} \times \dot{H}^{s-1}$  following the scheme in [1], that is,

$$u_0(x) = (\chi_{\{|\xi| < N\}} \hat{f})^\vee(x) + (\chi_{\{|\xi| \geq N\}} \hat{u}_0)^\vee(x) = u_{0,1}(x) + u_{0,2}(x) \quad (2.3)$$

and

$$u_1(x) = (\chi_{\{|\xi| < N\}} \hat{f})^\vee(x) + (\chi_{\{|\xi| \geq N\}} \hat{u}_0)^\vee(x) = u_{1,1}(x) + u_{1,2}(x), \quad (2.4)$$

where  $N$  is a large number to be chosen later.

From this splitting it follows that

$$\|D^\rho u_{0,1}\|, \|D^{\rho-1} u_{1,1}\| \leq cN^{\rho-s} \quad \text{for } \rho \geq s, \quad (2.5)$$

and

$$\|D^\rho u_{0,2}\|, \|D^{\rho-1} u_{1,2}\| \leq cN^{\rho-s} \quad \text{for } 0 \leq \rho \leq s. \quad (2.6)$$

As we already pointed out, for initial data  $u_{0,1}, u_{1,1}$ , IVP (1.1) has a unique global solution in  $\dot{H}^1 \cap L_x^{k+1} \times L^2$ . Moreover, this solution is expressed via Duhamel's principle in the integral form

$$v(t) = \partial_t W(t)u_{0,1} + W(t)u_{1,1} - \int_0^t W(t-t')(|v|^{k-1}v)(t') dt', \quad (2.7)$$

and it satisfies the conservation law (1.2), which in terms of the size of the initial data yields the estimate

$$\|v_t(t)\| + \|\nabla v(t)\| + \|v(t)\|_{L_x^{\frac{k+1}{2}}}^{\frac{k+1}{2}} \sim N^{1-s}. \quad (2.8)$$

Next, we consider the IVP with variable coefficients given by

$$\begin{cases} \partial_t^2 z - \Delta_x z = -|z+v|^{k-1}(z+v) + |v|^{k-1}v, & t \in \mathbb{R}, x \in \mathbb{R}^2, \\ z(0, x) = u_{0,2}(x) \in \dot{H}^s, \\ z_t(0, x) = u_{1,2}(x) \in \dot{H}^{s-1}, \end{cases} \quad (2.9)$$

with the initial data satisfying

$$\|D^\rho u_{0,2}\|, \|D^{\rho-1} u_{1,2}\| \leq cN^{\rho-s} \quad \text{for } 0 \leq \rho \leq s,$$

and its integral version

$$z(t) = \partial_t W(t)u_{0,1} + W(t)u_{1,1} - \int_0^t W(t-t')F(t') dt', \quad (2.10)$$

where

$$F = -|z+v|^{k-1}(z+v) + |v|^{k-1}v. \quad (2.11)$$

We will obtain the following estimates for the size of the solution of IVP (2.9) in terms of the size of the initial data. The local well posedness of IVP (2.9) easily follows from these estimates and the contraction principle applied to the integral equation (2.10). For the sake of simplicity, the details are omitted.

First, we consider the conformal and subconformal cases  $3 \leq k \leq 5$ . Let us define

$$\|z\|_\gamma = \|D^\gamma z\|_{L_{\Delta T}^\infty L_x^2} + \|z\|_{L_{\Delta T}^{3/\gamma} L_x^{6/(3-4\gamma)}}, \quad (2.12)$$

with  $\frac{3k-6}{4k-3} \leq \gamma \leq s$ .

From Strichartz estimate (2.2) and (2.5) we obtain

$$\begin{aligned} \|z\|_{L_{\Delta T}^{3/\gamma} L_x^{6/(3-4\gamma)}} &\leq c(\|D^\gamma u_{0,2}\| + \|D^{\gamma-1} u_{1,2}\|) + c \int_0^{\Delta T} \|D^{\gamma-1}(F(t'))\| dt' \\ &\leq cN^{\gamma-s} + c \int_0^t \|D^{\gamma-1} F(t')\| dt' \\ &\leq cN^{\gamma-s} + c\|F\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}}. \end{aligned} \quad (2.13)$$

In order to estimate the last term in (2.13), we observe that from (2.11) it is enough to estimate the terms  $F_1 = |z|^k$  and  $F_k = |z||v|^{k-1}$ .

For  $F_1$  we have

$$\begin{aligned}
 \|F_1\|_{L^1_{\Delta T} L^2_x} &\leq \|z\|_{L^k_{\Delta T} L^{2k/(2-\gamma)}_x}^k \\
 &\leq c \|D^\gamma z\|_{L^k_{\Delta T} L^2_x}^{\frac{6-3k-\gamma(3-4k)}{\gamma}} \|z\|_{L^{\frac{3k-6+\gamma(3-3k)}{\gamma/(3-4\gamma)}}_{\Delta T} L^6_x}^{\frac{3k-6+\gamma(3-3k)}{\gamma/(3-4\gamma)}} \\
 &\leq c \Delta T^{3-k+\gamma(k-1)} \|D^\gamma z\|_{L^\infty_{\Delta T} L^2_x}^{\frac{6-3k-\gamma(3-4k)}{\gamma}} \|z\|_{L^{\frac{3k-6+\gamma(3-3k)}{3/\gamma} L^{\frac{6}{\gamma/(3-4\gamma)}}_x}^{\frac{3k-6+\gamma(3-3k)}{\gamma/(3-4\gamma)}} \\
 &\leq c \Delta T^{3-k+\gamma(k-1)} \|z\|_\gamma^k,
 \end{aligned} \tag{2.14}$$

and similarly for  $F_k$ ,

$$\begin{aligned}
 \|F_k\|_{L^1_{\Delta T} L^2_x} &\leq \|zv^{k-1}\|_{L^1_{\Delta T} L^{2/(2-\gamma)}_x} \\
 &\leq c \Delta T \|z\|_{L^\infty_{\Delta T} L^{2/(1-\gamma)}_x} \|v\|_{L^{\frac{k-1}{2}}_{\Delta T} L^{2(k-1)}_x}^{k-1} \\
 &\leq c \Delta T \|D^\gamma z\|_{L^\infty_{\Delta T} L^2_x} \|Dv\|_{L^{\frac{k-3}{2}}_{\Delta T} L^2_x}^{\frac{k-3}{2}} \|v\|_{L^{\frac{k+1}{2}}_{\Delta T} L^{(k+1)}_x}^{\frac{k+1}{2}} \\
 &\leq c \Delta T N^{\frac{(k-3)(1-s)}{2}} N^{1-s} \|z\|_\gamma \leq c \Delta T N^{\frac{(k-1)(1-s)}{2}} \|z\|_\gamma.
 \end{aligned} \tag{2.15}$$

We remark that in order to incorporate the appropriate time space in (2.14), the above given restriction on  $k$  is needed.

The estimate for  $\|D^\gamma z\|_{L^\infty_{\Delta T} L^2_x}$  is exactly the same as in (2.13). Hence, it follows from (2.13), (2.14) and (2.15) and from the choice of  $\Delta T$  that for any  $\gamma \in (\frac{3k-6}{4k-3}, s)$ ,

$$\|z\|_\gamma \leq c N^{\gamma-s}. \tag{2.16}$$

It is also important to estimate the  $L^2_x$  norm of the solution of IVP (2.9). Thus, from (2.6) and Sobolev's inequality,

$$\|z\|_{L^\infty_{\Delta T} L^2_x} \leq \|u_{0,2}\| + \|u_{1,2}\| + \|D^{-1}F\|_{L^1_{\Delta T} L^2_x} \leq c N^{-s} + c \|F\|_{L^1_{\Delta T} L^1_x}. \tag{2.17}$$

To estimate  $F_1$  we apply (2.16) and a Gagliardo-Nirenberg type inequality, to obtain that

$$\begin{aligned}
 \|F_1\|_{L^1_{\Delta T} L^1_x} &\leq \Delta T \|z^k\|_{L^\infty_{\Delta T} L^1_x} \leq c \Delta T \|z\|_{L^\infty_{\Delta T} L^2_x}^k \\
 &\leq c \Delta T (\|z\|_{L^\infty_{\Delta T} L^2_x}^{\frac{1}{k}} \|D^{\frac{k-2}{k-1}} z\|_{L^\infty_{\Delta T} L^2_x}^{\frac{k-1}{k}})^k \\
 &\leq c \Delta T \|z\|_{L^\infty_{\Delta T} L^2_x} \|D^\gamma z\|_{L^\infty_{\Delta T} L^2_x}^{k-1} \\
 &\leq c \Delta T \|z\|_{L^\infty_{\Delta T} L^2_x} \|z\|_\gamma^{k-1} \\
 &\leq c N^{-\frac{(k-1)(1-s)}{2}} N^{(k-1)(\gamma-s)} \|z\|_{L^\infty_{\Delta T} L^2_x} \\
 &\leq c N^{\frac{(k-3)}{2} - \frac{(k-1)}{2}s} \|z\|_{L^\infty_{\Delta T} L^2_x},
 \end{aligned} \tag{2.18}$$

where  $\gamma = \frac{k-2}{k-1}$ . Since  $N$  is large and the exponent is negative, this part can be absorbed into the left hand side of (2.17).

In a similar manner, we have for  $F_k$

$$\begin{aligned} \|F_k\|_{L^1_{\Delta T} L^1_x} &\leq \|zv^{k-1}\|_{L^1_{\Delta T} L^1_x} \leq \Delta T \|z\|_{L^\infty_{\Delta T} L^2_x} \|v\|_{L^\infty_{\Delta T} L^{2(k-1)}_x} \\ &\leq c\Delta T \|z\|_{L^\infty_{\Delta T} L^2_x} \|Dv\|_{L^\infty_{\Delta T} L^2_x}^{\frac{k-3}{2}} \|v\|_{L^\infty_{\Delta T} L^{k+1}_x}^{\frac{k+1}{2}} \\ &\leq c\Delta T N^{\frac{(k-3)(1-s)}{2}} N^{1-s} \|z\|_{L^\infty_{\Delta T} L^2_x} \\ &\leq c\Delta T N^{-\frac{(k-1)(1-s)}{2}} \|z\|_{L^\infty_{\Delta T} L^2_x}, \end{aligned} \quad (2.19)$$

and therefore, from the choice of  $\Delta T$ , this term can also be absorbed into the left hand side of (2.17). Summing up, we have the estimate

$$\|z\|_{L^\infty_{\Delta T} L^2_x} \leq cN^{-s}. \quad (2.20)$$

Now it comes the nontrivial part of showing that the integral part  $I(t) = \int_0^t W(t-t')F(t')dt'$  in (2.10) is more regular than the initial data. More precisely,  $\partial_t I(t)$  and  $\nabla I(t)$  live in  $L^2_x$ , whereas the initial data are merely in  $\dot{H}^s \times \dot{H}^{s-1}$  with  $s < 1$ . Furthermore, the estimates in (2.16) and (2.20) will allow to measure the size of this piece of the solution of the IVP (2.9). In fact, Minkowski's inequality yields

$$\|(\partial_t I, \nabla I)(t)\|_{L^2_x} \leq \|F\|_{L^1_{\Delta T} L^2_x}. \quad (2.21)$$

For  $F_1$  we have

$$\begin{aligned} \|F_1\|_{L^1_{\Delta T} L^2_x} &\leq \|z^k\|_{L^1_{\Delta T} L^2_x} \leq \|z\|_{L^k_{\Delta T} L^{2k}_x}^k \leq \Delta T^{\frac{5-k}{4}} \|z\|_{L^{4k/(k-1)}_{\Delta T} L^{2k}_x}^k \\ &\leq \Delta T^{\frac{5-k}{4}} \|z\|_{L^{3/\gamma}_{\Delta T} L^{6/(3-4\gamma)}_x}^k \leq \Delta T^{\frac{5-3k}{4}} \|z\|_{\gamma}^k, \end{aligned} \quad (2.22)$$

where we have chosen  $\gamma = \frac{3k-3}{4k}$ .

Then, the choice of  $\Delta T$  and (2.16) give

$$\begin{aligned} \|F_1\|_{L^1_{\Delta T} L^2_x} &\leq cN^{-\frac{(5-k)(k-1)(1-s)}{8}} N^{k(\frac{3k-3}{4k}-s)} \\ &\leq cN^{\frac{(k-1)(k+1)}{8} - \frac{s}{8}(k^2+2k+5)} \leq cN^{\frac{k-1}{2} - \frac{(k+1)}{2}s}. \end{aligned} \quad (2.23)$$

The last inequality holds since  $s > \frac{k-3}{k-1}$ .

For  $F_k$  we use the Sobolev and a Gagliardo-Nirenberg type inequalities, and the estimates in (2.8), (2.16) and (2.20).

$$\begin{aligned} \|F_k\|_{L^1_{\Delta T} L^2_x} &\leq \|zv^{k-1}\|_{L^1_{\Delta T} L^2_x} \leq \Delta T \|z\|_{L^\infty_{\Delta T} L^{2/(1-2\varepsilon)}_x} \|v\|_{L^\infty_{\Delta T} L^{(k-1)/\varepsilon}_x}^{k-1} \\ &\leq c\Delta T \|D^{2\varepsilon} z\|_{L^\infty_{\Delta T} L^2_x} \|Dv\|_{L^\infty_{\Delta T} L^2_x}^{k-1-(k+1)\varepsilon} \|v\|_{L^\infty_{\Delta T} L^{k+1}_x}^{(k+1)\varepsilon} \\ &\leq cN^{-\frac{(k-1)(1-s)}{2}} N^{2\varepsilon-s} N^{(k-1-(k+1)\varepsilon)(1-s)} N^{2(1-s)\varepsilon} \\ &\leq cN^{\frac{(k-1)}{2} - \frac{(k+1)}{2}s + \varepsilon(2-(k-1)(1-s))}, \end{aligned} \quad (2.24)$$

where  $\varepsilon$  is a small positive number to be chosen.

In order to apply the scheme roughly described in the introduction, we shall also need an estimate on the growth of the  $L_x^{k+1}$  norm of the integral part  $I(t)$  at time  $\Delta T$ . We apply Sovolev's inequality to obtain

$$\begin{aligned} \|I(\Delta T)\|_{L_x^{k+1}} &\leq \int_0^{\Delta T} \left\| \frac{\sin D(\Delta T - t')}{D} F \right\|_{L_x^{k+1}} dt' \\ &\leq c \int_0^{\Delta T} \|D^{\frac{k-1}{k+1}} \frac{\sin D(\Delta T - t')}{D} F\|_{L_x^2} dt' \\ &\leq c \int_0^{\Delta T} \|D^{-\frac{2}{k+1}} F\|_{L_x^2} dt' \leq c \|D^{-\frac{2}{k+1}} F\|_{L_{\Delta T}^1 L_x^2}, \end{aligned} \quad (2.25)$$

and for  $F_1$ , Sovolev's inequality and the estimates in (2.16) with  $\gamma = \frac{(k^2-3)}{k(k+1)}$  yield

$$\begin{aligned} \|D^{-\frac{2}{k+1}} F_1\|_{L_{\Delta T}^1 L_x^2} &\leq \|D^{-\frac{2}{k+1}} |z|^k\|_{L_{\Delta T}^1 L_x^2} \leq \Delta T \|z\|_{L_{\Delta T}^\infty L_x^{2k(k+1)/(k+3)}}^k \\ &\leq c \Delta T \|D^\gamma z\|_{L_{\Delta T}^\infty L_x^2}^k \leq c \Delta T \|z\|_\gamma^k \\ &\leq c N^{-\frac{k-1}{2}(1-s)} N^{(\gamma-s)k} \\ &\leq c N^{-\frac{k-1}{2}(1-s)} N^{(\frac{k^2-3}{k(k+1)}-s)k} \\ &\leq c N^{\frac{k^2-5}{2(k+1)} - \frac{k+1}{2}s} \leq c N^{\frac{k-1}{2} - \frac{(k+1)}{2}s}. \end{aligned} \quad (2.26)$$

The estimates in (2.8) and (2.16) yield for  $F_k$  that

$$\begin{aligned} \|D^{-\frac{2}{k+1}} F_k\|_{L_{\Delta T}^1 L_x^2} &\leq \|D^{-\frac{2}{k+1}} (|z||v|^{k-1})\|_{L_{\Delta T}^1 L_x^2} \\ &\leq c \|z v^{k-1}\|_{L_{\Delta T}^1 L_x^{2(k+1)/(k+3)}} \\ &\leq c \Delta T \|z\|_{L_{\Delta T}^\infty L_x^2} \|v^{k-1}\|_{L_{\Delta T}^\infty L_x^{k+1}} \\ &\leq c \Delta T \|z\|_{L_{\Delta T}^\infty L_x^2} \|v\|_{L_{\Delta T}^\infty L_x^{k+1}}^{k-1} \\ &\leq c \Delta T \|z\|_{L_{\Delta T}^\infty L_x^2} \|D v\|_{L_{\Delta T}^\infty L_x^2}^{k-2} \|v\|_{L_{\Delta T}^\infty L_x^{k+1}} \\ &\leq c N^{-\frac{k-1}{2}(1-s)} N^{-s} N^{(1-s)(k-2)} N^{\frac{2}{k+1}(1-s)} \\ &\leq c N^{\frac{k^2-2k+1}{2(k+1)} - \frac{k^2+3}{2(k+1)}s} \leq c N^{\frac{k-1}{2} - \frac{(k+1)}{2}s}. \end{aligned} \quad (2.27)$$

Estimates (2.21)-(2.27) certainly prove our claim that the integral part in (2.10) is more regular than the initial data, and furthermore, that in the interval  $[0, \Delta T]$  it grows like

$$\|(\partial_t I, \nabla I)(\Delta T)\| + \|I(\Delta T)\|_{L_x^{\frac{k+1}{2}}} \leq c N^{\frac{(k-1)}{2} - \frac{(k+1)}{2}s + \varepsilon(2-(k-1)(1-s))}. \quad (2.28)$$



For higher powers of  $k$  we slightly modify the  $\|\cdot\|$  norm in (2.12) and consider

$$\|z\|_\gamma = \|D^\gamma z\|_{L_{\Delta T}^\infty L_x^2} + \|D^{\gamma-\theta} z\|_{L_{\Delta T}^{3/\theta} L_x^{6/(3-4\theta)}}, \quad (2.29)$$

where  $\gamma > \frac{k-3}{k-1}$  and  $\theta = 3 - 3\gamma + \frac{3\gamma-6}{k}$ .

Once again, Strichartz's estimate yields

$$\begin{aligned} \|D^{\gamma-\theta} z\|_{L_{\Delta T}^{3/\gamma} L_x^{6/(3-4\gamma)}} &\leq c(\|D^\gamma u_{0,2}\| + \|D^{\gamma-1} u_{1,2}\|) + c \int_0^{\Delta T} \|D^{\gamma-1}(F(t'))\| dt' \\ &\leq cN^{\gamma-s} + c\|F\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}}. \end{aligned} \quad (2.30)$$

We proceed as in (2.13)-(2.15) to obtain that

$$\begin{aligned} \|F_1\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}} &\leq \|z^k\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}} \leq \|z\|_{L_{\Delta T}^k L_x^{2k/(2-\gamma)}}^k \\ &\leq c\|D^{\gamma-\theta} z\|_{L_{\Delta T}^k L_x^{6/(3-4\theta)}}^k \\ &\leq c\Delta T^{\frac{3-k\theta}{3}} \|D^{\gamma-\theta} z\|_{L_{\Delta T}^{3/\theta} L_x^{6/(3-4\theta)}}^k \leq c\Delta T^{\frac{3-k\theta}{3}} \|z\|_\gamma^k \end{aligned} \quad (2.31)$$

and

$$\begin{aligned} \|F_k\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}} &\leq \|zv^{k-1}\|_{L_{\Delta T}^1 L_x^{2/(2-\gamma)}} \\ &\leq c\Delta T \|z\|_{L_{\Delta T}^\infty L_x^{2/(1-\gamma)}} \|v\|_{L_{\Delta T}^\infty L_x^{2(k-1)}} \\ &\leq c\Delta T \|D^\gamma z\|_{L_{\Delta T}^\infty L_x^2} \|Dv\|_{L_{\Delta T}^\infty L_x^2}^{\frac{k-3}{2}} \|v\|_{L_{\Delta T}^\infty L_x^{2(k+1)}}^{\frac{k+1}{2}} \\ &\leq c\Delta TN^{-\frac{(k-1)(1-s)}{2}} \|z\|_\gamma. \end{aligned} \quad (2.32)$$

From (2.30)-(2.32), we get

$$\begin{aligned} \|D^{\gamma-\theta} z\|_{L_{\Delta T}^{3/\gamma} L_x^{6/(3-4\gamma)}} &\leq \\ &cN^{\gamma-s} + c\Delta T^{\frac{3-k\theta}{3}} \|z\|_\gamma^k + c\Delta TN^{-\frac{(k-1)(1-s)}{2}} \|z\|_\gamma, \end{aligned} \quad (2.33)$$

and therefore, from the choice of  $\Delta T$  it follows that

$$\|z\|_\gamma \leq cN^{\gamma-s}, \quad (2.34)$$

for  $\gamma \in (\frac{k-3}{k-1}, s)$ .

The estimates for the  $L_{\Delta T}^\infty L_x^2$  norms found in (2.17)-(2.20) for the subconformal and conformal cases are the same for the superconformal case, and therefore

$$\|z\|_{L_{\Delta T}^\infty L_x^2} \leq cN^{-s}. \quad (2.35)$$

The same also holds in the analysis of the smoothing property of the integral part  $I(t)$  of the solution of the IVP (2.9), for all the estimates except those in (2.22) and (2.23), that require low values of  $k$ . For values of  $k > 5$ , we can give the following estimate, that demands additional regularity of the initial data in the statement of Theorem 1.2 in the superconformal case:

$$\begin{aligned} \|F_1\|_{L^1_{\Delta T} L^2_x} &\leq \|z^k\|_{L^1_{\Delta T} L^2_x} \leq \Delta T \|z\|_{L^\infty_{\Delta T} L^{2k}_x}^k \\ &\leq c \Delta T \|D^{\frac{k-1}{2}} z\|_{L^\infty_{\Delta T} L^2_x}^k \leq c \Delta T \|z\|_\gamma^k \\ &\leq c N^{-\frac{k-1}{2}(1-s)} N^{(\frac{k-1}{2}-s)k} \leq c N^{\frac{k-1}{2} - \frac{(k+1)}{2}s}, \end{aligned} \quad (2.36)$$

where  $s > \gamma = \frac{k-1}{k}$ .

With these estimates at hand, we obtain the same growth condition as in (2.28):

$$\|(\partial_t I, \nabla I)(\Delta T)\| + \|I(\Delta T)\|_{L^{k+1}_x}^{\frac{k+1}{2}} \leq c N^{\frac{(k-1)}{2} - \frac{(k+1)}{2}s + \varepsilon(2-(k-1)(1-s))}. \quad (2.37)$$

The closing argument that allows to yield the solution in the interval  $[0, T]$  for any  $T \gg 1$ , is based in an iterative process along with the growth condition (2.37).

In the interval  $[0, \Delta T]$ , we have that the solution of IVP (1.1) is given by

$$\begin{aligned} u(t) &= v(t) + z(t) \\ &= v(t) + I(t) + \partial_t W(t)u_{0,2} + W(t)u_{1,2}, \end{aligned} \quad (2.38)$$

where  $I(t) = -\int_0^t W(t-t')(F)(t')dt'$ , with  $F$  as in (2.11).

Next, we solve the IVP (1.1) in the interval  $[\Delta T, 2\Delta T]$  with initial data  $(v(\Delta T) + I(\Delta T), \partial_t v(\Delta T) + \partial_t I(\Delta T)) \in \dot{H}^1 \cap L^{k+1}_x \times L^2_x$ , which makes sense because the smoothing effect that takes place in the integral part  $I(t)$ . With this solution at hand, we consider IVP (2.9) with initial data

$$(\partial_t W(\Delta T)u_{0,2} + W(\Delta T)u_{1,2}, -\Delta W(\Delta T)u_{0,2} + \partial_t W(\Delta T)u_{1,2}). \quad (2.39)$$

This iterative procedure is continued until any given time  $T \gg 1$  is reached. In order to accomplish this goal we have to be sure that the process can be carried out uniformly, that is, that the growth condition in (2.8) should hold at all the steps up to reaching time  $T$ . It is here where estimate (2.37) turns out to be essential.

To reach time  $T$ , the number of iterations is

$$\frac{T}{\Delta T}. \quad (2.40)$$

For the problem with initial data in the energy space there is some growth of the initial data in the  $\|(\partial_t \cdot, \nabla \cdot)\| + \|\cdot\|_{L_x^{\frac{k+1}{2}}}$ -norm during each iteration, coming from the integral part  $I(t)$  and measured in (2.28). Since we want to keep (2.9) uniform, the following estimate should hold:

$$\frac{T}{\Delta T} N^{\frac{(k-1)}{2} - \frac{(k+1)}{2}s + \varepsilon(2 - (k-1)(1-s))} \leq cN^{1-s}; \quad (2.41)$$

and plugging into (2.39) the size of  $\Delta T$ , we have to check that

$$TN^{\frac{k-1}{2}(1-s)} N^{\frac{(k-1)}{2} - \frac{(k+1)}{2}s + \varepsilon(2 - (k-1)(1-s))} \leq cN^{1-s}. \quad (2.42)$$

Inequality (2.42) holds provided  $s > \frac{k-2}{k-1}$ ,  $\varepsilon$  is sufficiently small and  $N = N(T)$  is sufficiently large. We remark however, that (2.36) imposes an additional restriction on  $s$  for the superconformal case, namely,  $s > \frac{k-1}{k}$ . This completes the proof of Theorem 1.2.  $\square$

**Acknowledgments.** The author has been partially supported by the DIB, Universidad Nacional de Colombia, Bogotá.

## References

- [1] J. BOURGAIN, *Refinements of Strichartz's inequality and applications to 2D-NLS with critical nonlinearity*, IMRN **5** (1998), 253–283.
- [2] ———, *New global well-posedness results for the nonlinear Schrödinger equations*, AMS, 1999.
- [3] G. FONSECA, *Growth of the  $H^s$ -norm for the modified Korteweg-de Vries equation*, Diff. Integral Eq **13** no. 7-9 (2000), 1081–1093.
- [4] G. FONSECA, F. LINARES & G. PONCE, *Global well-posedness for the modified Korteweg-de Vries equation*, Comm. PDE. **24** (1999), 683–705.
- [5] M. KEEL & T. TAO, *Local and Global well-posedness of wave maps on  $\mathbb{R}^{1+1}$  for rough data*, IMRN **21** (1998), 1117–1156.
- [6] C. KENIG, G. PONCE & L. VEGA, *Global well-posedness for semi-linear wave equations*, Comm. PDE **25** (2000), 1741–1752.
- [7] H. LINBLAD, *A sharp counterexample to local existence of low regularity solutions to nonlinear wave equations*, Duke Math. J. **72** (1993), 503–539.
- [8] H. LINBLAD & C. SOGGE, *Global On existence and scattering with minimal regularity for semilinear wave equations*, J. Funct. Anal. **130** (1995), 357–426.
- [9] H. PECHER, *Nonlinear small data for the wave and Klein-Gordon equations*, Math Z **185** (1984), 261–270.
- [10] R. S. STRICHARTZ, *Restrictions of Fourier transforms to quadratic surfaces and decay of solutions of wave equations*, Duke Math. J. **44** (1977), 705–714.

(Recibido en noviembre de 2000)

DEPARTAMENTO DE MATEMÁTICAS  
UNIVERSIDAD NACIONAL DE COLOMBIA  
BOGOTÁ, COLOMBIA

e-mail: gfonseca@matematicas.unal.edu.co