

On n -th roots of meromorphic maps

Sobre raíces n -ésimas de funciones meromorfas

JUAN C. GARCÍA^{1,✉}, RUBÉN A. HIDALGO^{2,a}

¹Universidad Central del Ecuador, Quito, Ecuador

²Universidad de La Frontera, Temuco, Chile

ABSTRACT. Let S be a connected Riemann surface and let $\varphi : S \rightarrow \widehat{\mathbb{C}}$ be branched covering map of finite type. If $n \geq 2$, then we describe a simple geometrical necessary and sufficient condition for the existence of some n -th root, that is, a meromorphic map $\psi : S \rightarrow \widehat{\mathbb{C}}$ such that $\varphi = \psi^n$.

Key words and phrases. Riemann surfaces, holomorphic branched coverings, maps.

2010 Mathematics Subject Classification. 30F10, 57M12.

RESUMEN. Sea S una superficie de Riemann conexa y $\varphi : S \rightarrow \widehat{\mathbb{C}}$ un cubrimiento ramificado holomorfo de tipo finito. Para cada $n \geq 2$ describimos una condición geométrica necesaria y suficiente para la existencia de alguna raíz n -ésima, esto es, una función meromorfa $\psi : S \rightarrow \widehat{\mathbb{C}}$ de manera que $\varphi = \psi^n$.

Palabras y frases clave. Superficies de Riemann, cubrimientos ramificados holomorfos, mapas.

1. Introduction

In this paper, S will denote a connected (not necessarily compact or of finite type) Riemann surface and $\widehat{\mathbb{C}} = \mathbb{C} \cup \{\infty\}$ will be the Riemann sphere. A holomorphic surjective map $\varphi : S \rightarrow \widehat{\mathbb{C}}$ is a holomorphic branched covering if: (i) it has a finite set $B_\varphi \subset \widehat{\mathbb{C}}$ of branching points, (ii) $\varphi : S \setminus \varphi^{-1}(B_\varphi) \rightarrow \widehat{\mathbb{C}} \setminus B_\varphi$ is a holomorphic covering map and (iii) around each point $q \in B_\varphi$ there is a open disc Δ_q such that $\varphi^{-1}(\Delta_q)$ consists of a collection of pairwise disjoint discs V_j such that each of the restrictions $\varphi : V_j \rightarrow \Delta_q$ is a finite degree $d_{q,j}$ holomorphic map (i.e., there are biholomorphisms $z : V_j \rightarrow \mathbb{D}$ and $w : \Delta_q \rightarrow \mathbb{D}$, where \mathbb{D} is the unit disc, such that $w \circ \varphi \circ z^{-1}(z) = z^{d_{q,j}}$). For each $q \in B_\varphi$,

let $M_q \subset \{1, 2, \dots\}$ be the set of local degrees of φ at the points in the fiber $\varphi^{-1}(p)$. We say that φ is of *finite type* if the sets M_q are finite. This condition permits to define the *branch order* of $q \in B_\varphi$ as the minimum common multiple of the values in M_q .

Let $\varphi : S \rightarrow \widehat{\mathbb{C}}$ a holomorphic branched covering of finite type. If $n \geq 2$, then a meromorphic map $\psi : S \rightarrow \widehat{\mathbb{C}}$ such that $\varphi = \psi^n$ is called an n -th root of φ (the others n -th roots of φ are of the form $e^{2k\pi i/n}\psi$, where $k = 0, 1, \dots, n-1$).

The existence of an n -th root of φ necessarily implies that: (a) $\infty, 0 \in B_\varphi$ and (b) the branch orders of both 0 and ∞ are multiples of n . These two conditions are not sufficient for φ to have an n -root. For $S = \widehat{\mathbb{C}}$ the existence of an n -th root is equivalent for each zero and each pole of the rational map φ to have degree a multiple of n (which in particular asserts conditions (a) and (b)). But, for other Riemann surfaces, the above is not sufficient in general.

In [3] there was provided a simple geometrical necessary and sufficient condition for φ to have a 2-th root. In this paper, we generalize such a description for every $n \geq 2$ (Theorem 2.3).

In the final section we generalize some of the tools in the proof of the main result to the context of Kleinian groups of higher dimension.

Remark 1.1 (A connection to Fuchsian groups). Let K be a finitely generated Fuchsian group, acting on the hyperbolic plane \mathbb{H}^2 , such that \mathbb{H}^2/K is an orbifold of genus zero (so its underlying Riemann surface structure is $\widehat{\mathbb{C}}$) and let k_1, \dots, k_r be the orders of its cone points. Let Γ be a subgroup of K and let S be the underlying Riemann surface structure associated to the hyperbolic orbifold \mathbb{H}^2/Γ (if Γ is assumed to be torsion free, then this orbifold has no cone points). It is well known that S is of finite type if and only if Γ is finitely generated (in particular, by taking infinitely generated subgroups we obtain examples of surfaces of infinite type). The inclusion $\Gamma \leq K$ induces a holomorphic branched covering of finite type $S = \mathbb{H}^2/\Gamma \rightarrow \widehat{\mathbb{C}}$. The local degree of φ at each point of S is either 1 (the generic case) or a divisor of some k_j (if Γ is torsion free, then the local degree at each point over a cone point of order k_j is also k_j). Conversely, the uniformization theorem asserts that, for a connected Riemann surface S of hyperbolic type, each branched covering of finite type $\varphi : S \rightarrow \widehat{\mathbb{C}}$ is obtained in such a way for suitable choices of Γ and K .

2. Main results

Before stating our main result we need some definitions.

2.1. Admissible arcs and n -Z-orientability

Let us recall that a *map* on an orientable and connected surface X is a 2-cell decomposition of it, induced by the embedding of a connected graph \mathcal{H} for

which each of its vertices has a finite degree and each face (i.e., the connected components of $X \setminus \mathcal{H}$) are finite-sided polygons.

An *admissible arc* for a branched covering of finite type $\varphi : S \rightarrow \widehat{\mathbb{C}}$, with $0, \infty \in B_\varphi$, is a simple arc $\delta \subset \widehat{\mathbb{C}}$ whose end points are 0 and ∞ , and $B_\varphi \setminus \{0, \infty\} \subset \delta$. For such an admissible arc, the graph $\varphi^{-1}(\delta) = \mathcal{G}_\delta \subset S$ defines a map \mathcal{F}_δ on S ; each of its faces is a polygon with $2(r-1)$ sides, $r = \#B_\varphi$. We say that \mathcal{F}_δ is *n -Z-orientable* if we may label its faces with numbers inside $\{1, 2, \dots, n\}$ (see Figure 1 at the end for the case $n = 3$), such that the following two properties hold:

- (1) around each vertex $q \in \varphi^{-1}(0)$ (respectively, $q \in \varphi^{-1}(\infty)$), following the counterclockwise (respectively, the clockwise) orientation, the labelling is a finite consecutive sequence of the ordered tuple $(1, 2, \dots, n)$ (these correspond, respectively, to the first two figures at the left of Figure 1),
- (2) around each vertex $q \notin \varphi^{-1}\{0, \infty\}$, following the counterclockwise orientation we see a finite sequence of a same tuple $(i, i+1)$, if $i = 1, \dots, n-1$, or of the tuple $(1, n)$ (i.e., we see as labelling $i, i+1, i, i+1, \dots, i, i+1$, or we see $1, n, 1, n, \dots, 1, n$) (these correspond to the last three figures at the left of Figure 1).

Remark 2.1. For instance, if we let $\varphi : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ to be $\varphi(z) = z^3$, then it can be seen that the positive real line is an admissible arc. As in this case \mathcal{F}_δ has exactly three faces (cyclicly permuted by the rotation $A(z) = e^{2\pi i/3}z$), this cannot be n -Z-orientable for $n \neq 3$.

The 2-Z-orientable definition was introduced by Zapponi in [9, 10, 11] (he used the term “orientable”) in order to decide if a given Strebel quadratic meromorphic form \mathcal{Q} [8] on a closed Riemann surface has an square root (the 2-cell decomposition is obtained from the graph whose vertices are the zeroes of \mathcal{Q} and the edges are its non-compact horizontal trajectories).

Remark 2.2. If δ_1 and δ_2 are admissible arcs for φ , then there is an orientation-preserving homeomorphism $h : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$ fixing the points 0 and ∞ and such that $h(\delta_1) = \delta_2$.

2.2. n -Z-orientability is a necessary and sufficient condition

The following generalizes the results in [3] done for the case $n = 2$.

Theorem 2.3. *Let S be a connected Riemann surface. Let $\varphi : S \rightarrow \widehat{\mathbb{C}}$ be a holomorphic branched covering map of finite type with $0, \infty \in B_\varphi$ and each one with branch order a multiple of $n \geq 2$. If δ is an admissible arc for φ , then the existence of n -th roots of φ is equivalent for the map \mathcal{F}_δ to be n -Z-orientable.*

In terms of Fuchsian groups, Theorem 2.3 asserts the following.

Corollary 2.4. *Let $K < \mathrm{PSL}_2(\mathbb{R})$ be a co-compact Fuchsian group acting on the hyperbolic plane \mathbb{H}^2 such that \mathbb{H}^2/K has genus zero. Let $n \geq 2$ be an integer and assume that there are two cone points $p, q \in \mathbb{H}^2/K$ whose cone orders are multiples of n . Let $\delta \subset \mathbb{H}^2/K$ be a simple arc whose end points are p and q and containing all other cone points in its interior. Let \mathcal{F}_δ be the map of \mathbb{H}^2 induced by the lifting of δ to \mathbb{H}^2 . Then the existence of a normal subgroup Γ_n , of index $n \geq 2$, in K such that $K/\Gamma_n \cong \mathbb{Z}_n$ and \mathbb{H}^2/Γ_n has genus zero, is equivalent for \mathcal{F}_δ to be n -Z-orientable.*

In order to see the above, we identify \mathbb{H}^2/K with $\widehat{\mathbb{C}}$ and $p = 0, q = \infty$. Then in Theorem 2.3 we set $S = \mathbb{H}^2$ and take $\varphi : \mathbb{H}^2 \rightarrow \widehat{\mathbb{C}}$ a branched regular covering with K as its deck group.

2.3. Case of compact Riemann surfaces

If S is a compact Riemann surface, then every non-constant meromorphic map $\varphi : S \rightarrow \widehat{\mathbb{C}}$ is a holomorphic branched covering of finite type. In this way, Theorem 2.3 can be rewritten as follows.

Corollary 2.5. *Let S be a compact Riemann surface. Let $\varphi : S \rightarrow \widehat{\mathbb{C}}$ be a non-constant meromorphic map with $0, \infty \in B_\varphi$ and each one with branch order a multiple of $n \geq 2$. If δ is an admissible arc for φ , then the existence of n -th roots of φ is equivalent for the associated map \mathcal{F}_δ to be n -Z-orientable.*

The compact Riemann surface S can be defined by a complex projective algebraic curve inside \mathbb{P}^n and the meromorphic map $\varphi : S \rightarrow \widehat{\mathbb{C}}$ can be described by a rational map.

Let us assume S is defined as the zero locus of the homogeneous polynomials $P_1, \dots, P_r \in \mathbb{C}[x_1, \dots, x_{n+1}]$ and that φ corresponds to the quotient Q_1/Q_2 , where $Q_1, Q_2 \in \mathbb{C}[x_1, \dots, x_{n+1}]$ are homogeneous polynomials of the same degree. If $\sigma \in \mathrm{Gal}(\mathbb{C})$, the group of field automorphisms of \mathbb{C} , then we set S^σ (respectively, φ^σ) the projective algebraic curve defined by the polynomials $P_1^\sigma, \dots, P_r^\sigma$ (respectively, Q_1^σ/Q_2^σ), where P_j^σ (respectively, Q_j^σ) is obtained from P_j (respectively, Q_j) by applying σ to all of its coefficients. It can be checked that S^σ is again a compact Riemann surface and that $\varphi^\sigma : S^\sigma \rightarrow \widehat{\mathbb{C}}$ is a holomorphic branched covering map of finite type.

If (S_1, φ_1) and (S_2, φ_2) are isomorphic (i.e., there is an isomorphism $\psi : S_1 \rightarrow S_2$ such that $\varphi_1 = \varphi_2 \circ \psi$), then for every $\sigma \in \mathrm{Gal}(\mathbb{C})$ it holds that the two new pairs $(S_1^\sigma, \varphi_1^\sigma)$ and $(S_2^\sigma, \varphi_2^\sigma)$ are still isomorphic (by ψ^σ). This process provides of an action of $\mathrm{Gal}(\mathbb{C})$ on (equivalence classes) of pairs (S, φ) . As the property of having an n -th root is a $\mathrm{Gal}(\mathbb{C})$ -invariant, the above asserts the following.

Corollary 2.6. *Let S be a compact Riemann surface. Let $\varphi : S \rightarrow \widehat{\mathbb{C}}$ be a non-constant meromorphic map with $0, \infty \in B_\varphi$ and each one with branch order*

a multiple of $n \geq 2$. Then the n -Z-orientability property of a pair (S, φ) is a $\text{Gal}(\mathbb{C})$ -invariant.

Belyi's theorem [2] asserts that a compact Riemann surface S can be defined by a curve over the field $\overline{\mathbb{Q}}$ of algebraic numbers if and only if there is a non-constant meromorphic map, called a Belyi map, $\beta : S \rightarrow \widehat{\mathbb{C}}$ whose branching points are contained inside $\{\infty, 0, 1\}$. On S there is a 2-cell decomposition \mathcal{D}_β , called a dessin d'enfant [6], whose underlying graph $\beta^{-1}([0, 1])$ is bipartite (the black vertices are $\beta^{-1}(1)$ and the white ones are $\beta^{-1}(0)$). Corollary 2.5 provides a geometrical condition for the new Belyi map $\varphi = \beta/(\beta - 1)$ to have n -square roots. Such a geometrical condition, in terms of the dessin \mathcal{D}_β , is that its faces can be labeled using numbers in $\{1, 2, \dots, n\}$ such that around each black vertex (respectively, white vertex), following the counterclockwise orientation, we see a finite consecutive sequence of the tuple $(1, 2, \dots, n)$ (respectively, $(n, n - 1, \dots, 2, 1)$). This condition provides a Galois invariant for the dessin \mathcal{D}_β . In [4], for $n = 2$, it was observed that this is a new Galois invariant on dessins d'enfants. We expect (but we have no explicit evidence) that for each $n \geq 3$ it provides a new Galois invariant.

3. Proof of Theorem 2.3

3.1.

Let $r \geq 2$ be the cardinality of B_φ and let $\delta \subset \widehat{\mathbb{C}}$ an admissible arc for φ , starting at the branch point $p_1 = 0$, ending at the branch point $p_r = \infty$. We label the rest of the branch points of φ as p_2, \dots, p_{r-1} , such that p_j is between p_{j-1} and p_{j+1} . Let us denote by k_j be the branch order of p_j . We are assuming that k_1 and k_r are both multiples of $n \geq 2$. Set \mathbb{X} to be either $\widehat{\mathbb{C}}$, \mathbb{C} or \mathbb{H}^2 depending on if

$$\sum_{j=1}^r (1 - k_j^{-1}) - 2$$

is negative, zero or positive, respectively.

3.2.

Let K be a discrete group of isometries of \mathbb{X} such that $\mathbb{X}/K = \widehat{\mathbb{C}}$ and whose cone points are p_1, \dots, p_r , with respective cone orders k_1, \dots, k_r . Let $\pi_K : \mathbb{X} \rightarrow \widehat{\mathbb{C}}$ be a regular holomorphic branched covering with K as deck group.

The arc δ defines a fundamental domain P_δ for K (see Figure 2 at the end), with $2(r - 1)$ sides, and set of side pairings $\mathcal{A}_{P_\delta} = \{C_1, \dots, C_{r-1}\}$, such that

$$K = \langle C_1, \dots, C_{r-1} : C_1^{k_1} = (C_1^{-1}C_2)^{k_2} = \dots = (C_{r-2}^{-1}C_{r-1})^{k_{r-1}} = C_{r-1}^{k_r} = 1 \rangle.$$

The K -translates of P_δ produces a 2-cell decomposition $\mathcal{T}_{K, P_\delta}$ of \mathbb{X} , that is, a map on \mathbb{X} . As k_1 and k_r are multiples of n , the following produces a surjective

homomorphism

$$\theta_0 : K \rightarrow G = \langle \sigma \rangle \cong \mathbb{Z}/n\mathbb{Z} : C_j \mapsto \sigma, \quad j = 1, \dots, r-1.$$

As $\ker(\theta_0)$ is the group generated by the K -conjugates of the elements

$$C_1^n, C_1^{-1}C_2, C_1C_2^{-1}, \dots, C_1^{-1}C_{r-1}, C_1C_{r-1}^{-1},$$

it follows that adjacent faces of the map $\mathcal{T}_{K, P_\delta}$ have different labels.

Let $x_1, \dots, x_r \in \mathbb{X}$ be the fixed points of the elements $C_1, C_1^{-1}C_2, C_2^{-1}C_3, \dots, C_{r-2}^{-1}C_{r-1}$ and C_{r-1} , respectively. Then, $\pi_K(x_j) = p_j$, for $j = 1, \dots, r$.

Remark 3.1. The map $\mathcal{T}_{K, P_\delta}$ is n -Z-orientable. To see this, for each $T \in K$, we label the T -translated of P_δ by the element $\theta_0(T) \in G$. Now, in order to be consequent with our definition of n -Z-orientability as in the introduction, we make the identification of σ^j with the integer $j+1$, for $j = 0, 1, \dots, n-1$.

The orbifold $\mathbb{X}/\ker(\theta_0)$ can be identified with the Riemann sphere $\widehat{\mathbb{C}}$ with exactly $n(r-2) + 2$ cone points, these being of orders $k_1/n, k_2/n, \dots, k_{r-1}/n$. The pair $(K, \ker(\theta_0))$ induces a Möbius transformation A , of order n , and (by using the above identification) a degree n meromorphic map $\eta : \widehat{\mathbb{C}} \rightarrow \widehat{\mathbb{C}}$, whose deck group is $\langle A \rangle \cong \mathbb{Z}/n\mathbb{Z}$, branched at the end points of δ , i.e., 0 and ∞ . Up to conjugation by a suitable Möbius transformation, we may assume that $A(z) = e^{2\pi i/n}z$ and $\eta(z) = z^n$.

The set $\eta^{-1}(\delta)$ is a collection of n simple arcs (containing all the cone points of $\mathbb{X}/\ker(\theta_0)$ and whose end points are the two fixed points of A , i.e., 0 and ∞) which are cyclically permuted by A . This provides a n -Z-orientable map \mathcal{F}_0 on $\widehat{\mathbb{C}}$ and it is induced by the map $\mathcal{T}_{K, P_\delta}$.

Remark 3.2. Each index normal subgroup Γ_0 of K , such that $K/\Gamma_0 \cong \mathbb{Z}/n\mathbb{Z}$, is given as the kernel of a surjective homomorphism $\theta : K \rightarrow \mathbb{Z}/n\mathbb{Z}$. Let us assume that \mathbb{X}/Γ_0 has genus zero. In this case, the inclusion $\Gamma_0 \triangleleft K$ induces a regular holomorphic branched covering $\pi_{\Gamma_0} : \mathbb{X} \rightarrow \widehat{\mathbb{C}}$ with Γ_0 as its deck group, such that $\pi_K = R \circ \pi_{\Gamma_0}$, where $R(z) = z^n$. Let us restrict to those Γ_0 such that $\pi_{\Gamma_0}(x_1) = 0$ and $\pi_{\Gamma_0}(x_r) = \infty$. (Note that if each of the k_j , where $j = 2, \dots, r-1$, are not a multiple of n , then this is the only possibility). In this case, for $j = 2, \dots, r-1$, $\pi_{\Gamma_0}(x_j)$ is a cone point of order k_j (since that point is not critical point of R). This asserts that $C_{j-1}^{-1}C_j \in \Gamma_0$. If we set $\sigma = \theta(C_1)$, which is a generator of $\mathbb{Z}/n\mathbb{Z}$, the previous asserts that $\theta(C_j) = \sigma$, for every $j = 1, \dots, r-1$. In other words, up to post-composing by an automorphism of $\mathbb{Z}/n\mathbb{Z}$, we obtain θ_0 , i.e., Γ_0 is uniquely determined.

3.3.

As a consequence of the uniformization theorem, there is a proper subgroup Γ of K such that S is the Riemann surface structure of the orbifold \mathbb{X}/Γ , a

regular holomorphic (possible branched) covering $\pi_\Gamma : \mathbb{X} \rightarrow S$, with deck group Γ , such that $\pi_K = \varphi \circ \pi_\Gamma$ (i.e., φ is induced by the inclusion $\Gamma < K$).

Let us observe that the π_Γ -image of the n -Z-orientable map \mathcal{T}_{K,P_δ} on \mathbb{X} is the map \mathcal{F}_δ (which might or not be n -Z-orientable in principle).

Now, the existence of a meromorphic map $\psi : S \rightarrow \widehat{\mathbb{C}}$, such that $\varphi = \eta \circ \psi = \psi^n$ is equivalent to have that $\Gamma \leq \ker(\theta_0)$ (see Remark 3.2). By lemma 3.3 (whose arguments follow the same ideas as in [3] for $n = 2$), the previous is equivalent for the n -Z-orientability of \mathcal{F}_δ . This will provide the desired result.

Lemma 3.3. *The map \mathcal{F}_δ is n -Z-orientable if and only if $\Gamma \leq \ker(\theta_0)$.*

Proof. Let us assume $\Gamma \leq \ker(\theta_0)$. The idea is to push-down the (n -Z-orientable) labelling on the faces of \mathcal{T}_{K,P_δ} to the faces of \mathcal{F}_δ . Two faces F_1 and F_2 of \mathcal{T}_{K,P_δ} are projected to the same face if and only if there is some $T \in \Gamma$ such that $F_2 = T(F_1)$. As we are assuming $\Gamma \leq \ker(\theta_0)$, the induced labelling is well defined. It is not difficult to observe that induced labelling on the map \mathcal{F}_δ satisfies the condition for being n -Z-orientable.

In the other direction, let us assume we have a labelling for \mathcal{F}_δ satisfying the n -Z-orientability. By the connectivity of S , we may construct a fundamental (connected) domain \mathcal{Q} for Γ by gluing some copies K -translated of P_δ (as many as the index of Γ in K). The projection of those copies of P_δ , used in the construction of \mathcal{Q} , projects under π_Γ exactly to the faces of \mathcal{F}_δ . Now, lift the labelling of the n -Z-orientable map \mathcal{F}_δ to obtain labelling of these copies of P_δ included in \mathcal{Q} . Use the group K to translate these labels to the rest of K -translates of P_δ . This provides a labelling on \mathcal{T}_{K,P_δ} satisfying the n -Z-orientability property. By Remark 3.2, this can be assumed to be the labelling provided by θ_0 . As the above procedure of pulling-down the labelling from \mathcal{T}_{K,P_δ} to \mathcal{F}_δ induces the given labelling, it follows from the first part that $\Gamma \leq \ker(\theta_0)$. \checkmark

4. A remark: θ -Zapponi-orientability of Kleinian groups

In the previous section we have considered a Fuchsian group K , a fundamental polygon P , the set $\mathcal{A}_P \subset K$ of its side pairings, and a surjective homomorphism $\theta_0 : K \rightarrow G = \mathbb{Z}/n\mathbb{Z}$ such that $\ker(\theta_0) \cap \mathcal{A}_P = \emptyset$. The homomorphism θ_0 permitted to label each of the faces of the map $\mathcal{T}_{K,P}$, using as labelling the elements of G , and such that adjacent faces have different labels. This procedure can be generalized for any Kleinian group as follows.

Let K be a discrete group of isometries of \mathbb{X}^m , where \mathbb{X}^m is either the m -dimensional hyperbolic \mathbb{H}^m or the m -dimensional Euclidian space \mathbb{E}^m or the m -dimensional sphere \mathbb{S}^m . Let $P \subset \mathbb{X}^m$ be a fundamental polyhedron of K and let \mathcal{A}_P the subset of K consisting of the side-pairings of P . It is well known that \mathcal{A}_P is a set of generators for K and that a complete set of relations is provided by how the sides of P are glued by these side-pairings (Poincaré Polyhedron

Theorem, see [1, 5, 7]). The K -translates of P provides a n -tessellation $\mathcal{T}_{K,P}$ of \mathbb{X}^m .

4.1. (K, P) -admissible homomorphisms

Let $\theta : K \rightarrow G$, where G is a finite group, be a surjective homomorphism. For each $T \in K$ we proceed to label the n -face $T(P)$ using the element $\theta(T) \in G$. If adjacent faces have different labels, then we say that θ is (K, P) -admissible.

Lemma 4.1. θ is (K, P) -admissible if and only if $\mathcal{A}_P \cap \ker(\theta) = \emptyset$.

Proof. This follows from the fact that, for $T_1, T_2 \in K$, one has that $T_1(P)$ and $T_2(P)$ are adjacent if and only if there is some $L \in \mathcal{A}_P$ such that $T_2 = T_1 L$. \checkmark

Remark 4.2. For every surjective homomorphism $\theta : K \rightarrow G$, it is possible to find a fundamental polyhedron P for K such that θ is (K, P) -admissible.

Example 4.3. (1) If K_P is the subgroup of K generated by all the elements of the form AB , where $A, B \in \mathcal{A}_P$, then either $K_P = K$ or has index two in K . If $\theta : K \rightarrow G = \mathbb{Z}_2$ is any homomorphism, then $K_P \leq \ker(\theta)$. It follows that θ is (K, P) -admissible if and only if $K \neq K_P = \ker(\theta)$ (in particular, there is at most one (K, P) -admissible homomorphism onto \mathbb{Z}_2). (2) Let $n, r \geq 2$ and let us consider a Fuchsian group, acting in the hyperbolic plane \mathbb{H}^2 , with the following presentation

$$K = \langle C_1, \dots, C_{r-1} : C_1^{k_1} = (C_1^{-1}C_2)^{k_2} = \dots = (C_{r-2}^{-1}C_{r-1})^{k_{r-1}} = C_{r-1}^{k_r} = 1 \rangle.$$

Let P be a fundamental domain of K as shown in Figure 2. Its set of side-pairings is $\mathcal{A}_P = \{C_1, \dots, C_{r-1}\}$. Let $G = \langle \sigma \rangle \cong \mathbb{Z}/n\mathbb{Z}$. If k_1 and k_r are both multiples of n , then we may consider the surjective homomorphism $\theta_0 : K \rightarrow G$, defined by $\theta_0(C_j) = \sigma$, for every $j = 1, \dots, r-1$. As $\ker(\theta_0)$ is the group generated by the conjugates of the elements $C_1^n, C_1^{-1}C_2, C_1C_2^{-1}, \dots, C_1^{-1}C_{r-1}, C_1C_{r-1}^{-1}$, it follows that θ_0 is (K, P) -admissible. The induced labelling of $\mathcal{T}_{K,P}$ by θ_0 satisfies to be n - \mathbb{Z} -orientable.

4.2. θ - \mathbb{Z} -orientable subgroups

Let $\theta : K \rightarrow G$ be a (K, P) -admissible homomorphism. If Γ is a proper subgroup of K , then the tessellation $\mathcal{T}_{K,P}$ induces an m -dimensional tessellation $\mathcal{T}_{K,P,\Gamma}$ on the geometric orbifold $\mathcal{O}_\Gamma = \mathbb{X}^m/\Gamma$. The labelling on the faces of $\mathcal{T}_{K,P}$, provided by the (K, P) -admissible homomorphism θ , induces a labelling of the faces of the tessellation $\mathcal{T}_{K,P,\Gamma}$. It is not difficult to see that the adjacent m -faces of this last tessellation have different labels if and only if $\Gamma \leq \ker(\theta)$. If this is the situation, we say that (K, P, Γ) is θ - \mathbb{Z} -orientable.

We summarize all the above in the following.

Lemma 4.4. *Let K be a discrete group of isometries of \mathbb{X}^m , $P \subset \mathbb{X}^m$ be a fundamental polyhedron for it and $\mathcal{A}_P \subset K$ be the set of side-pairings of P . Let $\theta : K \rightarrow G$ be a (K, P) -admissible homomorphism onto a finite group G (equivalently, $\ker(\theta) \cap \mathcal{A}_P = \emptyset$). Then (K, P, Γ) is θ -Z-orientable if and only if $\Gamma \leq \ker(\theta)$.*





FIGURE 1. The labelling for 3-Z-orientability

FIGURE 2. The fundamental polygon P_δ

Acknowledgment. We thank the anonymous referee for her/his comments and suggestions which contributed to improving this manuscript.

References

- [1] B. N. Apanasov, *Conformal Geometry of Discrete Groups and Manifolds*, De Gruyter Expositions in Mathematics **32**, 2000.
- [2] G. V. Belyi, *On Galois extensions of a maximal cyclotomic field*, Mathematics of the USSR-Izvestiya **14** (1980), no. 2, 247–256.
- [3] J. C. García and R. A. Hidalgo, *On square roots of meromorphic maps*, Results in Mathematics **74** (2019), no. 3, Art. 118, 12 pp., <https://doi.org/10.1007/s00025-019-1042-7>.
- [4] E. Girono, G. González-Diez, R. A. Hidalgo, and G. A. Jones, *Zapponi-orientable dessins d'enfants*, Revista Matemática Iberoamericana **36** (2020), no. 2, 549–570.
- [5] L. Greenberg, *Fundamental polyhedra for kleinian groups*, Annals of Mathematics, Second Series **84** (1966), 433–441.

- [6] A. Grothendieck, *Esquisse d'un Programme. (1984). In Geometric Galois Actions, 1. L. Schneps and P. Lochak eds.*, London Math. Soc. Lect. Notes Ser **242** (1997), 5–47, Cambridge University Press, Cambridge.
- [7] J. G. Ratcliffe, *Foundations of hyperbolic manifolds*, Berlin, New York: Springer-Verlag, 1994.
- [8] K. Strebel, *Quadratic differentials*, Springer, New-York, 1984.
- [9] L. Zapponi, *Grafì, differenziali di strebel e curve. Tesi di laurea*, Università degli studi di Roma “La Sapienza”, 1995.
- [10] ———, *Dessins d'enfants en genre 1. in Geometric Galois Actions, 2. L. Schneps and P. Lochak eds.*, London Math. Soc. Lecture Note Ser. (1997), 79–116, Cambridge Univ. Press, Cambridge.
- [11] ———, *Fleurs, arbres et cellules: un invariant galoisien pour une famille d'arbres*, Compositio Mathematica **122** (2000), 113–133.

(Recibido en diciembre de 2019. Aceptado en abril de 2020)

FACULTAD DE CIENCIAS
UNIVERSIDAD CENTRAL DEL ECUADOR
QUITO, ECUADOR
e-mail: jcg70@gmail.com

DEPARTAMENTO DE MATEMÁTICA Y ESTADÍSTICA
UNIVERSIDAD DE LA FRONTERA
TEMUCO, CHILE
e-mail: ruben.hidalgo@ufrontera.cl