

Exact solutions for a new fifth-order integrable system

CÉSAR A. GÓMEZ S.

Universidad Nacional de Colombia, Bogotá

ABSTRACT. We consider a new fifth-order integrable system. The exact solutions are obtained by the generalized tanh method.

Keywords and phrases. Nonlinear differential equation; travelling wave solution; generalized tanh method.

2000 Mathematics Subject Classification. Primary: 35Q53. Secondary: 35Q51.

RESUMEN. Nosotros consideramos un nuevo sistema integrable de quinto orden. Obtenemos las soluciones exactas de dicho sistema por el método tanh generalizado.

1. Introduction

The knowledge of closed form solutions of nonlinear PDEs facilitates the testing of numerical solvers, and aids in the stability analysis of solutions. There are a number of methods for the construction of exact solutions to differential partial differential equations of the mathematical physics [1], [2], [3], [4] [5], [6],[6], [9], [10]. Travelling wave solutions of many nonlinear PDEs, from solitons theory can be expressed as polynomials of the hyperbolic tangent and secant functions. In this brief note, we present exact solutions for the new fifth-order integrable system using the generalized tanh method.

2. Mikhailov–Novikov–Wang system

Mikhailov–Novikov–Wang system [8], [12] reads

$$\begin{cases} u_t = u_{xxxxx} - 20uu_{xxx} - 50u_xu_{xx} + 80u^2u_x + w_x \\ w_t = -6wu_{xxx} - 2u_{xx}w_x + 96wuu_x + 16w_xu^2. \end{cases} \quad (2.1)$$

We search for exact solutions of system (2.1) in the form

$$\begin{cases} u(x, t) = v(\xi) \\ w(x, t) = w(\xi) \\ \xi = x + \lambda t. \end{cases} \quad (2.2)$$

As a result system (2.1) reduces to the nonlinear system ordinary differential equation

$$\begin{cases} \lambda v'(\xi) - v^{(5)}(\xi) + 20v(\xi)v^{(3)}(\xi) + 50v'(\xi)v''(\xi) - 80v^2(\xi)v'(\xi) - w'(\xi) = 0 \\ \lambda w'(\xi) + 6w(\xi)v'''(\xi) + 2v''(\xi)w'(\xi) - 96w(\xi)v(\xi)v'(\xi) - 16w'(\xi)v^2(\xi) = 0. \end{cases} \quad (2.3)$$

We obtain

$$w'(\xi) = \lambda v'(\xi) - v^{(5)}(\xi) + 20v(\xi)v^{(3)}(\xi) + 50v'(\xi)v''(\xi) - 80v^2(\xi)v'(\xi). \quad (2.4)$$

The first equation of (2.3) can be written as follows:

$$\left(\lambda v(\xi) - v^{(4)}(\xi) + 20v(\xi)v''(\xi) + 15 \left((v'(\xi))^2 \right) - \frac{80}{3}v^3(\xi) - w(\xi) \right)' = 0. \quad (2.5)$$

Integrating (2.5) once with respect to ξ we get

$$\lambda v(\xi) - v^{(4)}(\xi) + 20v(\xi)v''(\xi) + 15 (v'(\xi))^2 - \frac{80}{3}v^3(\xi) - w(\xi) = c, \quad (2.6)$$

where c is integration constant. As we look for the exact solutions of special form, we set $c = 0$ for simplicity. Therefore

$$w(\xi) = \lambda v(\xi) - v^{(4)}(\xi) + 20v(\xi)v''(\xi) + 15 (v'(\xi))^2 - \frac{80}{3}v^3(\xi). \quad (2.7)$$

Substituting (2.4) and (2.7) into second equation of (2.3), we obtain after simplifications the following ordinary differential equation:

$$\begin{aligned} & \lambda^2 v'(\xi) - \lambda v^{(5)}(\xi) + 26\lambda v(\xi)v'''(\xi) + 52\lambda v'(\xi)v''(\xi) - 192\lambda v^2(\xi)v'(\xi) \\ & - 2v''(\xi)v^{(5)}(\xi) + 160v(\xi)v''(\xi)v'''(\xi) + 100v'(\xi)(v''(\xi))^2 \\ & - 2880v^2(\xi)v'(\xi)v''(\xi) + 16v^2(\xi)v^{(5)}(\xi) - 480v^3(\xi)v'''(\xi) + 3840v^4(\xi)v'(\xi) \\ & - 6v'''(\xi)v^{(4)}(\xi) + 90(v'(\xi))^2v'''(\xi) + 96v(\xi)v'(\xi)v^{(4)}(\xi) - 1440v(\xi)(v'(\xi))^3 \\ & = 0. \end{aligned} \quad (2.8)$$

To obtain exact solution for equation (2.8), we use the generalized tanh method [10] which consist of the following five steps:

step 1

We consider solutions of (2.8) in the form

$$v(\xi) = \sum_{k=0}^m a_k \phi^k(\xi), \quad (2.9)$$

where $\phi(\xi)$ satisfies the Riccati equation

$$\phi'(\xi) = \phi^2(\xi) + k \quad (2.10)$$

whose solutions are given by

$$\phi(\xi) = \begin{cases} -\frac{1}{\xi}, & k = 0 \\ \sqrt{k} \tan(\sqrt{k}\xi) & k > 0 \\ -\sqrt{k} \cot(\sqrt{k}\xi) & k > 0 \\ -\sqrt{-k} \tanh(\sqrt{-k}\xi) & k < 0 \\ -\sqrt{-k} \coth(\sqrt{-k}\xi) & k < 0. \end{cases} \quad (2.11)$$

The exponent m must be determined by balancing the linear term of the highest order with the nonlinear term in (2.8). We obtain $m = 2$.

step 2

Substituting (2.9) into (2.8) and using (2.10) we obtain a polynomial in $\phi(\xi)$. Equating the coefficients of this polynomial to zero, we get the following system of algebraic equations:

- (1) $-16k^3\lambda a_1 + k\lambda^2 a_1 + 52k^2\lambda a_0 a_1 + 256k^3 a_0^2 a_1 - 192k\lambda a_0^2 a_1 - 960k^2 a_0^3 a_1 + 3840k a_0^4 a_1 + 180k^4 a_1^3 - 1440k^3 a_0 a_1^3 - 256k^5 a_1 a_2 + 104k^3\lambda a_1 a_2 + 2176k^4 a_0 a_1 a_2 - 5760k^3 a_0^2 a_1 a_2 + 400k^5 a_1 a_2^2 = 0.$
- (2) $-136k^2\lambda a_1 + \lambda^2 a_1 + 208k\lambda a_0 a_1 + 2176k^2 a_0^2 a_1 - 192\lambda a_0^2 a_1 - 3840k a_0^3 a_1 + 3840 a_0^4 a_1 + 3912k^3 a_1^3 - 192k\lambda a_1^3 - 18720k^2 a_0 a_1^3 + 23040k a_0^2 a_1^3 - 5824k^4 a_1 a_2 + 1196k^2\lambda a_1 a_2 + 37120k^3 a_0 a_1 a_2 - 1152k\lambda a_0 a_1 a_2 - 66240k^2 a_0^2 a_1 a_2 + 46080k a_0^3 a_1 a_2 - 15840k^3 a_1^3 a_2 + 22048k^4 a_1 a_2^2 - 51840k^3 a_0 a_1 a_2^2 = 0.$
- (3) $-256k^4 a_1^2 + 156k^2\lambda a_1^2 + 2688k^3 a_0 a_1^2 - 384k\lambda a_0 a_1^2 - 8640k^2 a_0^2 a_1^2 + 15360k a_0^3 a_1^2 - 1440k^3 a_1^4 - 272k^3\lambda a_2 + 2k\lambda^2 a_2 + 416k^2\lambda a_0 a_2 + 4352k^3 a_0^2 a_2 - 384k\lambda a_0^2 a_2 - 7680k^2 a_0^3 a_2 + 7680k a_0^4 a_2 + 5136k^4 a_1^2 a_2 - 20160k^3 a_0 a_1^2 a_2 - 2624k^5 a_2^2 + 208k^3\lambda a_2^2 + 8192k^4 a_0 a_2^2 - 11520k^3 a_0^2 a_2^2 + 800k^5 a_2^3 = 0.$
- (4) $a_1^4 - 1232k^2\lambda a_2 + 2\lambda^2 a_2 + 1040k\lambda a_0 a_2 + 19712k^2 a_0^2 a_2 - 384\lambda a_0^2 a_2 - 19200k a_0^3 a_2 + 7680 a_0^4 a_2 + 51344k^3 a_1^2 a_2 - 768k\lambda a_1^2 a_2 - 146880k^2 a_0 a_1^2 a_2 + 92160k a_0^2 a_1^2 a_2 - 26176k^4 a_2^2 + 1456k^2\lambda a_2^2 + 71168k^3 a_0 a_2^2 - 768k\lambda a_0 a_2^2 - 80640k^2 a_0^2 a_2^2 + 30720k a_0^3 a_2^2 - 48960k^3 a_1^2 a_2^2 + 21152k^4 a_2^3 - 34560k^3 a_0 a_2^3 = 0.$
- (5) $-240k\lambda a_1 + 156\lambda a_0 a_1 + 3840k a_0^2 a_1 - 2880 a_0^3 a_1 + 14112k^2 a_1^3 - 192\lambda a_1^3 - 38880k a_0 a_1^3 + 23040 a_0^2 a_1^3 + 3840k a_1^5 - 27008k^3 a_1 a_2 + 2392k\lambda a_1 a_2 + 122624k^2 a_0 a_1 a_2 - 1152\lambda a_0 a_1 a_2 - 132480k a_0^2 a_1 a_2 + 46080 a_0^3 a_1 a_2 - 92640k^2 a_1^3 a_2 + 76800k a_0 a_1^3 a_2 + 151168k^3 a_1 a_2^2 - 960k\lambda a_1 a_2^2 - 296640k^2 a_0 a_1 a_2^2 + 115200k a_0^2 a_1 a_2^2 - 57600k^3 a_1 a_2^3 = 0.$
- (6) $-1856k^3 a_1^2 + 416k\lambda a_1^2 + 12928k^2 a_0 a_1^2 - 384\lambda a_0 a_1^2 - 23040k a_0^2 a_1^2 + 15360 a_0^3 a_1^2 - 11040k^2 a_1^4 + 15360k a_0 - 120\lambda a_1 + 1920 a_0^2 a_1 +$

- $$17464 k a_1^3 - 21600 a_0 a_1^3 + 3840 a_1^5 - 49664 k^2 a_1 a_2 + 1300 \lambda a_1 a_2 + 144128 k a_0 a_1 a_2 - 72000 a_0^2 a_1 a_2 - 147360 k a_1^3 a_2 + 76800 a_0 a_1^3 a_2 + 356320 k^2 a_1 a_2^2 - 960 \lambda a_1 a_2^2 - 466560 k a_0 a_1 a_2^2 + 115200 a_0^2 a_1 a_2^2 + 53760 k a_1^3 a_2^2 - 265920 k^2 a_1 a_2^3 + 107520 k a_0 a_1 a_2^3 = 0.$$
- (7) $-4288 k^2 a_1^2 + 260 \lambda a_1^2 + 18304 k a_0 a_1^2 - 14400 a_0^2 a_1^2 - 19680 k a_1^4 + 15360 a_0 a_1^4 - 1680 k \lambda a_2 + 624 \lambda a_0 a_2 + 26880 k a_0^2 a_2 - 11520 a_0^3 a_2 + 142928 k^2 a_1^2 a_2 - 768 \lambda a_1^2 a_2 - 256320 k a_0 a_1^2 a_2 + 92160 a_0^2 a_1^2 a_2 + 23040 k a_1^4 a_2 - 87680 k^3 a_2^2 + 2496 k \lambda a_2^2 + 185856 k^2 a_0 a_2^2 - 768 \lambda a_0 a_2^2 - 138240 k a_0^2 a_2^2 + 30720 a_0^3 a_2^2 - 247680 k^2 a_1^2 a_2^2 + 138240 k a_0 a_1^2 a_2^2 + 116736 k^3 a_2^3 - 384 k \lambda a_2^3 - 172800 k^2 a_0 a_2^3 + 46080 k a_0^2 a_2^3 - 23040 k^3 a_2^4 = 0.$
- (8) $-4032 k a_1^2 + 8064 a_0 a_1^2 - 10080 a_1^4 - 720 \lambda a_2 + 11520 a_0^2 a_2 + 153392 k a_1^2 a_2 - 129600 a_0 a_1^2 a_2 + 23040 a_1^4 a_2 - 133248 k^2 a_2^2 + 1248 \lambda a_2^2 + 192000 k a_0 a_2^2 - 69120 a_0^2 a_2^2 - 365760 k a_1^2 a_2^2 + 138240 a_0 a_1^2 a_2^2 + 246144 k^2 a_2^3 - 384 \lambda a_2^3 - 253440 k a_0 a_2^3 + 46080 a_0^2 a_2^3 + 61440 k a_1^2 a_2^3 - 99840 k^2 a_2^4 + 30720 k a_0 a_2^4 = 0.$
- (9) $-12096 a_1 a_2 + 119808 a_1 a_2^2 - 164160 a_1 a_2^3 + 34560 a_1 a_2^4 = 0.$
- (10) $7084 a_1^3 - 40320 k a_1 a_2 + 56448 a_0 a_1 a_2 - 70560 a_1^3 a_2 + 346608 k a_1 a_2^2 - 221760 a_0 a_1 a_2^2 + 53760 a_1^3 a_2^2 - 372480 k a_1 a_2^3 + 107520 a_0 a_1 a_2^3 + 34560 k a_1 a_2^4 = 0.$
- (11) $-25920 a_2^2 + 73440 a_2^3 - 57600 a_2^4 + 7680 a_2^5 = 0$
- (12) $-1344 a_1^2 + 56672 a_1^2 a_2 - 95040 k a_2^2 + 69120 a_0 a_2^2 - 167040 a_1^2 a_2^2 + 223200 k a_2^3 - 115200 a_0 a_2^3 + 61440 a_1^2 a_2^3 - 134400 k a_2^4 + 30720 a_0 a_2^4 + 7680 k a_2^5 = 0.$

step 3:

Upon solving the above system with respect to the unknowns k , λ , a_0 , a_1 , a_2 , and using the elimination method of Wu [11] and the Mathematica software [7], we consider only the following solutions:

$$a_1 = 0, \quad a_2 = \frac{3}{4}, \quad k = \pm\sqrt{\lambda}, \quad a_0 = \pm\frac{\sqrt{\lambda}}{2}.$$

(The other solutions of the above system are given by $a_1 = 0$, $a_2 = 0$, k, λ, a_0 arbitrary constants. But this case is not interesting).

step 4:

According to (2.11) and taking into account the results of the previous step (solutions of the system of algebraic equations) after simplifications, we restrict ourselves to considering only the following solutions for (2.8) :

For $\lambda > 0$,

- (1) $v_1(\xi) = \frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \cot^2\left(\lambda^{\frac{1}{4}}\xi\right).$..
- (2) $v_2(\xi) = -\frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \coth^2\left(\lambda^{\frac{1}{4}}\xi\right).$

$$(3) v_3(\xi) = \frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \tan^2 \left(\lambda^{\frac{1}{4}} \xi \right).$$

$$(4) v_4(\xi) = -\frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \tanh^2 \left(\lambda^{\frac{1}{4}} \xi \right).$$

step 5 :

We obtain solutions for system (2.1). By (2.2) and (2.7), the solutions for $u(x, t)$ and $w(x, t)$ are given by (for $\lambda > 0$):

$$(1) u_1(x, t) = \frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \cot^2 \left(\lambda^{\frac{1}{4}} (x + \lambda t) \right)$$

$$w_1(x, t) = \frac{\lambda^{\frac{3}{2}}}{6}$$

$$(2) u_2(x, t) = -\frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \coth^2 \left(\lambda^{\frac{1}{4}} (x + \lambda t) \right).$$

$$w_2(x, t) = -\frac{\lambda^{\frac{3}{2}}}{6}$$

$$(3) u_3(x, t) = \frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \tan^2 \left(\lambda^{\frac{1}{4}} (x + \lambda t) \right).$$

$$w_3(x, t) = \frac{\lambda^{\frac{3}{2}}}{6}$$

$$(4) u_4(x, t) = -\frac{\sqrt{\lambda}}{2} + \frac{3}{4}\sqrt{\lambda} \tanh^2 \left(\lambda^{\frac{1}{4}} (x + \lambda t) \right).$$

$$w_4(x, t) = -\frac{\lambda^{\frac{3}{2}}}{6}$$

Remark 1: One also can consider solutions with $k = -\sqrt{\lambda}$.

Remark 2: In the case $w(x, t) = 0$, system (2.1) is reduced to well-known Kaup–Kupershmidt equation [6], [8], [9], [12]

$$u_t = u_{xxxxx} - 20uu_{xxx} - 50u_x u_{xx} + 80u^2 u_x \tag{2.12}$$

which is a particular case of a class of fifth order PDEs with four parameters

$$u_t + \rho u_{xxxxx} + \gamma u u_{xxx} + \beta u_x u_{xx} + \alpha u^2 u_x = 0 \tag{2.13}$$

which has known solutions in some cases:

- For $\alpha = 30, \beta = 20, \gamma = 10, \rho = 1$ [6].
- For $\alpha = 5, \beta = 5, \gamma = 5, \rho = 1$ (2.13) reduces to Sawada and Kotera (SK) equation [6].
- For $\alpha = 20, \beta = 25, \gamma = 10, \rho = 1$ [6].

Remark 3: A plenty of exact solutions for the Kaup–Kupershmidt equation can be found using the inverse scattering transform for the corresponding spectral problem [13]. In [8], Sergyeyev present a *zero curvature representation ZCR* for the M–N–W system. Knowing a *ZCR* involving an essential (spectral) parameter enables one to use the inverse scattering transform and construct plenty of exact solutions for the system in question, including multisoliton and finite-gap solutions [8].

Acknowledgments: I want to express my gratitude to professor A. Sinitsyn for his enthusiastic guidance and help. I also would like to express my sincere thanks to the anonymous referee for the comments suggestions and corrections that have improved the presentation of this paper.

References

- [1] R. CONTE & M. MUSETTE, Link between solitary waves and projective Riccati equations, *J. Phys. A Math.* **25** (1992), 5609–5623.
- [2] E. INC & M. ERGÜT, New Exact Travelling Wave Solutions for Compound KdV-Burgers Equation in Mathematical Physics, *Applied Mathematics E-Notes* **2** (2002), 45–50.
- [3] J. MEI, H. ZHANG & D. JIANG, New exact solutions for a Reaction-Diffusion equation and a Quasi-Camassa-Holm Equation, *Applied Mathematics E-Notes* **4** (2004), 85–91.
- [4] Z. YAN, The Riccati equation with variable coefficients expansion algorithm to find more exact solutions of nonlinear differential equation, *Comput. Phys. Comm.* **152** (2003) 1, 1–8. Preprint version available at <http://www.mmrc.iss.ac.cn/pub/mm22.pdf/20.pdf>
- [5] Z. YAN, An improved algebra method and its applications in nonlinear wave equations, *Chaos Solitons Fractals* **21** (2004) 4, 1013–1021. Preprint version available at <http://www.mmrc.iss.ac.cn/pub/mm22.pdf/19.pdf>
- [6] D. BALDWIN, U. GOKTAS, W. HEREMAN, L. HONG, R.S. MARTINO & J.C.MILLER, Symbolic computation of exact solutions expressible in hyperbolic and elliptic functions for nonlinear PDFs, *J. Symbolic Compt.* **37** (2004) 6, 669–705; Preprint version: [nlin.SI/0201008\(arXiv.org\)](http://nlin.SI/0201008(arXiv.org))
- [7] A. SALAS & C. GOMEZ, El Software mathematica en la busqueda de soluciones exactas de ecuaciones diferenciales no lineales en derivadas parciales mediante la ecuacion de Riccati, *Memorias Primer Seminario Internacional de Tecnologías en Educación Matemática*, Universidad Pedagógica Nacional **1** (2005), 379–387
- [8] A. SERGYEYEV, Zero curvature representation for a new fifth-order integrable system, to appear in *Fund. Prink. Mat.* **12** 2006 (in Russian); English translation to appear in *J. Math. Sci.*; preprint version (in English): [nlin.SI/0604064\(arXiv.org\)](http://nlin.SI/0604064(arXiv.org))
- [9] N.A. KUDRYASHOV, Nonlinear differential equations with exact solutions expressed via the Weierstrass function, *Z. Naturforsch. A* **59** (2004) 7-8, 443–454. Preprint version: [nlin.CD/0312035\(arXiv.org\)](http://nlin.CD/0312035(arXiv.org))
- [10] E. FAN & Y.C. HON, Generalized tanh Method Extended to Special Types of Nonlinear Equations, *Z. Naturforsch. A* **57**(2002) 8, 692–700.
- [11] L. ZHI, Y. NOTAKE, H. KAI, M.T. NODA & K. I. SHIRAISHI, Hibrid method for solving Polinomial Equations, *NLA99: Computer algebra* (Saitama, 1999), 91–104, *Josai Math. Monogr.* **2**, Josai Univ. Sakado, 2000. Preprint version available at <http://www.mmrc.iss.ac.cn/~lzhi/Publications/atcm99.pdf>
- [12] A.V. MIKHAILOV, V. NOVIKOV & J.P.WANG, On clasification of integrable non-evolutionary equation, Preprint: [nlin.SI/0601046\(arXiv.org\)](http://nlin.SI/0601046(arXiv.org))
- [13] D.J. KAUP, On the inverse scattering problem for cubic eingevalue problems of the class $\phi_{xxx} + 6q\phi_x + 6r\phi = \lambda\phi$, *Stud. Appl. Math.* **62** (1980), 189–216.

(Recibido en junio de 2006. Aceptado en septiembre de 2006)

DEPARTAMENTO DE MATEMÁTICAS
UNIVERSIDAD NACIONAL DE COLOMBIA
CR. 30 CLL. 45
BOGOTÁ, COLOMBIA
e-mail: cagomezsi@unal.edu.co

