

Antifungal activity of extracts, essential oil and constituents from *Petroselinum crispum* against *Colletotrichum acutatum*

Actividad antifúngica de extractos, aceite esencial y constituyentes de *Petroselinum crispum* contra *Colletotrichum acutatum*

doi: 10.15446/rfnam.v71n3.68284

Rodrigo Pineda¹, Samuel Vizcaíno², Carlos M. García³, Jesús H. Gil⁴ and Diego Durango^{3*}

ABSTRACT

Keywords:

Apiaceae
Parsley
Extracts
Essential oil
GC-MS
Parsley-apiole

The effect of extracts, essential oil, and their major constituents from parsley (*Petroselinum crispum* (Mill.) Fuss.) against the phytopathogenic fungus *Colletotrichum acutatum* was evaluated by the poisoned agar method. Results showed that all extracts, along with the essential oil, significantly inhibit the radial growth of *C. acutatum* at concentrations higher than 100 µg mL⁻¹. The higher activity was found for the essential oil followed by the *n*-hexane extract. Analysis by gas chromatography with mass spectroscopy (GC-MS) of *n*-hexane extract and the essential oil of *P. crispum* showed that the major components correspond to the phenylpropanoids myristicin and *parsley*-apiole. Both compounds were isolated by conventional chromatographic techniques and their structures elucidated by spectroscopic methods. Myristicin and *parsley*-apiole displayed a significant inhibitory effect against *C. acutatum*. The highest fungistatic activity was found to *parsley*-apiole with IC₅₀ value of 40 µg mL⁻¹. In conclusion, parsley may be a good source of antifungal compounds to control *C. acutatum*.

RESUMEN

Palabras clave:

Apiaceae
Perejil
Extractos
Aceite esencial
CG-EM
Parsley-apiol

Se evaluó el efecto de extractos, el aceite esencial y los componentes principales del perejil (*Petroselinum crispum* (Mill.) Fuss.) contra el hongo fitopatógeno *Colletotrichum acutatum* mediante el método del agar envenenado. Los resultados mostraron que todos los extractos, junto con el aceite esencial, inhibieron significativamente el crecimiento radial de *C. acutatum* en concentraciones superiores a 100 µg mL⁻¹. La mayor actividad se encontró para el aceite esencial, seguido por el extracto de *n*-hexano. El análisis por cromatografía de gases con espectroscopia de masas (CG-EM) del extracto de *n*-hexano y el aceite esencial de *P. crispum* mostró que los componentes principales corresponden a los fenilpropanoides miristicina y *parsley*-apiol. Ambos compuestos fueron aislados por técnicas cromatográficas convencionales y sus estructuras elucidadas por métodos espectroscópicos. La miristicina y el *parsley*-apiol mostraron un efecto inhibitorio significativo contra *C. acutatum*. La actividad fungistática más alta se encontró para *parsley*-apiol con un valor CI₅₀ de 40 µg mL⁻¹. En conclusión, el perejil puede ser una buena fuente de compuestos antimicrobianos para controlar el *C. acutatum*.

¹ Universidad de Talca. Av Lircay, Talca, Región del Maule, Chile.

² Facultad de Ciencias Exactas y Naturales. Universidad de Antioquia. AA 1226, Medellín, Colombia.

³ Facultad de Ciencias. Universidad Nacional de Colombia. AA . 3840, Medellín, Colombia.

⁴ Facultad de Ciencias Agrarias. Universidad Nacional de Colombia. AA 1779, Medellín, Colombia.

* Corresponding author: <dldurango@unal.edu.co>

Anthraxnose, a disease caused by the phytopathogenic fungus *Colletotrichum acutatum* Simmonds affects different plants, such as tamarillo (*Solanum betaceum* Cav.), tomato (*Solanum lycopersicum* L.), strawberry (*Fragaria x ananassa* Duch.), apple (*Malus domestica* Borkh.), blackberry (*Rubus glaucus* Benth.), mango (*Mangifera indica* L.), among other fruits of economic importance (Damm *et al.*, 2012; Wharton and Diéguez-Uribeondo, 2004). Its presence occurs in temperate and tropical climates around the world. Conventionally, its control has been based on the application of synthetic fungicides, which can cause adverse effects to the environment and to man because of their low selectivity (Gao *et al.*, 2017; Wharton and Diéguez-Uribeondo, 2004; Gaviria-Hernández *et al.*, 2013). In addition, as a result of the gradual development of resistance by the microorganism to this class of substances used for its control, it has become necessary to apply higher and more constant doses in the field, which has led to an increase in the production costs and furthers the problem of food security (Jílková *et al.*, 2015).

In response to the above, new alternatives of control disease in important crops are needed. One of the approaches that have become more prevalent in recent years is the one that uses essential oils and extracts of plants or their major components for the control of phytopathogenic microorganisms (Grande-Tovar *et al.*, 2018; Nazzaro *et al.*, 2017). Alternative methods to control *C. acutatum* have been developed; these include, the use of natural antimicrobials, biocontrol agents, ultraviolet radiation and resistant cultivars (Falconí and Yáñez-Mendizábal, 2018). For example, He *et al.* (2018) determined the efficacy and possible mechanism of cinnamon essential oil on inhibition of *C. acutatum* isolated from kiwi fruit. Meanwhile, Rashid *et al.* (2018) evaluated the antifungal effects of *Rhus coriaria* L. fruit extracts against tomato anthracnose caused by *Colletotrichum acutatum*. The results indicate that *R. coriaria* extract can be utilised as a deterrent and curative natural product for the anthracnose disease in tomatoes. Some substances of this type have demonstrated a high antimicrobial activity, good selectivity, low toxicity, and little persistence in the environment (Tripathi and Dubey, 2004).

Edible and herbal plants/spices such as garlic, oregano, cinnamon, clove, parsley, among other are an important source of essential oils and have been employed

alone or combined as natural food preservatives with antimicrobial properties (Pisoschi *et al.*, 2018). Particularly, parsley is an easy plant to grow with low production costs, and is an aromatic herb commonly used in Colombia. The essential oil and different extracts obtained from parsley have shown antimicrobial activity against Gram positive and negative bacteria (Khalil *et al.*, 2018; Semeniuc *et al.*, 2017; Farzai *et al.*, 2013). To the extent of our knowledge, there are few works to establish the antifungal potential of compounds obtained from parsley. In the present work, the antifungal activity against *C. acutatum* of *n*-hexane, dichloromethane, ethyl acetate, and methanol extracts, the essential oil, and their main constituents from parsley was evaluated.

MATERIALS AND METHODS

Chemicals

NaOH, H₂SO₄, HCl, Na₂SO₄, glucose, K₂HPO₄, NaNO₃, MgSO₄, FeSO₄ were purchased from Merck KGaA (Darmstadt, Germany) and Sigma-Aldrich (St. Louis, MO, USA). Solvents methanol (MeOH), dichloromethane (CH₂Cl₂), ethyl acetate (EtOAc), and *n*-hexane were acquired from Merck. All chemicals and reagents used were of analytical grade.

General methods

Thin layer chromatography was performed with silica gel F₂₅₄ (0.2 mm Merck) plates. Compounds were detected by UV fluorescence (254 and 365 nm) and/or spraying with AcOH-H₂SO₄-H₂O (143:28:30, v/v) "Universal stain", followed by heating at 80 °C for 1–2 min. ¹H and ¹³C NMR spectra were recorded in CDCl₃ on a Bruker AMX 300 spectrometer (¹H at 300 MHz; ¹³C 75.5 MHz). Chemical shifts (δ) are reported in ppm relative to the residual solvent signals (CDCl₃: δ_H 7.24 and δ_C 77 ppm) and coupling constants (*J*) in Hz. Mass spectrometry analysis was performed using a Hewlett-Packard 6890 (Agilent Technologies) gas chromatograph coupled with a HP 5973 MSD (Mass selective detector-Quadrupole type). A HP-5 MS capillary column (30 m x 0.25 mm x 0.25 μm, Agilent Technologies) was used. Hydrogen was the carrier gas, having a constant flow rate of 1.0 mL min⁻¹ and split ratio 1:10. The temperature was programmed as follows: column temperature, 50 °C; injector temperature, 150 °C; detector temperature, 280 °C. The oven temperature was programmed from 50 to 250 °C at 10 °C min⁻¹ and, then, held isothermally

for 6 min. The MS system operated with an electron impact of 70 eV, an acceleration voltage of 1.1 kV, and scan range of m/z 40-500. The products were identified by comparison of their mass spectra with those of the US National Institute of Standards and Technology (NIST) mass spectral library (Version 2008). Quantitative analysis of each essential oil component (expressed in percentages) was carried out by peak area normalization measurement.

Plant material and extraction procedure

Extracts: Fresh parsley plant materials (curly leaf) were purchased from a local supermarket in Medellín, Colombia. Samples were washed under tap water and roots were discarded. Leaves and stems were homogenized in a laboratory blender. Then, vegetal material (900 g) was extracted sequentially with 700 mL of each solvent (*n*-hexane (HEX), dichloromethane (DCM), ethyl acetate (EA), and methanol (ME)) by percolation (5 cycles per 24 h) a room temperature until exhaustion. Extracts were dried over anhydrous sodium sulfate, filtered, evaporated using a rotary evaporator (Buchi Heating Bath B-490, Buchi Rotavapor R-200) under reduced pressure at 40 °C, and stored in a refrigerator at 4 °C in amber vials. **Essential oil:** The essential oil (EO) was obtained by hydrodistillation. A quantity of 900 g of fresh leaves and stems of parsley was coarsely chopped and immersed in 1.5 L of distilled water contained in a 3 L flask. Distillation was carried out using a modified Clevenger apparatus. The extraction process was performed for 2 h after the first drop of distillate until complete exhaustion of the plant. Then, sodium chloride (1 g) and 40 mL of dichloromethane (2x20 mL) was added with the aqueous distillate in a separating funnel. The organic layer was separated, dried over sodium sulfate, filtered, evaporated using a rotary evaporator (Buchi Heating Bath B-490, Buchi Rotavapor R-200) under reduced pressure at 40 °C, and stored in a refrigerator at 4 °C in amber vials.

Isolation and identification of major constituents in bioactive materials

The bioactive *n*-hexane extract and essential oil from *P. crispum* were measured chromatographically with silica gel (Merck, Darmstadt, Germany, silica gel 60, 230-400 mesh) column and eluted with gradients of *n*-hexane:EtOAc (100:0, 95:5, 90:10, 80:20, 70:30, 0:100 v/v). Next, the

n-hexane and *n*-hexane:EtOAc (95:5) fractions were measured in a silica gel column chromatograph eluted with *n*-hexane and gradients *n*-hexane:CH₂Cl₂ (98:2, 95:5, and 90:10 v/v). Fractions 2 to 4 were mixed and measured through a Sephadex LH-20 column chromatograph eluted with *n*-hexane:CH₂Cl₂:MeOH, 50:25:25, v/v affording two compounds, which were identified through spectroscopic methods (one-dimensional (¹H and ¹³C) and two-dimensional (COSY, HSQC and HMBC) NMR, as well as EI/MS analysis.

Antifungal activity

The strain of phytopathogenic fungus *C. acutatum* was isolated from commercial fruits of tamarillo (*S. betaceum*) affected by the disease and purchased at the local markets in the city of Medellín (Colombia). Identification was performed by the Phytopathology Laboratory of the Universidad Nacional de Colombia-Sede Medellín, evaluating morphological and molecular characteristics (Afanador-Kafuri *et al.*, 2014). The evaluation of antifungal activity was carried out using the poisoned agar method (Balouiri *et al.*, 2016). Essential oil, extracts, and pure compounds were mixed with ethanol (<0.2% EtOH) and tested at 100, 200, and 400 µg mL⁻¹. Thymol (2-isopropyl-5-methylphenol) and carbendazim (methyl benzimidazol-2-ylcarbamate) at 100 µg mL⁻¹ were used as positive controls. Mycelial growth (mm) were determined as mean values ± standard deviation of three replicates of mycelium diameter. In addition, for bioactive materials and their major constituents, the concentration that inhibits 50% fungal mycelia growth (IC₅₀) was determined according to Rivillas and Soriano (2007).

RESULTS AND DISCUSSION

Table 1 shows the extraction yield for the four solvents of different polarities –*n*-hexane (HEX), dichloromethane (DCM), ethyl acetate (EA), and methanol (ME)–, and the essential oil (EO) collected by hydrodistillation from *P. crispum*. The order of increasing yield was HEX < DCM < EA < ME, coinciding with a growing polarity of the extracting solvent. Thus, the highest yield was achieved by the most polar solvent, methanol. The *n*-hexane extract presented approximately 10 times less of the weight reached by the methanol extract. This suggests that the proportion of low polarity products extractable with hexane in *P. crispum* is low, compared to those of high polarity present in extracts in ethyl acetate and methanol. The yield from essential oil hydrodistillation

extraction was closer to the extraction yield from *n*-hexane. Overall, the total amount of extractable material with organic solvent in *P. crispum* was 13.73 g (1.42%).

Then, all extracts and essential oil were tested for antifungal activity using the poisoned-food technique at 100, 200, and 400 $\mu\text{g mL}^{-1}$, according to Balouiri *et al.* (2016).

Table 1. Yields of essential oil and extracts for the leaves and stems of *P. crispum*.

Extract	Mass (g) ^a	%(p/p) ^b
HEX extract	0.72	0.08
DCM extract	1.67	0.19
EA extract	2.03	0.22
ME extract	8.32	0.92
Essential oil	0.99	0.11

^a Crude weight results from 900 g of fresh aerial parts. ^b Yield (%) = [crude weight/900 g]×100

The inhibitory effects of organic extracts and essential oil of *P. crispum* at 400 $\mu\text{g mL}^{-1}$ on the growth of *C. acutatum* are presented in Figure 1. All extracts displayed a significant inhibitory effect in the growth of *C. acutatum* after 264 h. In general, the growth of fungal mycelium was dependent of the type and amount of extract in the culture medium. The *n*-hexane extract caused the least growth of *C. acutatum*; after 264 hours, the radial growth was almost 30 mm less than the control. Higher

polarity extracts, methanol followed by ethyl acetate, had the lowest inhibitory effect on *C. acutatum*. After 72 h, the *n*-hexane extract showed an inhibition percentage of 57%. Then, the fungistatic effect presented a slight decrease, reaching 48% inhibition after 264 h.

The essential oil of *P. crispum* displayed the highest inhibitory effect. At 400 $\mu\text{g mL}^{-1}$, essential oil of *P. crispum* inhibited the radial growth of *C. acutatum*




Figure 1. Effect of extracts and essential oil of *Petroselinum crispum* at 400 $\mu\text{g mL}^{-1}$ on radial mycelial growth of *Colletotrichum acutatum*.

completely during the first 72 h. Similarly, the inhibition percentage was slightly decreased at 78% after 264 h. Even at 100 µg mL⁻¹, essential oil exhibited inhibition percentages of *C. acutatum* of almost 60% during the 264 h of the analysis (*data not shown*). Thymol and carbendazim® presented a significantly higher antifungal effect (inhibition percentages of 100%) as compared to all evaluated extracts. Carbendazim is commonly used for control of anthracnose disease, both in the field and postharvest. However, some plant pathogens, such as *C. acutatum* and *C. gloeosporioides*, have strains resistant to carbendazim (Zhang and Huang, 2007). Meanwhile, thymol has been recognized for its potent antifungal activity (Numpaque *et al.*, 2011). The IC₅₀ values calculated for the *n*-hexane extract and the essential oil of *P. crispum* were 316.7 and 185.1 µg mL⁻¹, respectively. The high antifungal activity of essential oil of *P. crispum* found in the present study is in agreement

with that reported by Linde *et al.* (2016). The chemical composition of essential oil and *n*-hexane extract from leaves and stems of *P. crispum* was analyzed by means of GC-MS. The detected constituents (relative amounts higher than 0.1%) are shown in Tables 2 and 3. Analysis by GC-MS led to the identification and quantification of about 23 components. Among them were 12 in the essential oil and 22 in the *n*-hexane extract, comprising between 93.5-99.1% of the total composition of the essential oil/*n*-hexane extract. Phenylpropenes were the main components detected in the essential oil (77.0%) and *n*-hexane extracts (48.7%) for *P. crispum*. Parsley- or dill-apiole (27.6 and 43.2% in *n*-hexane extract and essential oil, respectively) and myristicin or sarisan (18.7 and 30.8% in *n*-hexane extract and essential oil, respectively) were the major constituents on *P. crispum*. The phenylpropene safrole was also detected in low proportion.

Table 2. Relative abundance (%RA) of major compounds obtained by GC-MS (%) from *n*-hexane extract of *P. crispum*.

TR (min)	Component	%RA	Fragment (m/z)
5.57	D-Limonene	0.2	68(100), 93(59), 67(45), 79(23), 94(23), 136(23)
5.71	β-Phellandrene	8.9	93(100), 77(27), 91(25), 136(18), 79(17), 94(15)
6.87	Terpinolene	1.5	121(100), 93(98), 136(73), 79(34), 77(30), 91(31)
7.52	p-Cymene	5.7	117(100), 132(81), 115(63), 91(54), 92(23), 65(23)
7.76	1,3,8-p-Menthatriene	11.2	119(100), 91(74), 134(66), 41(30), 39(27), 105(21)
9.34	Geranial	2.1	69(100), 41(46), 84(29), 94(17), 83(14), 67(12)
10.34	Mellilotal	0.8	119(100), 91(64), 134(37), 65(14), 120(10), 39(8)
10.73	Pentylidenecyclohexane	0.6	67(100), 109(77), 81(60), 96(54), 41(37), 152(37)
11.19	3-Guaiacylpropanol	1.5	91(100), 121(63), 166(30), 77(25), 65(19), 51(18)
12.13	Safrole	2.4	162(100), 104(39), 131(35), 135(31), 103(27), 77(27)
13.19	β-Caryophyllene	0.6	93(100), 133(92), 91(86), 41(77), 79(76), 69(75)
14.27	2-Methylisoborneol	1.2	95(100), 107(25), 43(22), 108(21), 41(16), 110(15)
15.33	β-Sesquiphellandrene	0.9	69(100), 41(68), 93(46), 55(29), 91(28), 92(24)
15.97	Ledol	1.2	43(100), 41(61), 69(53), 109(50), 81(34), 55(34)
17.79	Myristicin or sarisan	18.7	192(100), 91(25), 165(23), 65(16), 119(15), 161(14)
21.17	Parsley- or dill-apiole	27.6	222(100), 207(24), 149(16), 177(16), 223(14), 195(11)
23.86	Hexadecanoic acid	3.8	43(100), 73(97), 60(90), 41(77), 57(77), 55(65)
24.14	Ethyl linoleate	0.6	79(100), 67(63), 95(60), 93(52), 55(46), 81(44)
24.35	Methyl linoleate	1.8	79(100), 67(78), 41(61), 55(57), 93(47), 81(43)
24.78	Phenethyl hexanoate	2.1	104(100), 105(18), 43(12), 99(6), 71(5), 91(5)
28.82	Linoleic acid	3.2	67(100), 81(88), 82(73), 95(63), 68(60), 55(60)
29.60	α-Linolenic acid	2.5	41(100), 79(78), 67(73), 55(62), 81(38), 80(37)

Monoterpenes (27.5%), fatty acids (11.9%), oxygenated monoterpenes (3.3%), sesquiterpenes (2.7%), and others (5.0%) were also found in the extract. Meanwhile, 7.5%

monoterpenes, 0.6% oxygenated monoterpenes, 3.9% sesquiterpenes, and 4.5% others were detected in the essential oil.

Table 3. Relative abundance of major compounds obtained by GC-MS (%) from essential oil of *P. crispum*.

TR (min)	Component	%RA	Fragment (m/z)
8.08	D-Limonene	0.7	68(100), 93(59), 67(45), 79(23), 94(23), 136(23)
8.30	β -Phellandrene	1.6	93(100), 77(27), 91(25), 136(18), 79(17), 94(15)
8.53	<i>p</i> -Cymene	4.4	117(100), 132(81), 115(63), 91(54), 92(23), 65(23)
10.75	1,3,8- <i>p</i> -Menthatriene	0.8	119(100), 91(74), 134(66), 41(30), 39(27), 105(21)
12.87	β -Phenylethanol	0.7	91(100), 92(56), 65(23), 122(22), 39(12), 51(9)
13.36	Geranial	0.6	69(100), 41(46), 84(29), 94(17), 83(14), 67(12)
14.81	Mellilotal	3.8	119(100), 91(64), 134(37), 65(14), 120(10), 39(8)
17.24	Safrole	3.0	162(100), 104(39), 131(36), 103(35), 77(27), 161(26)
18.68	β -Caryophyllene	1.1	41(100), 69(98), 93(94), 133(65), 79(61), 91(55)
22.03	Ledol	2.8	43(100), 41(61), 69(53), 109(50), 81(34), 55(34)
24.72	Myristicin or sarisan	30.8	192(100), 91(24), 165(22), 119(16), 65(15), 39(14)
29.41	<i>Parsley-</i> or <i>dill-</i> apiole	43.2	222(100), 207(24), 149(16), 177(16), 223(14), 195(11)

Previous reports differ on the major constituents in essential oil of curly leaf parsley. Some authors have found that monoterpenes are the dominant compounds. Thus, El-Zaedi *et al.* (2016b) reported that 1,3,8-*p*-menthatriene (38.4-48.8%) and β -phellandrene (22.2-29.5%) are the major constituents in the essential oil of parsley grown in Spain, obtained by hydrodistillation using a Deryng system (the Polish version of the Clevenger apparatus). Kurowska and Galazka (2006) found α -pineno (32.0%) to be the major constituent of the essential oil of *P. crispum* cultivated in Poland, also having β -pineno (19.0%), myristicin (18.3%), and apiole (10.1%). On the other hand, Camilotti *et al.* (2015) reported that phenylpropanoides were the principal compounds, comprising 52.07% of the oil and consisting mainly of apiole (41.05%) and myristicin (5.08%). According to Zhang *et al.* (2006), myristicin (32.75%) and apiole (17.54%) were the major constituent in essential oil of parsley cultivated in China. Stankovic *et al.* (2004) found that apiole (57.0%) was the major constituent of essential oil of *P. crispum* cultivated in Serbia and Montenegro. Linde *et al.* (2016) reported apiole (50.3%) and myristicin (14.0%) as major constituents in the essential oil of parsley cultivated in Brazil. The chemical

composition of the parsley is also dependent on the variety analyzed; in the flat-leaf parsley (*Petroselinum sativum*) contents of myristicin and apiol have been found that vary between 25-47% and 18-23%, respectively (Romeilah *et al.*, 2010; El-Zaedi *et al.*, 2016a). In this way, apiole and/or myristicin have always been among the major constituents of essential oils obtained from parsley. This diverse chemical composition could be a result of several parameters including geographical origin, vegetative stage of plant, parsley cultivar, storage condition, and extraction method, as well as their interactions with microorganisms and insects (Chatzopoulou and Katsiotis, 1995). In addition, abiotic factors like luminosity, temperature, rainfall, nutrition, time of the day for harvest, and post-harvest techniques are also important to the chemical composition of essential oil (Morais and Castanha, 2012). The results by Camilotti *et al.* (2015), Zhang *et al.* (2006), Linde *et al.* (2016) and Stankovic *et al.* (2004) agree quite well with the results of this study.

Then, the *n*-hexane extract and essential oil from *P. crispum* were submitted to chromatography on silica gel, using *n*-hexane and *n*-hexane:EtOAc (95:5). Further

sequential purification by column chromatography using silica gel (eluted with *n*-hexane and *n*-hexane:CH₂Cl₂, 98:2) and Sephadex LH-20 (eluted with *n*-hexane:CH₂Cl₂:MeOH, 50:25:25), which were identified through one-dimensional (¹H and ¹³C) and two-dimensional (COSY, HSQC, and HMBC) NMR spectroscopic methods, in addition to EI/MS analysis.

Parsley-apiole: The compound was isolated as a yellowish oil; C₁₂H₁₄O₄. EI-MS m/z: 222(100)[M]⁺, 207 (26), 195 (11), 191 (11), 177 (18), 149 (16). ¹H NMR (300 MHz, CDCl₃): δ 6.32 (1H, s, H-2), 5.95 (2H, s, -OCH₂O-), 5.90-6.00 (1H, m, H-8), 5.08 (1H, dd, J = 1.5, 12.9, H-9b), 5.03 (1H, dd, J = 1.5, 9.6, H-9a), 3.89 (3H, s, -OCH₃), 3.86 (3H, s, -OCH₃), 3.31 (2H, dd, J = 1.5, 6.3, H-7). ¹³C NMR (75 MHz, CDCl₃): δ: 138.9 (C-3), 138.6 (C-5), 137.2 (C-8), 136.1

(C-2), 135.0 (C-4), 125.6 (C-9), 115.1 (C-1), 108.1 (C-6), 101.3 (-OCH₂O-), 59.9 (-OCH₃), 56.7 (-OCH₃), 33.9 (C-7).

Myristicin: The compound was isolated as a colorless oil; C₁₁H₁₂O₃. EI-MS m/z: 192(100)[M]⁺, 165 (22), 161 (13), 119 (16), 91 (24). ¹H NMR (300 MHz, CDCl₃): δ 6.29 (1H, d, J = 1.4, H-2), 6.26 (1H, d, J = 1.4, H-6), 5.83 (2H, s, -OCH₂O-), 5.77 (1H, m, H-8), 5.03 (1H, dd, J = 1.5, 17.0, H-9a), 4.92 (1H, dd, J = 1.5, 8.1, H-9b), 3.79 (3H, s, -OCH₃), 3.21 (2H, d, J = 6.3, H-7). ¹³C NMR (75 MHz, CDCl₃): δ: 144.9 (C-4), 143.6 (C-5), 137.4 (C-3), 136.7 (C-8), 125.6 (C-1), 115.9 (C-9), 107.6 (C-2), 102.7 (C-6), 101.3 (-OCH₂O-), 50.9 (-OCH₃), 33.3 (C-7). Spectral analyses were in agreement with those reported in the literature (Di Stefano *et al.*, 2011; Razzaghi-Abyaneh *et al.*, 2007). Chemical structures are presented in Figure 2.




Figure 2. Isolated compounds from *P. crispum*.

Antifungal activity of isolated compounds was evaluated for *C. acutatum*. Results are presented in Figures 3 and 4. The myristicin (a phenylpropanoid with one methylenedioxy and one methoxy group) presented a moderate fungistatic activity for all the evaluated concentrations, as compared to thymol and the carbamate carbendazim®. However, it was observed that even at the concentration of 100 µg mL⁻¹, myristicin presented a significant reduction of mycelial growth of the microorganism, as compared to the control and solvent control. In general, the fungistatic activity of myristicin was concentration dependent. At 200 µg mL⁻¹, the inhibition percentage was greater than 35%, even after 264 h. At this time, the concentration of 400 µg mL⁻¹ showed a 55% inhibition.

The moderate antifungal property of myristicin has been previously reported (Moreira-Valente *et al.*, 2015;

Meepagala *et al.*, 2005). According to Moreira-Valente *et al.* (2015), myristicin at 0.3% inhibited *Aspergillus flavus* and *A. ochraceus* 85 and 80%, respectively.

Parsley-Apiole (a phenylpropanoid with one methylenedioxy and two methoxy groups) had a high fungistatic activity against the phytopathogenic fungus *C. acutatum*. The development of fungal colonies was initiated only after 24 hours of inoculation at a concentration of 75 µg mL⁻¹ and after 48 hours for other concentrations evaluated. Growth inhibition percentages were found to be between 100 and 65% after 96 h at concentrations greater than 100 µg mL⁻¹. The IC₅₀ calculated for *parsley-apiole* was 40.0 µg mL⁻¹. In an earlier work on the antifungal properties of *parsley-apiole*, it was shown to have a varying degree of growth inhibitory effects against *B. cinerea*, *C. acutatum*, *C. fragariae*, *C. gloeosporioides*, and *F. oxysporium*




Figure 3. Effect of myristicin on radial mycelial growth of *C. acutatum*.




Figure 4. Effect of parsley-apiole on radial mycelial growth of *C. acutatum*.

(Meepagala *et al.*, 2005). The authors showed that *parsley*-apiole exhibited high antifungal activity against *B. cinerea* but low activity against *C. acutatum*, *C. fragariae*, *C. gloeosporioides*, and *F. oxysporium* in the microdilution assay (Meepagala *et al.*, 2005). This apparent discrepancy with respect to the present results could be attributed to the fact that different strains of *C. acutatum* and different methods were used to evaluate antifungal activity. In addition, *parsley*-apiole and myristicin have been reported as inhibitors of aflatoxin G₁ production in *Aspergillus parasiticus* (Razzaghi-Abyaneh *et al.*, 2007).

Both oxygenated phenylpropenes have an allyl side chain bonded to the aromatic ring which confers a lipophilic characteristic. Although not much is known about the mechanism of antifungal action of phenylpropenes, it has been reported that their lipophilic characteristic enables them to enter between the fatty-acid-chain constituents of the membrane lipid bilayers and alter the fluidity and permeability of the cell membranes (de Almeida *et al.*, 2009). Remarkably, although the structural difference between *parsley*-apiole and myristicin is only seen in the presence of one methoxyl group on the aromatic ring, *parsley*-apiole was significantly more active than myristicin against *C. acutatum*. These results may suggest that electronic and/or steric factors in phenylpropenes might be important for antifungal activity. Further research to obtain information on the structure-activity relationship is required.

CONCLUSIONS

In the present study, the mycelial growth inhibition of *Colletotrichum acutatum* using essential oil obtained by hydrodistillation and four successive extracts (*n*-hexane, dichloromethane, ethyl acetate, and methanol) from fresh leaves and stems of parsley (*P. crispum* curly leaf) was studied. Essential oil and *n*-hexane extract exhibited the strongest antifungal activity against *C. acutatum*. GC/MS analysis demonstrated that the dominant components of both bioactive materials were phenylpropenes, specifically *parsley*-apiole (or some isomer) and myristicin (or some isomer). Both constituents were isolated, and their structures unambiguously confirmed by spectroscopic methods. Then, antifungal activity against *C. acutatum* of these phenylpropenoids was evaluated. *Parsley*-apiole exhibited a strong fungistatic activity. Accordingly, *parsley*-apiole could be suggested as a structural template for developing new antifungal agents.

ACKNOWLEDGMENTS

This work was supported by the Universidad Nacional de Colombia. We thank Facultad de Ciencias and Escuela de Química for granting scholarships to some authors. The authors are grateful to Dr. W. Quiñones for NMR measurements.

REFERENCES

- Afanador-Kafuri L, González A, Gañán L, Mejía JF, Cardona N, Álvarez E. 2014. Characterization of the *Colletotrichum* species causing anthracnose in andean blackberry in Colombia. *Plant Disease* 98(11): 1503-1513. doi: 10.1094/PDIS-07-13-0752-RE.
- Balouiri M, Sadiki M, Ibsnouda SK. 2016. Methods for *in vitro* evaluating antimicrobial activity: A review. *Journal of Pharmaceutical Analysis* 6(2): 71-79. doi: 10.1016/j.jpha.2015.11.005.
- Camilotti J, Ferarrese L, Bortolucci WC, Gonçalves JE, Takemura OS, Junior RP, Alberton O, Linde GA, Gazim ZC. 2015. Essential oil of parsley and fractions to *in vitro* control of cattle ticks and dengue mosquitoes. *Journal of Medicinal Plants Research* 9(40): 1021-1030. doi: 10.5897/JMPR2015.5941.
- Chatzopoulou PS, Katsiotis ST. 1995. Procedures influencing the yield and the quality of the essential oil from *Juniperus communis* L. berries. *Pharmaceutica Acta Helvetiae* 70(3): 247-253. doi: 10.1016/0031-6865(95)00026-6.
- Damm U, Cannon PF, Woudenberg JHC, Crous PW. 2012. The *Colletotrichum acutatum* species complex. *Studies in Mycology* 73(1): 37-113. doi: 10.3114/sim0010.
- De Almeida RRP, Souto RNP, Bastos CN, da Silva MHL, Maia JGS. 2009. Chemical variation in *Piper aduncum* and biological properties of its dillapiole-rich essential oil. *Chemistry and Biodiversity* 6(9): 1427-1434. doi: 10.1002/cbdv.200800212.
- Di Stefano V, Pitonzo R, Schillaci D. 2011. Antimicrobial and antiproliferative activity of *Athamanta sicula* L. (Apiaceae). *Pharmacognosy Magazine* 7(25): 31-34. doi: 10.4103/0973-1296.75893.
- El-Zaedi H, Calín-Sánchez A, Martínez-Tomé J, Noguera-Artiaga L, Burló F, Carbonell-Barrachina AA. 2016a. Irrigation dose and plant density affect the essential oil content and sensory quality of parsley (*Petroselinum sativum*). *Scientia Horticulturae* 206(1): 1-6. doi: 10.1016/j.scienta.2016.04.028.
- El-Zaedi H, Martínez-Tomé J, Calín-Sánchez A, Burló F, Carbonell-Barrachina AA. 2016b. Volatile composition of essential oils from different aromatic herbs grown in Mediterranean regions of Spain. *Foods* 5(2): 41. doi: 10.3390/foods5020041.
- Farzaei MH, Abbasabadi Z, Ardekani MRS, Rahimi R, Farzaei F. 2013. Parsley: a review of ethnopharmacology, phytochemistry and biological activities. *Journal of Traditional Chinese Medicine* 33(6): 815-826. doi: 10.1016/S0254-6272(14)60018-2.
- Falconí CE, Yáñez-Mendizábal V. 2018. Efficacy of UV-C radiation to reduce seedborne anthracnose (*Colletotrichum acutatum*) from Andean lupin (*Lupinus mutabilis*). *Plant Pathology* 67(4): 831-838. doi: 10.1111/ppa.12793.
- Gao YY, He LF, Li BX, Mu W, Li J, Liu F. 2017. Sensitivity of *Colletotrichum acutatum* to six fungicides and reduction in incidence and severity of chili anthracnose using pyraclostrobin. *Australasian Plant Pathology* 46(6): 521-528. doi: 10.1007/s13313-017-0518-8.

- Gaviria-Hernández V, Patiño-Hoyos LF, Saldarriaga-Cardona A. 2013. Evaluación *in vitro* de fungicidas comerciales para el control de *Colletotrichum* spp., en mora de castilla. *Corpoica Ciencia y Tecnología Agropecuaria* 14(1): 67-75.
- Grande-Tovar CD, Chaves-Lopez C, Serio A, Rossi C, Paparella A. 2018. Chitosan coatings enriched with essential oils: Effects on fungi involved in fruit decay and mechanisms of action. *Trends in Food Science and Technology* 78(1): 61-71. doi: 10.1016/j.tifs.2018.05.019.
- He J, Wu D, Zhang Q, Chen H, Li H, Han Q, Lai X, Wang H, Wu Y, Yuan J, Dong H, Qin W. 2018. Efficacy and mechanism of cinnamon essential oil on inhibition of *Colletotrichum acutatum* isolated from 'Hongyang' kiwifruit. *Frontier in Microbiology* 9(1): 1-12. doi: 10.3389/fmicb.2018.01288.
- Jilková B, Víchová J, Pokorný R, Vejražka K. 2015. Sensitivity of *Colletotrichum acutatum* isolates to selected fungicides. *Acta Universitatis Agriculturae et Silviculturae Mendelianae Brunensis* 63(4): 1111-1119. doi: 10.11118/actaun201563041111.
- Khalil N, Ashour M, Fikry S, Singab AN, Salama O. 2018. Chemical composition and antimicrobial activity of the essential oils of selected Apiaceous fruits. *Future Journal of Pharmaceutical Sciences* 4(1): 88-92. doi: 10.1016/j.fjps.2017.10.004.
- Kurowska A, Galazka I. 2006. Essential oil composition of the parsley seed of cultivars marketed in Poland. *Flavour and Fragrance Journal* 21(1): 143-147. doi: 10.1002/ffj.1548.
- Linde GA, Gazim ZC, Cardoso BK, Jorge LF, Tešević V, Glamčičija J, Sokovic M, Colauto NB. 2016. Antifungal and antibacterial activities of *Petroselinum crispum* essential oil. *Genetics and Molecular Research* 15(3). doi: 10.4238/gmr.15038538.
- Meepagala KM, Sturtz G, Wedge DE, Schrader KK, Duke SO. 2005. Phytotoxic and antifungal compounds from two Apiaceae species, *Lomatium californicum* and *Ligusticum hultenii*, rich sources of Z-ligustilide and apiol, respectively. *Journal of Chemical Ecology* 31(7): 1567-1578. doi: 10.1007/s10886-005-5798-8.
- Morais LAS, Castanha RF. 2012. Chemical composition of sweet basil essential oil naturally submitted to *Planococcus citri* infestation. *Horticultura Brasileira* 30(2): S2178-S2182.
- Moreira-Valente VM, Newandran-Jham G, Marangon-Jardim C, Dev-Dhingra O, Ghiviriga I. 2015. Major antifungals in nutmeg essential oil against *Aspergillus flavus* and *A. ochraceus*. *Journal of Food Research* 4(1): 51-57. doi: 10.5539/jfr.v4n1p51.
- Nazzaro F, Fratianni F, Coppola R, De Feo V. 2017. Essential oils and antifungal activity. *Pharmaceuticals* 10(4): 1-20. doi: 10.3390/ph10040086.
- Numpaque MA, Oviedo LA, Gil JH, García CM, Durango DL. 2011. Thymol and carvacrol: biotransformation and antifungal activity against the plant pathogenic fungi *Colletotrichum acutatum* and *Botryodiplodia theobromae*. *Tropical Plant Pathology* 36(1): 3-13. doi: 10.1590/S1982-56762011000100001.
- Pisoschi AM, Pop A, Georgescu C, Turcus V, Olah NK, Mathe E. 2018. An overview of natural antimicrobials role in food. *European Journal of Medicinal Chemistry* 143(1): 922-935. doi: 10.1016/j.ejmech.2017.11.095.
- Rashid TS, Awla HK, Sijam K. 2018. Antifungal effects of *Rhus coriaria* L. fruit extracts against tomato anthracnose caused by *Colletotrichum acutatum*. *Industrial Crops & Products* 113(1): 391-397. doi: 10.1016/j.indcrop.2018.01.066.
- Razzaghi-Abyaneh M, Yishinari T, Shams-Ghahfarokhi MS, Rezaee MB, Nagasawa H, Sakuda S. 2007. Dillapiol and apiol as specific inhibitors of the biosynthesis of aflatoxin G1 in *Aspergillus parasiticus*. *Bioscience, Biotechnology and Biochemistry* 71(9): 2329-2332. doi: 10.1271/bbb.70264.
- Rivillas-Acevedo LA, Soriano-García M. 2007. Isolation and biochemical characterization of an antifungal peptide from *Amaranthus hypochondriacus* seeds. *Journal of Agricultural and Food Chemistry* 55(25): 10156-10161. doi: 10.1021/jf072069x.
- Romeilah RM, Fayed SA, Mahmoud GI. 2010. Chemical Compositions, Antiviral and Antioxidant Activities of Seven Essential Oils. *Journal of Applied Sciences Research* 6(1): 50-62.
- Semeniuc CA, Pop CR, Rotar AM. 2017. Antibacterial activity and interactions of plant essential oil combinations against Gram-positive and Gram-negative bacteria. *Journal of Food and Drug Analysis* 25(2): 403-408. doi: 10.1016/j.jfda.2016.06.002.
- Stankovic M, Nikolic N, Stanojevic L, Cacic MD. 2004. The effect of hydrodistillation technique on the yield and composition of essential oil from the seed of *Petroselinum crispum* (Mill.) Nym. Ex. A.W. Hill. *Chemistry and Industry* 58(9): 409-412. doi: 10.2298/HEMIND0409409S.
- Tripathi P, Dubey NK. 2004. Exploitation of natural products as an alternative strategy to control postharvest fungal rotting of fruit and vegetables. *Postharvest Biology and Technology* 32(3): 235-245. doi: 10.1016/j.postharvbio.2003.11.005.
- Wharton PS, Diéguez-Urbeondo J. 2004. The biology of *Colletotrichum acutatum*. *Anales del Jardín Botánico de Madrid* 61(1): 3-22. doi: 10.3989/ajbm.2004.v61.i1.61.
- Zhang H, Chen F, Wang X, Yao HY. 2006. Evaluation of antioxidant activity of parsley (*Petroselinum crispum*) essential oil and identification of its antioxidant constituents. *Food Research International* 39(8): 833-839. doi: 10.1016/j.foodres.2006.03.007.
- Zhang RL, Huang JS. 2007. Cloning of a carbendazim resistant gene from *Colletotrichum gloeosporioides* of mango in South China. *African Journal of Biotechnology* 6(2): 143-147.